

ONLINE SEARCH REQUEST FORM

1-120

USER J. Chambers SERIAL NUMBER 08/319,745

ART UNIT 1804 PHONE 308-2035 DATE 1/17/96

Please give a detailed statement of requirements. Describe as specifically as possible the subject matter to be searched. Define any terms that may have special meaning. Give examples or relevant citations, authors, or keywords, if known.

You may include a copy of the broadest and or relevant claim(s).

Search Seq. I. D. 3, 4 and 10 please

FOR OFFICIAL USE ONLY

STAFF USE ONLY

COMPLETED 1-31-96

SEARCHER —

ONLINE TIME 20 TOTAL TIME 20
(in minutes)

NO. OF DATABASES 1

SYSTEMS

☐ CAS ONLINE

☐ DARC/QUESTEL

☐ DIALOG

☐ SDC

☒ OTHER IF

Jan 17 18:01

US-08-319-745-3.rge

1

Jan 17 18:01

US-08-319-745-3.rge

2

WATERBURY (TM)

Release 2.1D John F. Collings, Biocomputing Research Unit.
Copyright (c) 1993, 1994, 1995 University of Edinburgh, U.K.
Distribution rights by IntelliGenetics, Inc.

MParch_nn n.a. - n.a. database search, using Smith-Waterman algorithm

Run on: Wed Jan 17 17:09:59 1996; MasPar time 3098.56 Seconds
1086.012 Million cell updates/sec

Tabular output not generated.

Title: >US-08-319-745-3
Description: (1:4448) from US08319745.seq
Perfect Score: 4448
N.A. Sequence: 1 GCCCAGGTGCGACGCTGT.....CTTTTTTTTTTTTTTTTG 4448
Comp: CGGTCACAGCTGTGCCACA.....GAAAAAAAAAAAAAAC

Scoring table: TABLE default
Gap 6

Nmatch STD : Dbase 0; Query 0

Searched: 478145 seqs, 378268764 bases x 2

Database: emb1-new7

1 BCT
2 FUN
3 INV1
4 INV2
5 MAM
6 ORG
7 PLN
8 PRI
9 PRO
10 ROD
11 SYN
12 UNC
13 VRT
14 VIR

Database:

EST-STS

15 EST1
16 EST2
17 EST3
18 EST4
19 EST5
20 EST6
21 EST7
22 EST8
23 EST9
24 EST10
25 EST11

26 EST12

27 EST13

28 EST14

29 EST15

30 EST16

31 EST17

32 EST18

33 EST19

34 EST20

35 EST21

36 EST22

37 EST23

38 EST24

39 EST25

40 EST26

41 EST27

42 EST28

43 EST29

44 EST30

45 EST31

46 EST32

47 EST33

48 EST34

49 EST35

50 EST36

51 EST37

52 EST38

53 EST39

54 EST40

55 EST41

56 EST42

57 EST43

58 EST44

59 EST45

60 EST46

61 EST47

62 EST48

63 EST49

64 EST50

65 EST51

66 EST52

67 EST53

68 EST54

69 EST55

70 EST56

71 EST57

72 EST58

73 STS1

74 STS2

75 STS3

76 STS4

77 gnEST1

78 gnEST2

79 gnEST3

80 gnEST4

81 gnEST5

82 gnEST6

83 gnEST7

84 gnEST8

85 gnEST9

Database:

genbank89

86 BCT1

87 BCT2

88 BCT3
89 BCT4
90 BCT5
91 BCT6
92 INV1
93 INV2
94 INV3
95 INV4
96 INV5
97 MAM1
98 MAM2
99 PAT1
100 PAT2
101 PHG
102 PLN1
103 PLN2
104 PLN3
105 PLN4
106 PLN5
107 PLN6
108 PLN7
109 PRI1
110 PRI2
111 PRI3
112 PRI4
113 PRI5
114 PRI6
115 PRI7
116 PRI8
117 PRI9
118 ROD1
119 ROD2
120 ROD3
121 ROD4
122 ROD5
123 ROD6
124 ROD7
125 STR
126 SYN
127 UNA
128 VRL1
129 VRL2
130 VRL3
131 VRL4
132 VRL5
133 VRL6
134 VRT1
135 VRT2
136 VRT3

Database:

genbank-new7

137 BCT
138 INV
139 MAM
140 PHG
141 PLN
142 PRI
143 ROD
144 STR
145 SYN
146 UNA
147 VRL
148 VRT

Database: u-emb143.89
149 part1

Statistics: Mean 12.432; Variance 3.452; scale 3.601

Pred. No. is the number of results predicted by chance to have a score greater than or equal to the score of the result being printed, and is derived by analysis of the total score distribution.

SUMMARIES

Result	No.	Score	Query	Match	Length	DB	ID	Description	Pred. No.
	1	203		4.6	5536	93	DMPTCR	D. melanogaster patch	6.73e-241
	2	141		3.2	5665	94	DROMPP2	D. melanogaster membra	7.58e-151
	3	55		1.2	1301	116	HUMRDP	Human RD protein (RD)	3.59e-34
	4	55		1.2	1875	115	HUMMRD4	Human MHC class III H	3.59e-34
	5	55		1.2	1226	111	HSRD	Human mRNA for RD pro	3.59e-34
	6	49		1.1	1805	103	DDIDPP5A	Dictyostelium purpure	5.73e-27
	7	47		1.1	1961	103	DDIDPP4A	Dictyostelium purpure	1.26e-24
	8	46		1.0	1831	136	XLU170KR	Xenopus laevis UI 70K	1.81e-23
	9	44		1.0	3963	93	DMO09306	Drosophila melanogast	3.54e-21
	10	44		1.0	255	57	T40389	ya33d03.r3 Homo sapie	3.54e-21
	11	43		1.0	6800	121	MUSIGMD	Mouse germline IgM ch	4.78e-20
	12	41		0.9	1332	136	XLU170K0	Xenopus laevis UI 70K	8.13e-18
	13	41		0.9	593	76	HOMU1265	Human chromosome 2 ST	8.13e-18
	14	41		0.9	1812	121	MUSMRDOK	Mouse MHC class III R	8.13e-18
	15	40		0.9	243	49	T02581	0224C3 Plasmodium fal	1.02e-16
	16	39		0.9	22703	93	CEZK675	Caenorhabditis elegans	1.25e-15
	17	39		0.9	22703	4	CEZK675	Caenorhabditis elegans	1.25e-15
	18	39		0.9	22703	138	CEZK675	Caenorhabditis elegans	1.25e-15
c	19	38		0.9	21805	4	CEF21H12	Caenorhabditis elegans	1.49e-14
c	20	38		0.9	21805	92	CELF21H12	Caenorhabditis elegans	1.49e-14
	21	38		0.9	486	65	T71726	yc62a01.rl Homo sapie	1.49e-14
	22	37		0.8	1770	103	DDIDPP3A	Dictyostelium purpure	1.72e-13
	23	37		0.8	756	100	A25270	Human IFN-gamma antag	1.72e-13
	24	37		0.8	756	117	S74221	IK-IK factor [human,	1.72e-13
	25	37		0.8	756	8	HS221157	Rattus norvegicus (cl	1.92e-12
c	26	36		0.8	451	123	RATSAT	H. sapiens partial CD	1.92e-12
	27	36		0.8	215	22	HSC25B012	Mus musculus (clone l	2.08e-11
c	28	35		0.8	11026	122	MUSMVDPA	Rat substance P recep	2.08e-11
c	29	35		0.8	3408	123	RATSPR	IB1866 Homo sapiens c	2.08e-11
	30	35		0.8	279	51	T15769	M. musculus DNA for vi	2.08e-11
	31	35		0.8	189	10	MMVIMV31	Human STS UT266, 5' p	2.08e-11
	32	35		0.8	189	143	MMVIMV31	Rat substance P recep	2.08e-11
	33	35		0.8	173	76	HUMUT266A	Ovis aries subspecies	2.17e-10
c	34	35		0.8	2552	123	RATSPR05	Rat gamma-crystallin	2.17e-10
	35	35		0.8	471	98	U00207	Capra aegagrus Saanen	2.17e-10
	36	34		0.8	54670	122	RATCRYG	Mouse Hox-3.1 gene an	2.17e-10
	37	34		0.8	550	98	U00186	Mouse Hox-3.1 gene an	2.17e-10
	38	34		0.8	434	34	RZ0917	Ovis aries Perendale	2.17e-10
c	39	34		0.8	10272	120	MUSHOXMAA	R. norvegicus (Sprague	2.19e-09
	40	34		0.8	479	98	U00206	Chlamydomonas reinhar	2.19e-09
c	41	33		0.7	3551	123	RNAALB	D. melanogaster membra	2.19e-09
	42	33		0.7	8670	103	CRAG7	Mus musculus 129/Sv c	2.19e-09
	43	33		0.7	3376	94	DROMPP1	R. norvegicus N-myc ge	2.19e-09
c	44	33		0.7	6125	119	MMU10098		
	45	33		0.7	6951	124	RNNMYC		

ALIGNMENTS

RESULT 1

Jan 17 18:01

US-08-319-745-3.rge

5

LOCUS DMPCTC 5536 bp RNA INV 09-APR-1991
DEFINITION D. melanogaster patched (ptc) mRNA involved in segment polarity.
ACCESSION X17558
KEYWORDS developmental regulation; glycoprotein; patched gene; segmentation gene; transmembrane protein.
SOURCE fruit fly.
ORGANISM Drosophila melanogaster
Eukaryotae; mitochondrial eukaryotes; Metazoa/Eumycota group;
Metazoa; Eumetazoa; Bilateria; Coelomata; Protostomia; Arthropoda;
Tracheata; Insecta; Pterygota; Neoptera; Holometabola; Diptera;
Brachycera; Cyclorrhapha; Drosophilidae; Drosophila; Sophophora;
melanogaster group; melanogaster subgroup.
REFERENCE 1 (bases 1 to 5536)
Ingham, P.W.
AUTHORS Direct Submission
TITLE Submitted (10-DEC-1990) to the EMBL/GenBank/DBJ databases. Ingham
JOURNAL P.W., Imperial Cancer Research Fund Developmental Biology Unit,
JOURNAL Nature 341 (6242), 508-513 (1989)
MEDLINE 90015164
COMMENT See also M28418.
See also M28418 for related sequence.

FEATURES
source

mRNA

CDS

NCBI gi: 8389 Location/Qualifiers
1..5536
/organism="Drosophila melanogaster"
/dev_stage="embryo"
/chromosome="2"
/map="440"
1..5536
/gene="ptc"
752..4651
/gene="ptc"
/standard_name="patched"
/note="NCBI gi: 8390"
/codon_start=1
/product="patched protein"
/translation="MDRSLRPVDPDTHGVVDKLFSDLYIRTSWVDAQVALDQIDKG
KARGSRATYILRSVFSQSHLETLGSSVQKHAGVLFVALVILSTFCVGLKSAQIHKSVH
QWIOEGGGLAEALAYTKTIGEDSATHQLLIQTHDPNLSVLPQALLAHLELVK
ATAVRHLDTMGLRDMCNPSFSEFIEQILRHLPSCIIITPDLDFEGSQL
LGPFAVPIPNQRLMTTINPASMVQMKMSSEKISDFETVEQYMKRAATASG
YMEKPCINPNCPDTPAKNSATOPDVCAILSGCYAANKHWPPEELIVGAKR
NHSGLHRAAALQSVQVLTQEMEDQDNKYVHLGWTQKAAEVLNMQNRSRE
VEQLRQSRATNYDIYVSSAALDDI LAKFSHP SALSIVIGVATVLYAFCTILRW
RDPVSGSVGAVGLMCFSTAAAGLIGLALLGVFNAASTQVVPFLAIGLVDFIEM
LTAAYESNRREQTKLILKKVPSILFSACSTAGSFFMAAFIPVPAIKVFCLOALIVM
CSNLAALVILPAMISDLRRRTAGRADIFCCFPVMEQPKVAPVLPVLPNNNGRGA
RHPSKNNNRVAPLPAQNPLEORADIPGSSHSLSFSLATFAFOHYTPFLMRISWKEFL
TVWGF LAALISLSVASTRLDGLDIIIDLPKSNHEKFLDQATRLFGFSYNAVTQGN
FEYPTQQLRDYHDVSVVRVPIKNDNGGGLDFWLLIFSLGNLQKIFDEEYRDR
LTKECWEPNASSDAI LAYKLI VQTGHVNDPDKELVLTNRLVNSDGLINORAEYNYLS
RHPSKNNNRVAPLPAQNPLEORADIPGSSHSLSFSLATFAFOHYTPFLMRISWKEFL
MFLYHLGLDTISQILTIHIDLSVYEGFLNPSPGIPFIFMEQYTLRSLAMI
LACVILAAVILSVLLISVAAVILSVLASIAOIFCMTLLIGLISAPVILILS
VGMMLCNVLISLGEMTSVGNRQVRQLSNQMSLGLVHGMILTSGVAFHLSISPEF
VIRHFCWLLVLCVGCANSLIVFPILLSMVGEAELVP LEHPDRISTPSPFLPVRSSK
RSGKSVYQGSRSRSGSCQKSHHHHKKDINDPSLTTITEEPQSWKSSNSSIQPNWDIT

Jan 17 18:01

US-08-319-745-3.rge

6

mat_peptide
752..4648
/gene="ptc"
/standard_name="patched"
/note="signal peptide sequence was not determined"
/product="patched protein"
misc_feature
4896..4900
/gene="ptc"
/note="mRNA destabilizing motif"
polyA_signal
4900..4905
/gene="ptc"
misc_feature
5004..5008
/note="atypical"
misc_feature
5140..5144
/note="mRNA destabilizing motif"
misc_feature
5246..5250
/note="mRNA destabilizing motif"
misc_feature
5456..5460
/note="mRNA destabilizing motif"
polyA_signal
5503..5508
/gene="ptc"
polyA_signal
5503..5508
/gene="ptc"
polyA_signal
5536
/gene="ptc"
BASE COUNT 1360 a 1575 c 1407 g 1194 t
ORIGIN

DB 93; Score 203; Match 59.9%; OryMatch 4.6%; Pred. No. 6.73e-241;
Matches 885; Conservative 0; Mismatches 574; Indels 18; Gaps 14;
Db 821 gatctttacataccagcagctgggtggagcccaagtgccgcgcagatagataag 880
||||| ||||| ||||| ||||| ||||| ||||| ||||| ||||| ||||| ||||| |||||
Qy 300 CATCTTTACATAGCTACAGTTGGTGGACGCCGCTAGCTCTCTGAACTCGAAAG 359
||||| ||||| ||||| ||||| ||||| ||||| ||||| ||||| ||||| |||||
Db 881 ggcaagcgctggcgagcgagcgatctctgcgcatcagtcattccagtcaccactc 940
||||| ||||| ||||| ||||| ||||| ||||| ||||| ||||| ||||| |||||
Qy 360 GGTAACTCGAAGGAGGAAGAACCTCTCTGTGGATACGAGCGTGGCTACAAAGACGCTC 419
||||| ||||| ||||| ||||| ||||| ||||| ||||| ||||| ||||| |||||
Db 941 gaaacctcgagcagctccggtgcaaaagcagcgagcgagctgctattcgtgctatcctg 1000
||||| ||||| ||||| ||||| ||||| ||||| ||||| ||||| ||||| |||||
Qy 420 TTTATTTGGCTGCTTTCTTCAAGCGACGCGGGGAAAGTCTCTTCGTTGCCATCCTC 479
||||| ||||| ||||| ||||| ||||| ||||| ||||| ||||| ||||| |||||
Db 1001 gtctgagcacttcctgctgcgctgaagagcgccagatccactccaagtgcaaccag 1060
||||| ||||| ||||| ||||| ||||| ||||| ||||| ||||| ||||| |||||
Qy 480 GTTCTCTGAGCTTCTGCGTGGTCTCAAGTCACGCAAAATACATACAAGGTCGACCAA 539
||||| ||||| ||||| ||||| ||||| ||||| ||||| ||||| ||||| |||||
Db 1061 ctgtgagcagagcgagcgagcgagcgagcgagcgagcgagcgagcgagcgagcgagcgagc 1120
||||| ||||| ||||| ||||| ||||| ||||| ||||| ||||| ||||| |||||
Qy 540 CTCTGGGTTCAAGAGGCTGTGATAGAGCCGAGTTCATACAAAGTCCAGGACGCTTTG 599
||||| ||||| ||||| ||||| ||||| ||||| ||||| ||||| ||||| |||||
Db 1121 ggcagagcagagtcggcgagcagcagtcagtcgctattcagagcagcagcagcagcagc 1180
||||| ||||| ||||| ||||| ||||| ||||| ||||| ||||| ||||| |||||
Qy 600 GGGAGGCGGACTCCTCGACGCGACGAGCTGTCTATACAAAGTCCAGGACGCTC 659
||||| ||||| ||||| ||||| ||||| ||||| ||||| ||||| ||||| |||||
Db 1181 tcgctcgcagcagcagcagcagcagcagcagcagcagcagcagcagcagcagcagcagc 1240
||||| ||||| ||||| ||||| ||||| ||||| ||||| ||||| ||||| |||||
Qy 660 TCCTCTGCTACATCCAGGCGCTTCTTGAACACCTTGAAGTGGTGCACGCGGACGCTCGG 719
||||| ||||| ||||| ||||| ||||| ||||| ||||| ||||| ||||| |||||

REFERENCE 1 (bases 1 to 1276)
AUTHORS Spritz, R. A.
TITLE Direct Submission
JOURNAL Submitted (16-AUG-1989) to the EMBL/GenBank/DBJ databases. Spritz R.A., University of Wisconsin-Madison, 309 Laboratory of Genetics, 445 Henry Mall. Madison WI 53706. U S A

AUTHORS	Surovy, C.S., Hoganson, G., Gosink, J., Strunk, K. and Spritz, R.A.
TITLE	The human RD protein is closely related to nuclear RNA-binding proteins and has been highly conserved
JOURNAL	Gene 90 (2), 299-302 (1990)
MEDLINE	90387680
COMMENT	A second cDNA sequence (clone hRD2 from lglt11 placental library) corresponds to bases 1-1082 and is identical.

[illegible]


```
exon          583..684
              /number=8
exon          685..772
              /number=9
exon          773..1831
              /number=10
polyA signal  1810..1817
polyA site    1831
BASE COUNT   602 a 335 c 524 g 370 t
ORIGIN
DB 136; Score 46; Match 65.3%; QryMatch 1.0%; Pred. No. 1.81e-23;
Matches 98; Conservative 0; Mismatches 52; Indels 0; Gaps 0;
Db 1228 gaagcgacccatagacagagtagagagagagataggacagagacagagacagga 1287
      ||||| |||| ||||| ||||| ||||| ||||| ||||| ||||| ||||| |||||
Qy 3946 GAAGGATCGCATGACATAGGATCGAGACCGCTGMAAGGCACAGATCGGACAGGC 4005
Db 1288 ggcgagataggaccgcatagagagagagagagatcacaaacgagagcgagacagag 1347
      || ||||| |||| ||||| |||| ||||| |||| ||||| |||| ||||| ||||
Qy 4006 ATCGGATAGGATCCTGACCGGACAGGATAGGATAGGATAGGATAGGATAGGATAG 4065
Db 1348 gtgacgcgagcaaaagagagagagaggg 1377
      || ||||| |||| ||||| |||| ||||| |||| ||||| |||| ||||| ||||
Qy 4066 ACAGCGGAGACCGATATAGACGMAAGGG 4095

RESULT 9
LOCUS Dm009306 3963 bp mRNA INV 15-MAY-1994
DEFINITION Drosophila melanogaster shuttle craft protein (etc) mRNA, complete cds.
ACCESSION U09306
KEYWORDS fruit fly.
SOURCE Drosophila melanogaster
ORGANISM Drosophila melanogaster
          Eucaryotae; Metazoa; Arthropoda; Tracheata; Insecta; Pterygota;
          Diptera; Brachycera; Cyclorrhapha; Drosophilidae; Drosophila;
          (Sophophora); melanogaster group; melanogaster subgroup.
REFERENCE 1 (bases 1 to 3963)
AUTHORS Strumbakis, N.D., Li, Z. and Tolias, P.P.
TITLE Dynamic subcellular distribution in egg chambers and embryos of a novel Drosophila melanogaster protein encoded by the embryonic lethal gene shuttle craft
JOURNAL Unpublished
REFERENCE 2 (bases 1 to 3963)
AUTHORS Tolias, P.P.
TITLE Direct Submission
JOURNAL Submitted (29-APR-1994) Peter P. Tolias, Public Health Research Institute, 455 First Ave., New York, NY 10016, USA
COMMENT NCBI gi: 487399
FEATURES             Location/Qualifiers
     source           1..3963
                     /clone_lib="ovarian gt22A of P. Tolias, ovarian gtl1 of
                     L. Kalfayan and st.10 of A. Spradling"
                     /chromosomes="2L"
                     /organism="Drosophila melanogaster"
                     /map="35C1-C3"
                     /sex="female"
                     /tissue_type="ovary"
                     /dev_stage="adult"
     note="Several overlapping clones used to assemble
sequence of shuttle craft/1(2)35Cb. Originally isolated as
a partial clone (containing the C-terminal 222 amino
acids) in expression cloning screens for cDNAs encoding
```

```
single stranded nucleic acid binding proteins"
1..223
/map="35C1-C3"
CDS
/feature="etc"
/standard_name="shuttle craft"
/note="C-terminal 222 amino acids encode a novel
single-stranded DNA binding domain; NCBI gi: 487400"
/codon_start=1
/evidence=experimental
/product="shuttle craft protein"
/translation="MAEYWOQLTNGPGGAGPGNGESSAMVDQNGGHESAAGVGCNRRHS
NNYVNFNQIFQHNILGGGAPSNATSTQHPVGSSTYTFSLGGGGAGGAGIAPPVASAS
TSFANVSHQSPNYSQAMPTTONGDGIARVTVTSSYGSNSNSFSSYTFPGNN
PDFSASKLOASAPFEVNFPAKLSLEETPAATTCNGSTASLETAINETRPTLRAGE
PAERCANQCSNNINREERERDRDRDRDRDRDRDRDRDRDRDRDRDRDRDRDRDRDRDR
QRSDYRDREDYERSDRARPQOQDYDNHRNKRDRDRDRDRDRDRDRDRDRDRDRDRDRDR
TSNSAHPSPKSOLOQI1SPRGPPLPPADNEKLSQREKLVDRDIEQRRLCGLVCVEA
IKSHQPTWSCRCNYHMLHLKCTITWASSKSEVGRCPACQNVLDLPRYLCLFCCKL
KNPPVSRTELAHSGEGCCRIEGCSHACTLILCHPGPCPPQANVVRSCGGRSTKTMQ
CAMKEVLGCEICDKLLACGEHRCOAEGSKCAACSEQVWQCHGCKQERKVPCTRE
SQDKTYSCDKSCGP1PCGHHKCKDSCHAGSCPCCKLSPEQITSCPCCKIPVAGOR
SSCLDPIPTCEGICRTLRGCGPAHPHQGSKCHLGCPQPCQKQTVGKCRGCHMDQMI
KCRQLGNRADDARCKRCKTCKRCKGKCNVECCIDIDHDGPIPCNRTLSCKGKCDQ
PCHRGNCPPCYSSFEELYCECGAEVYVPVPGCTKKPICKLPSSRIHPCHDPQHNIC
PQSAGEICRQSCTRPTPCGHRCAACHGACPEPPCKELVEVQCECGNRKRQSRQ
HSGTTPCPMIFTYTKLGHNDWRKTIPCSQPNFSCMGACGKPLPCGHHKCKIKFCHGG
ELAREHSRIATIQLASMAEMSRGNMELSELLAPAKSKNTLDCNDECRLLERNRL
AALSSGNSDTKOKCLTKYSEFVRGAKNPALTYSVETLTLDVLKAKESQKRSRH
SPTPMREKRLVHELCEVFGESVYDKENRNNVATAHDKRCWFATPSIMEVLAHE
SQGRVPVPPSNNAWGLKK"
variation
replace(548..568,"")
/feature="etc"
/note="21 bp from 548..568 are not present in some cDNAs
due to alternative splicing. This results in an in-frame 7
amino acid deletion of the etc protein."
923..1027
repeat_region
/feature="etc"
/note="RD-domain; a repeat region of R followed by D or E"
1109..1159
/feature="etc"
/note="Bipartite nuclear localization site (putative)"
1967..2005
repeat_region
/feature="etc"
/note="PCC repeat (PXXXXXXXXXXCXXC)"
2699..2737
repeat_region
/feature="etc"
/note="PCC repeat (PXXXXXXXXXXCXXC)"
2744..2782
repeat_region
/feature="etc"
/note="PCC repeat (PXXXXXXXXXXCXXC)"
2792..2830
repeat_region
/feature="etc"
/note="PCC repeat (PXXXXXXXXXXCXXC)"
2837..2875
repeat_region
/feature="etc"
/note="PCC repeat (PXXXXXXXXXXCXXC)"
3545..3963
3'UTR
/map="35C1-C3"
polyA signal 3937..3942
polyA site 3963
/map="35C1-C3"
BASE COUNT 1043 a 1055 c 1111 g 754 t
```


polyA_signal	163..168	/note="for Cmu(s)"	
polyA_signal	676..681	/note="putative"	
repeat_region	1097..2096	/note="(CA)33 repeat"	
exon	1863..1978	/rpt_unit=1097..1098	
CDS	join(1863..1978,2097..2105)		
	/partial		
	/note="C-mu-membrane; NCBI gi: 202417"		
	/codon_start=3		
	/product="immunoglobulin mu-chain"		
intron	1979..2096	/translation="GEVNAEEGFENLWTASTIVLFLLSLFTVTTVLFKKV"	
exon	2097..2105	/note="C-mu-membrane intron A"	
polyA_signal	2348..2353	/note="Ig H-chain C-mu-membrane, exon m2"	
repeat_region	2556..2627	/note="for Cmu (membrane); putative"	
repeat_region	2636..2691	/note="(GGGAGA)12-repeat"	
polyA_signal	3367..3372	/rpt_unit=2556..2561	
promoter	3479..3901	/note="(CA)28-repeat"	
CDS	3367..3372	/rpt_unit=2636..2638	
	/note="for Cmu (membrane); putative"		
	3367..3372		
	/note="TATA-like motive for ORF 146; putative"		
	3479..3901		
	/note="ORF 146 with stretch of 27 hydrophobic amino acids, membrane binding segment; putative; NCBI gi: 457144"		
	/codon_start=1		
	/translations="MGLVTKFQIYCHSPYGVTSHTSYFCYCPMLGQSAATHGSLSS		
	RLTFPPSRGKQCGQALPCQPTTLGMCCKFKLVLIWLSQSEATAVGSLSPGTNI		
	PVTVCPLYMPCFKNPVLGLMPTLIQHKFISTRHPP"		
repeat_unit	3509..3680		
repeat_unit	3936..4098	/note="unique sequence inverted repeat 1"	
CDS	join(4653..4934,5311..5415,6415..6735)	/note="unique sequence inverted repeat 1"	
	/partial		
	/note="C-delta, CH1 domain; NCBI gi: 554163"		
	/codon_start=3		
	/product="immunoglobulin mu-chain"		
	/translations="DKKPDPMFLISECKAPENEKINLGLVIGSQPLKISWEPKSS		
	IYEVFPSEMRNGYTWLQYTVLASELNIAHTKINKPKRKPKEPESWDSSQSK		
	RVPTLQAKNHSTEATKAITTKQDIEGAWASNLINILFTSTHPENSWLCEVSGF		
	FPENIHLMLGLVSHKRSKSTFEVTAHPNTAQPGGTQTQMSVLRPLVALSSLDITTCVWVE		
	HEASKTKLNASKSLAIS"		
exon	4653..4934		
CDS	4653..4934		
	/pseudo		
	/codon_start=1		
polyA_signal	4895..4900	/note="for ORF 146; putative"	
intron	4935..5310		
exon	5311..5415	/note="C-delta intron A"	
	/partial		
intron	5416..6414	/note="Ig H-chain C-delta, hinge domain"	

repeat_region	/note="C-delta intron B"
5851..5910	
/note=" (CT) 30-repeat"	
/rpt_unit=5851..5852	
5911..5970	
/note=" (CA) 30-repeat"	
5989..6325	
/note="C gamma 3; putative"	
/pseudo	
/codon_start=1	
6415..6735	
/note="Ig H-chain C-delta, CH3 domain"	
6736..6800	
/note="C-delta intron C"	
BASE COUNT 1953 a 1758 c 1500 g 1589 t	
ORIGIN 387 bp upstream of HindIII site on chromosome 12.	
DB 121; Score 43; Match 59.0%; QryMatch 1.0%; Pred. No. 4.78e-20; Matches 87; Conservative 0; Mismatches 38; Indels 1; Gaps 1;	
Db 2558 gagagggagagagagggagagggagagggagagggagagggagagggagagggaggg 2617	
Qy 3963 GATAGGATCGAGACCGTGAAGGGACAGACATCGGACAGGATCGGATCGGATCGT 4022	
Db 2618 gagagggagagggaggga 2677	
Qy 4023 GACCGGACAGGATAGGATAGAGACCATCGAGAGACAGACAGACAG-CCGAGACCCATA 4081	
Db 2678 gagaga 2683	
Qy 4082 TAGAGA 4087	
RESULT 12	
LOCUS X1D170K0 1332 bp DNA	VRT 05-APR-1990
DEFINITION Xenopus laevis U1 70K gene exon 10.	
ACCESSION X12429	
KEYWORDS U1 small nuclear RNA.	
SOURCE African clawed frog.	
ORGANISM Xenopus laevis	
Eukaryotes; mitochondrial eukaryotes; Metazoa/Eumycota group; Metazoa; Eumetazoa; Bilateria; Coelomata; Deuterostomia; Chordata; Vertebrata; Gnathostomata; Osteichthyes; Sarcopterygii; Chondata; Tetrapoda; Amphibia; Anura; Mesobatrachia; Pipidae; Pipidae; Xenopodinae; Xenopus.	
1 (bases 1 to 1332)	
Etterodt,M., Vignali,R., Scherly,D., Mattaj,I.W., Ciliberto,G. and Philipson,L.	
Title Direct Submission	
Journal Submitted (18-JUL-1988) to the EMBL/GenBank/DBJ databases.	
Etterodt M., University of Aarhus, Dept. of Molecular Biology, C.F. Moellers Allee 130, DK-8000 Aarhus C, Denmark	
2 (bases 1 to 1332)	
Etterodt,M., Vignali,R., Ciliberto,G., Scherly,D., Mattaj,I.W. and Philipson,L.	
Title Structure and expression of a Xenopus gene encoding an snRNP protein (U1 70K)	
Journal EMBO J. 7 (13), 4311-4321 (1988)	
Medline 89210819	
Comment NCBI gi: 65172	
Features Location/Qualifiers	
source 1..1332	
/organism="Xenopus laevis"	
/clone_lib="lambda EMBL4"	

WATERMAN

(TM)

Release 2.1D John F. Collins, Biocomputing Research Unit.
Copyright (c) 1993, 1994, 1995 University of Edinburgh, U.K.
Distribution rights by IntelliGenetics, Inc.

MParch_pp protein - protein database search, using Smith-Waterman algorithm

Run on: Wed Jan 17 17:22:34 1996; MasPar time 17.16 Seconds
502.247 Million cell updates/sec

Tabular output not generated.

Title: >US-08-319-745-10
Description: (1:1356) from US08319745.pep
Perfect Score: 9913
Sequence: 1 MASAGNARRGPGQARRREA.....TORPPWALCPATSPSPL 1356

Scoring table: PAM 150
Gap 11

Searched: 53402 seqs, 6354270 residues

Database: a-geneseq18
1 part1
2 part2
3 part3
4 part4
5 part5
6 part6
7 part7
8 part8
9 part9
10 part10

Statistics: Mean 40.815; Variance 192.679; scale 0.212

Pred. No. is the number of results predicted by chance to have a
score greater than or equal to the score of the result being printed,
and is derived by analysis of the total score distribution.

SUMMARTIES

Result No.	Score	Query Match %	Length	ID	Description	Pred. No.
1	115	1.2	439	5 R28150	Sugar beet chitinase	4.21e+00
2	114	1.2	1229	10 R54074	CryET5.	4.88e+00
3	114	1.2	928	1 R06289	Predicted retinoblast	4.88e+00
4	111	1.1	1280	2 P70452	Sequence encoded by h	7.60e+00
5	111	1.1	1280	8 R44297	Sequence encoded by h	7.60e+00
6	110	1.1	1280	1 R04868	Protein encoded by Mu	8.80e+00
7	109	1.1	521	8 R47068	Mammalian chromaffin	1.07e+01
8	108	1.1	487	4 R22380	Antigen mc-35c.	1.18e+01

9	108	1.1	195	8 R47339	Peptide fragment of m	1.18e+01
10	108	1.1	380	8 R48063	Sequence of polypepti	1.18e+01
11	106	1.1	928	1 R05305	Cancer suppressing gen	1.57e+01
12	106	1.1	286	1 P82590	Polypeptide with glyc	1.57e+01
13	106	1.1	816	1 P82112	Human retinoblastoma	1.57e+01
14	106	1.1	928	7 R36534	Retinoblastoma (RB) p	1.57e+01
15	106	1.1	970	1 P90599	Human retinoblastoma.	1.57e+01
16	104	1.0	272	1 P93560	Plasmodium berghei ci	2.09e+01
17	103	1.0	380	3 R14857	Cry1X protein.	2.41e+01
18	101	1.0	458	10 R54834	Human derived adrenal	3.20e+01
19	101	1.0	253	2 R10535	Prod. of pMG3C9 used	3.20e+01
20	101	1.0	864	4 R24042	Lipoxigenase.	3.20e+01
21	100	1.0	719	2 R08041	81 kD endotoxin deduc	3.68e+01
22	99	1.0	402	2 P70644	Pseudorabies virus gp	4.24e+01
23	98	1.0	575	10 R57139	Interleukin-10 recept	4.87e+01
24	98	1.0	504	10 R54682	Mouse brain 5HT2C ser	4.87e+01
25	98	1.0	1253	5 R28337	SFV4 structural poly	4.87e+01
26	98	1.0	685	2 R11331	Human luteinising hor	4.87e+01
27	97	1.0	316	3 R15878	Cholesterol esterase	5.59e+01
28	97	1.0	316	3 R15868	Cholesterol esterase	5.59e+01
29	97	1.0	316	3 R15873	Cholesterol esterase	5.59e+01
30	97	1.0	657	5 R28964	Notch hN5k full lengt	5.59e+01
31	97	1.0	316	3 R15866	Cholesterol esterase	5.59e+01
32	97	1.0	316	3 R15881	Cholesterol esterase	5.59e+01
33	97	1.0	316	3 R15870	Cholesterol esterase	5.59e+01
34	97	1.0	316	3 R15880	Cholesterol esterase	5.59e+01
35	97	1.0	316	3 R15883	Cholesterol esterase	5.59e+01
36	97	1.0	316	3 R15877	Cholesterol esterase	5.59e+01
37	97	1.0	316	3 R15869	Cholesterol esterase	5.59e+01
38	97	1.0	316	3 R15874	Cholesterol esterase	5.59e+01
39	97	1.0	316	3 R15865	Cholesterol esterase	5.59e+01
40	97	1.0	316	3 R15867	Cholesterol esterase	5.59e+01
41	97	1.0	316	3 R15882	Cholesterol esterase	5.59e+01
42	97	1.0	316	3 R15879	Cholesterol esterase	5.59e+01
43	97	1.0	316	3 R15872	Cholesterol esterase	5.59e+01
44	97	1.0	316	3 R15675	Cholesterol esterase	5.59e+01
45	97	1.0	457	3 P50361	Human acetyl choline	5.59e+01

ALIGNMENTS

RESULT 1
ID R28150 standard; Protein; 439 AA.
AC R28150;
DT 17-MAR-1993 (first entry)
DE Sugar beet chitinase 1.
KW SBC-1; fungicide; anti-fungal agent; extensine.
OS Beta vulgaris cv monova.
FH Key Location/Qualifiers
FT Peptide 1..26
FT /label= leader
FT Domain 27..46
FT /label= hevein domain
FT Domain 47..178
FT /label= proline rich
FT /note= "possibly involved in anchoring
chitinase 1 protein to the cell wall
after modification of the prolines to
FT glycosylated hydroxyprolines, as in
FT extensines"
FT Domain 179..416
FT /label= functional domain
FT Region 417..439
FT /note= "probably directs the protein to

PR	28-MAR-1986; US-845610.
PR	01-AUG-1986; US-892575.
PA	(UNII) UNIV OF ILLINOIS.
PI	Roninson IB, Pastan IH, Gottesman MM;
DR	WPI; 87-291656/41.
DR	N-PSDB; N70752.
PT	DNA for multi-drug resistance in human cells - used to detect
PT	chemotherapy-resistant tumour cells and for producing
PT	polypeptide(s) for diagnosis and therapy
PS	Claim 4(a); Table 5, pp30-39; 61pp; English.
CC	The human multi-drug resistant KB carcinoma cell lines were used as
CC	the source of the mdrl gene nucleic acid sequences (N70751). To
CC	obtain cDNA clones of the mdrl gene (N70752), poly (A) and RNA was
CC	used. Analysis of the AA sequence presented in (N70752) indicates
CC	that the mdrl gene product is likely to be a transmembrane protein.
CC	The presence of transmembrane domains and potential glycosylation
CC	sites is consistent with the mdrl protein being related to the
CC	p-glycoprotein.
SQ	Sequence 1280 AA;
DB	2; Score 111; Match 22.8%; QryMatch 1.1%; Pred. No. 7.60e+00;
Matches	29; Conservative 43; Mismatches 46; Indels 9; Gaps 8;
Db	216 lvlaispvlglsaaawakilsftskellayagavaveevlaairtviafggqkkele 275 : ::: : : : : : : : : : : : : :
Qy	462 LVALSVAAGLIGCSLI-GISFNA-ATTQVLPFLIAGVGVDVFLLAHA-FSETGQNKRI- 517 :: : : : : : : : : : : : : : : : : :
Db	276 rynklee-akrigikkaitanisgaafllliyasy-al-afwygttlvlageysigqvl 332 :: : : : : : : : : : : : : : : : : :
Qy	518 PFEDRTGCILKRTCASVALTSISNWTAFFMAALIPALRAFSLQAQAVVV--FNFAVL 575 333 tvffsvl 339 : : : :
Qy	576 LIFPAIL 582
RESULT	5
ID	R44297 standard; Protein; 1280 AA.
AC	R44297;
DT	24-JUN-1994 (first entry)
DE	Sequence encoded by human multi-drug resistant gene mdrl.
KW	p-glycoprotein; multi-drug resistant gene; mdrl.
OS	Homo sapiens.
PN	W09324613-A.
PD	09-DEC-1993.
PF	14-MAY-1993; U04707.
PR	22-MAY-1992; US-887712.
PA	(GENE-) GENETIC THERAPY INC.
PA	(USSH) US DEPT HEALTH & HUMAN SERVICES.
PI	McDonagh KT, Nienhuis A, Tolstoshev P;
DR	WPI; 93-405805/50.
DR	N-PSDB; O52726.
PT	DNA or RNA sequence for human multi-drug resistant gene MDRL -
PT	encodes P-glycoprotein with altered splice site, used in cancer
PT	therapy
PS	Example; Fig 4; 64pp; English.
CC	pMDR2000 contains an mdrl cDNA sequence (O52726) described in PCT
CC	application no. W087/0943m wherein the first 282 bp of the 5' UTR
CC	and the last 23 bp of the 3' UTR of the cDNA sequence have been
CC	removed.
SQ	Sequence 1280 AA;
DB	8; Score 111; Match 22.8%; QryMatch 1.1%; Pred. No. 7.60e+00;
Matches	29; Conservative 43; Mismatches 46; Indels 9; Gaps 8;

Db	216	lvlaaispyldgsaaawakillssftdkellayakagavaeevlaairtviafggqkkele	275		
		:::: : : : :: : :::: :::: ::: ::: ::: :::			
Qy	462	LVALSVAGLIGLSLI-GISFNA-ATTQVLFPLALGVGDDVFLLAHA-FSETGNKRIP-	517		
Db	276	rynkleek-akrigikkaitanisigaafillyasy-al-afwygttlvlsgeysigqvlt	332		
		:: :: :: :: :: :: :: :: :: :: :: ::			
Qy	518	PFEDRTGECKIRTGASVALTSISNVTAFFFMAALIPIPALRAFSIQAAVVVV--FNFAWVL	575		
Db	333	tvffsvl	339		
		: ::			
Qy	576	LIFPAIL	582		
<hr/>					
RESULT 6					
ID	R04868	standard; protein; 1280 AA.			
AC	R47068;				
DT	01-OCT-1990	(first entry)			
DE	Protein encoded by Multidrug Resistance A1 gene				
KW	Multidrug Resistance A1 gene;	drug resistance of cancer cells; sa.			
OS	Homo sapiens.				
PN	J02100680-A.				
PD	12-APR-1990.				
PF	05-OCT-1988;	251475.			
PR	05-OCT-1988;	JO-251475.			
PA	(SUNR) Suntory Ltd.				
DR	WPI; 90-159707/21.				
DP	N-PSDB; Q04522.				
PT	MDR related gene derived from human normal cells -				
PT	codes specific amino acid sequence, used for diagnosing drug				
PT	resistance of cancer cells				
PS	Disclosure; P; Japanese.				
CC	The gene that encodes this protein is useful for diagnosis of drug				
CC	resistance of cancer cells.				
SC	Sequence 1280 AA;				
<hr/>					
DB	1;	Score 110; Match 23.8%; QryMatch 1.1%; Pred. No. 8.80e+00;			
Matches	30; Conservative 43; Mismatches 46; Indels 7; Gaps 6;				
Db	216	lvlaaispyldgsaaawakillssftdeellayakagavaeevlaairtviafggqkkele	275		
		:::: : : : :: : :::: :::: ::: ::: ::: :::			
Qy	462	LVALSVAGLIGLSLI-GISFNAATTQ-VLPFLALGVGDDVFLLAHA-FSETGNKRIP	518		
Db	276	rynkleekakrigikkaitanisigaafillyasy-al-afwygttlvlsgeysigqvlt	333		
		:: :: :: :: :: :: :: :: :: :: ::			
Qy	519	FEDRTGECKIRTGASVALTSISNVTAFFFMAALIPIPALRAFSIQAAVVVV--FNFAWVL	576		
Db	334	vffsvl	339		
		: ::			
Qy	577	LIFPAIL	582		
<hr/>					
RESULT 7					
ID	R47068	standard; Protein; 521 AA.			
AC	R47068;				
DT	01-JUL-1994	(first entry)			
DE	Mammalian chromaffin granule amine transporter protein.				
KW	Vesicle membrane transport protein; gene therapy; screening;				
KW	Parkinsons disease; neurotoxin; identification; detection;				
KW	antibody; probe;chromaffin granule amine transporter protein.				
OS	Rattus rattus.				
FH	Key	Location/Qualifiers			
FT	Domain	22..42			
FT	/label= Transmembrane domain.				

RESULT	10	
ID	R48063 standard; Protein; 380 AA.	
AC	R48063;	
DT	20-JUL-1994 (first entry)	
DE	Sequence of polypeptide encoded by the first open reading frame in	
DE	the unique short (Us) region of bovine herpes virus (BHV) genome	
KW	Insertion region; unique short region; Us; vaccine; antigen.	
OS	Bovine herpes virus type 1, strain ST.	
PN	W09400586-A.	
PD	06-JAN-1994.	
PF	25-JUN-1993; F00642.	
PR	26-JUN-1992; FR-007930.	
PI	(INMR) RHONE MERIEUX SA.	
PA	Audonnet JF, Legastelois ICMA, Leung-tack P, Riviere MEA;	
DR	WPI; 94-036222/03.	
DR	N-PSDB; Q33350.	
PT	New insertion region sequence of bovine herpes virus genomic DNA	
PT	- used for recombinant virus with this region deleted or	
PT	inactivated, useful in vaccines allowing differentiation between	
PT	vaccinated and infected cattle	
PS	Claim 9; Fig 2; 47pp; French.	
CC	The 4190 bp sequence in Q35350 encodes polypeptides homologous to	
CC	HSV-1 gI, gE and US9; these are neither essential for in vitro	

```

DB 1; Score 106; Match 21.3%; QryMatch 1.1%; Pred. No. 1.57e+01;
Matches 30; Conservative 43; Mismatches 56; Indels 12; Gaps 11;

Db 669 lleshpelehiwt-1fghtlqn-eyelmrdrhldqimmsygyckvknidlkfkiiiv- 725
      : : | : | : | : | : | : | : | : | : | : | : | : | : | : | : |
Qy 819 MLEENKQLPQM-WLHYFRDWLGGIQAEDSDWETGRIMP--NNYKNGSDDGV-LAYKLLVQ 875

Db 726 laykdlphavqetkfrvlkeeydsiiivfysfm-qrlktnllqvastrrptlesiph 784
      : : | : | : | : | : | : | : | : | : | : | : | : | : | : | : |
Qy 876 TGSBRKPTDISQLTKORLVDADGIINPSAFY--YLTVAWSNDPVAVAASQANIRPHRPE 933

Db 785 -ip-rspsykfssplripqgn 803
      : : : | : | : | : | : | : | : | : | : | : | : | : | : | : |
Qy 934 WVHKADYN-PETLRIPAAE 953

RESULT 12
ID P82590 standard; protein; 286 AA.
AC P82590;

```

DT 05-NOV-1990 (first entry)
 DE Polypeptide with glycosylation inhibiting factor activity.
 KW Glycosylation; G1F; TGIF4; IgE; immunoglobulin; antiinflammatory agent.
 PN EP-285405-A.
 PD 05-OCT-1988.
 PF 30-MAR-1988; 302862.
 PR 31-MAR-1987; US-032859.
 PA (UJJO) John Hopkins University.
 PI Iehizaka K, Martens CL, Moore KW;
 DR WPI; 88-280165/40.
 DR N-PSDB; N81168.
 PT Nucleic acid encoding glycosylation inhibiting peptide - capable of
 PT suppressing IgE immunoglobulin responses, for use as antiinflammatory
 PT agents.
 PS Disclosure; i 29pp; English.
 CC The polypeptide has the following biological activities: IgE binding
 CC factor switching activity, inhibition of IgE-induced Fc-epsilon receptor
 CC expression, and inhibition of phospholipase A2 activity after
 CC dephosphorylation.
 SQ Sequence 286 AA;

DB 1; Score 106; Match 32.1%; QryMatch 1.1%; Pred. No. 1.57e+01;
 Matches 17; Conservative 13; Mismatches 19; Indels 4; Gaps 4;

Db 168 ysyvglasspmrpylwkkvqitqgqlvqfvtlqtcgvfwpcsfplg-w 218

! : ! ! ! ! : ! : ! : : : : : : ! : ! : ! : !

QY 987 YTSGLSSYNGYPFLEWEQYISIRHLLLSISWLA-C-TFLVCAVFLAPW 1037

RESULT 13

ID P82112 standard; protein; 816 AA.
 AC P82112;
 DT 22-OCT-1990 (first entry)
 DE Human retinoblastoma gene product
 KW Retinoblastoma; neoplasm; osteocarcoma.
 OS Homo sapiens.
 PN EP-259031-A.
 PD 09-MAR-1988.
 PF 11-AUG-1987; 307095.
 PR 11-AUG-1986; US-895163.
 PA (MASS-) Massachusetts Eye; (WHIT-) Whitehead Inst Biomed Re.
 PI Dryja TP, Friend S;
 DR WPI; 88-065827/10.
 DR N-PSDB; N81369.
 PT Diagnosis of retinoblastoma - using genetic material corresp. to a normal
 PT human retinoblastoma gene or a unique sub-region
 PS Disclosure; p; English.
 CC The absence of this protein is associated with a neoplasm. The corresp.
 CC gene is used to produce an antibody which is contacted with the tumour
 CC sample and immune complexes, if detected, indicate the presence in the
 CC tumour sample of the protein. The whole sequence or a unique sub-region
 CC can be used. Method identifies patients lacking the defective retino-
 CC blastoma (Rb) allele and thus are not at risk of developing the disease.
 SQ Sequence 816 AA;

DB 1; Score 106; Match 21.3%; QryMatch 1.1%; Pred. No. 1.57e+01;
 Matches 30; Conservative 43; Mismatches 56; Indels 12; Gaps 11;

Db 557 llshepellehiwt-lfghtlqn-eyelmrdrldqimmsmygickvknldkfkiiw- 613

! : ! : ! : ! : ! : ! : ! : ! : ! : ! : ! : ! : ! : ! : !

QY 819 MLEENKQLPQM-WLHYFRDWLQIQDAFSDWETGRIMP-NNYKNGSDDGV-LAYKLLVQ 875

Db 614 taykdplhavqetfkrvlikeeydsiivfynsvfm-qrktnilgyaetrptlsiph 672
 ! : ! : ! : ! : ! : ! : ! : ! : ! : ! : ! : ! : ! : ! : !

QY 876 TGRDKP IDISQLTKQRLVDADGIINPSAFY--IYLTAWVSNDFVAYAAQANIRPHRPE 933

Db 673 -ip-rspykfpssplripogn 691

: : : ! : ! : ! ! ! : : :

QY 934 WVHDKADYV-PETRLRIPAAE 953

RESULT 14

ID R36534 standard; Protein; 928 AA.
 AC R36534;
 DT 24-AUG-1993 (first entry)
 DE Retinoblastoma (RB) protein.
 KW RB gene product; p5688 portion; cell cycle progression control;
 KW combination; therapeutic methods; arrest; tumorigenesis;
 KW regulation; physiological processes; blood cell prodn.;
 KW gamete prodn.; ss.
 OS Homo sapiens SR-40 cell line.
 PN W09308267-A.
 PD 29-APR-1993.
 PF 16-OCT-1992; U08918.
 PR 17-OCT-1991; US-778510.
 PA (REGC) UNIV CALIFORNIA.
 PI Goodrich DW, Lee EYHF, Lee WH, Wang NP.
 DR WPI; 93-152462/18.
 DR N-PSDB; Q41545.
 PT Method of controlling cell cycle progression - uses purified
 PT retinoblastoma protein or fragment, for use in combination with
 PT therapeutic methods to arrest tumorigenesis
 PS Claim 10; Fig 9; 68pp; English.
 CC The sequence is that of the retinoblastoma (RB) protein which may be
 CC used as part of a method of controlling cell cycle progression which
 CC may be used in combination with therapeutic methods to arrest
 CC tumorigenesis in organisms. The cell cycle can be reversibly
 CC arrested in a convenient and safe manner. The protein is used in
 CC a compsn. is relatively inexpensive and readily obtainable, and
 CC shows little or no toxic effects on healthy cells. It is also
 CC compatible with other methods and devices for regulating certain
 CC physiological processes of the body, such as blood cell prodn. and
 CC gamete prodn. Fragments of the protein are soluble in low concns.
 CC of glycerol thus enhancing their value in pharmaceutical applicns.
 SQ Sequence 928 AA;

DB 7; Score 106; Match 21.3%; QryMatch 1.1%; Pred. No. 1.57e+01;
 Matches 30; Conservative 43; Mismatches 56; Indels 12; Gaps 11;

Db 669 llshepellehiwt-lfghtlqn-eyelmrdrldqimmsmygickvknldkfkiiw- 725

! : ! : ! : ! : ! : ! : ! : ! : ! : ! : ! : ! : ! : ! : !

QY 819 MLEENKQLPQM-WLHYFRDWLQIQDAFSDWETGRIMP-NNYKNGSDDGV-LAYKLLVQ 875

Db 726 taykdplhavqetfkrvlikeeydsiivfynsvfm-qrktnilgyaetrptlsiph 784
 ! : ! : ! : ! : ! : ! : ! : ! : ! : ! : ! : ! : ! : ! : !

QY 876 TGRDKP IDISQLTKQRLVDADGIINPSAFY--IYLTAWVSNDFVAYAAQANIRPHRPE 933

Db 785 -ip-rspykfpssplripogn 803

: : : ! : ! : ! ! ! : : :

QY 934 WVHDKADYV-PETRLRIPAAE 953

RESULT 15

ID P90599 standard; protein; 970 AA.
 AC P90599;
 DT 30-OCT-1989 (first entry)
 DE Human retinoblastoma.
 KW Human retinoblastoma; screening; tumours; probes.

RESULT	3	
ID	DACB BACSU	STANDARD; PR7; 382 AA.
AC	P35150;	
DT	01-FEB-1994	(REL. 28, CREATED)
DT	01-FEB-1994	(REL. 28, LAST SEQUENCE UPDATE)
DT	01-FEB-1995	(REL. 31, LAST ANNOTATION UPDATE)
DE	PENICILLIN-BINDING PROTEIN 5* PRECURSOR (D-ALANYL-D-ALANINE	
DE	CARBOXYPEPTIDASE) (EC 3.4.16.4) (DD-PEPTIDASE) (DD-CARBOXYPEPTIDASE)	
DE	(PBP-5*).	
GN	DACB.	
OS	BACILLUS SUBTILIS.	
OC	PROKARYOTA; FIRMICUTES; ENDOSPORE-FORMING RODS AND COCCI; BACILLACEAE.	
RN	[1]	
RP	SEQUENCE FROM N.A., AND PARTIAL SEQUENCE.	
RM	92193254	
RA	BUCHANAN C.E., LING M.-L.;	
RL	J. BACTERIOL. 174:1717-1725(1992).	
RL	[2]	
RP	SEQUENCE FROM N.A.	
RC	STRAIN=168 / MARBURG;	
RM	95020538	
RA	SOROKIN A.V., ZUMSTEIN E., AZEVEDO V., EHRLICH S.D., SERROR P.;	
RL	MOL. MICROBIOL. 10:385-395(1993).	
CC	!- FUNCTION: REMOVES C-TERMINAL D-ALANYL RESIDUES FROM SUGAR-PEPTIDE	

DB 2; Score 137; Match 27.6%; QryMatch 1.4%; Pred. No. 3.40e-03;
Matches 43; Conservative 39; Mismatches 62; Indels 12; Gaps 10;

RESULT	6	STANDARD;	PRT;	413 AA.
ID	YMB5 CAEEL			
AC	Q03602;			
DT	01-FEB-1994	(REL. 28, CREATED)		
DT	01-FEB-1994	(REL. 28, LAST SEQUENCE UPDATE)		
DT	01-JUN-1994	(REL. 29, LAST ANNOTATION UPDATE)		
DE	HYPOTHETICAL 47.0 KD PROTEIN F54G8.5 IN CHROMOSOME III.			
GN	F54G8.5.			

RP SEQUENCE FROM N.A.
RC STRAIN=K12;
RM 88228015
RA MAIDEN M.C.J., JONES-MORTIMER M.C., HENDERSON P.J.F.;
RL J. BIOL. CHEM. 263:8003-8010(1988).
RN [3]
RP PRELIMINARY SEQUENCE OF 1-28 FROM N.A.
RM 84114868
RA STONER C., SCHLEIF R.F.;
RL J. MOL. BIOL. 171:369-381(1983).

RP SEQUENCE FROM N.A., AND SEQUENCE OF 4-25.

FT	TRANSMEM	177	197	POTENTIAL.
----	----------	-----	-----	------------

Jan 17 17:11

US-08-319-745-10.rsp

15

```
FT TRANSMEM 242 262 POTENTIAL.
*FT TRANSMEM 281 301 POTENTIAL.
FT TRANSMEM 325 345 POTENTIAL.
FT TRANSMEM 347 367 POTENTIAL.
FT TRANSMEM 385 405 POTENTIAL.
FT TRANSMEM 407 427 POTENTIAL.
SQ SEQUENCE 435 AA; 48248 MW; 1105824 CN;

DB 8; Score 121; Match 19.9%; QryMatch 1.2%; Pred. No. 2.40e-01;
Matches 29; Conservative 43; Mismatches 68; Indels 6; Gaps 6;

Db 286 wlistisgiavgwlv-dyfkikgypntkvyrtvliivgms-fgffflgsilttnnitvalli 343
Qy 1013 WLLSISVVLACTFIVCAVFLIAPWTAGIIVVIALMTVELFGMGGLIGIKLSAVPWIL 1072

Db 344 cisiglagiatapvgwseisaelapigsvmslsmvnlannlfggiaaaltgylfdvtg 403
Qy 1073 IASVGI-GVEFTVHVALAFLTAIGDKNHRAMIA-LEHMFAPVLDGAVSTLLGVLMLAGSE 1130

Db 404 -sftlsflvagfvillgl-vfyvfv1 427
Qy 1131 FDEIVRYFFAVLAILTVLGVNLGLVL 1156

Search completed: Wed Jan 17 17:18:47 1996
Job time : 101 secs.
```

WPSRA (TM)

Release 2.1D John F. Collins, Biocomputing Research Unit.
Copyright (c) 1993, 1994, 1995 University of Edinburgh, U.K.
Distribution rights by IntelliGenetics, Inc.

MParch_pp protein - protein database search, using Smith-Waterman algorithm

Run on: Wed Jan 17 17:19:05 1996; MasPar time 45.33 Seconds
713.356 Million cell updates/sec

Tabular output not generated.

Title: >US-08-319-745-10

Description: (1:1356) from US08319745.pep

Perfect Score: 9913

Sequence: 1 MASAGNARGPGQGRREA.....TORPPWAALCPATASPSPL 1356

Scoring table: PAM 150

Gap 11

Searched: 78488 seqs, 23849247 residues

Database: pir45

1 ann1
2 ann2
3 ann3
4 unann1
5 unann2
6 unann3
7 unann4
8 unann5
9 unann6
10 unann7
11 unrev1
12 unrev2

Statistics: Mean 55.272; Variance 153.526; scale 0.360

Pred. No. is the number of results predicted by chance to have a score greater than or equal to the score of the result being printed, and is derived by analysis of the total score distribution.

SUMMARIES

Result No.	Score	Query Match	Length	ID	Description	Pred. No.
1	2806	28.3	1286	9	A33468 probable membrane pr	0.00e+00
2	2703	27.3	1299	9	S06119 membrane protein pat	0.00e+00
3	410	4.1	1170	11	S52525 hypothetical protein	7.72e-36
4	236	2.4	633	9	S44795 F09G8.3 protein - Ca	4.06e-13
5	145	1.5	464	8	S22697 extensin - Volvox ca	8.31e-03
6	143	1.4	382	7	S45552 penicillin-binding p	1.32e-02

7	143	1.4	382	7	B42274	penicillin-binding p	1.32e-02
8	137	1.4	711	8	C40046	antibiotic transport	5.23e-02
9	137	1.4	712	3	BVBRGB	cyaB protein - Borde	5.23e-02
10	135	1.4	215	7	S16564	nolH protein - Rhizo	8.20e-02
11	134	1.4	413	9	S28276	hypothetical protein	1.03e-01
12	130	1.3	472	7	B26430	arabinose transport	2.49e-01
13	130	1.3	403	12	S52796	prpL_2 protein - hum	2.49e-01
14	130	1.3	1063	7	A33830	cation efflux system	2.49e-01
15	129	1.3	295	10	B48013	proline-rich proteog	3.09e-01
16	128	1.3	226	6	S33024	hypothetical protein	3.85e-01
17	128	1.3	964	7	S18537	envD protein - Esche	3.85e-01
18	128	1.3	464	10	A47655	spliceosome-associat	3.85e-01
19	127	1.3	286	7	F43670	integral membrane pr	4.78e-01
20	123	1.2	1188	8	S49915	extensin-like protei	1.12e+00
21	123	1.2	246	11	S15378	H+-transporting ATP	1.12e+00
22	119	1.2	915	9	S44797	F09G8.4 protein - Ca	2.60e+00
23	119	1.2	353	9	S36438	EPPT protein - hydro	2.60e+00
24	119	1.2	1596	9	A41216	guanine nucleotide e	3.19e+00
25	118	1.2	430	8	JC2301	hypothetical 47.8K p	3.92e+00
26	117	1.2	199	7	JQ2137	NADH dehydrogenase (3.92e+00
27	117	1.2	1711	10	A47392	chromodomain-helicas	3.92e+00
28	117	1.2	666	7	A44797	low-calcium-respons	3.92e+00
29	116	1.2	502	10	A55197	Wiskott-Aldrich synd	4.81e+00
30	116	1.2	307	6	B36125	branched-chain amino	4.81e+00
31	116	1.2	126	7	S30744	hypothetical protein	4.81e+00
32	116	1.2	126	11	S30707	hypothetical protein	4.81e+00
33	115	1.2	439	8	S45025	chitinase (EC 3.2.1.	5.89e+00
34	115	1.2	381	1	S36011	ubiquinol--cytochrom	5.89e+00
35	115	1.2	436	6	S29891	secY protein - Micro	5.89e+00
36	114	1.2	226	7	C29504	hypothetical 24K pro	7.20e+00
37	112	1.1	454	7	S16565	nolI protein - Rhizo	1.07e+01
38	112	1.1	472	7	S47089	arabinose-proton sym	1.07e+01
39	112	1.1	254	7	S35738	guFA protein - Myxoc	1.07e+01
40	112	1.1	1063	3	GNWVRA	structural polyprote	1.07e+01
41	112	1.1	254	7	S39876	hypothetical protein	1.07e+01
42	112	1.1	1733	3	B45344	probable nuclear ant	1.07e+01
43	112	1.1	254	7	S33148	guFA protein - Myxoc	1.07e+01
44	111	1.1	376	7	S47693	hypothetical protein	1.31e+01
45	111	1.1	514	3	W2WLB5	E2 protein - human p	1.31e+01

ALIGNMENTS

RESULT	1
ENTRY	A33468
TITLE	#type complete probable membrane protein patched - fruit fly (Drosophila melanogaster)
ORGANISM	#formal_name Drosophila melanogaster
DATE	20-Dec-1989 #sequence_revision 20-Dec-1989 #text_change 27-Jan-1995
ACCESSIONS	A33468
REFERENCE	A33468
#authors	Hooper, J.E.; Scott, M.P.
#journal	Cell (1989) 59:751-765
#title	The Drosophila patched gene encodes a putative membrane protein required for segmental patterning.
#cross-references	MUID:90058658
#accession	A33468
#status	preliminary; not compared with conceptual translation
#molecule	type mRNA
#residues	1-1286 #label HOO
#cross-references	GB:M28418; GB:M28999
#note	nucleotide sequence is not given
KEYWORDS	membrane protein

Qy 868 LAYLLVQTGSRDKPIDISLTKQLVDADGGINPSAFYIYLTAWVSNDFVAYAASQANI 927

DB 9; Score 2703; Match 40.1%; QryMatch 27.3%; Pred. No. 0.00e+00; Mismatches 471; Conservative 304; Mismatches 306; Indels 95; Gaps 5


```
* DB 7; Score 143; Match 28.0%; QryMatch 1.4%; Pred. No. 1.32e-02;
Matches 21; Conservative 22; Mismatches 28; Indels 4; Gaps 3;

Db 233 staskdgidliavtindpdwdchmkmfnyfeyhytyliakgdpikgtfy-skaf 291
Qy 754 TTRVRDGLDITDIPRETREYDFIAAQFKY-FSFYNMVIVTQAKDYPNTQHLVDLHRSF 812

Db 292 ikrdityllteeke 306
Qy 813 --SNVKYVMLENKQ 825

RESULT 8
ENTRY C40046 #type complete
TITLE antibiotic transport-associated protein actII-3 -
Streptomyces coelicolor
ORGANISM #formal_name Streptomyces coelicolor
DATE 30-Jun-1992 #sequence_revision 30-Jun-1992 #text_change
18-Jun-1993
ACCESSIONS C40046
REFERENCE #authors Fernandez-Moreno, M.A.; Caballero, J.L.; Hopwood, D.A.;
Malpartida, F.
#journal Cell (1991) 66:769-780
#title The act cluster contains regulatory and antibiotic export
genes, direct targets for translational control by the bldA
crRNA gene of Streptomyces.
#cross-references MUID:91347376
#accession C40046
##molecule_type DNA
##residues 1-711 ##label FER
##cross-references GB:M64683
SUMMARY #length 711 #molecular-weight 74862 #checksum 4136

DB 8; Score 137; Match 20.1%; QryMatch 1.4%; Pred. No. 5.23e-02;
Matches 31; Conservative 57; Mismatches 60; Indels 6; Gaps 5;

Db 184 lliivtiivtyzspilwlpisagm-slvisgaivllknagltvn-aqtamiltvl 241
Qy 1015 LLSISWVLACTFLVCAVFLNPWTAGIIVWVLMTVELFGMMGLIGIKLSAVPVILIA 1074

Db 242 vlgaatdyallvayreelrrhrdheamavalrragpaivaasaatvavmvlilaal 301
Qy 1075 SVGIGVETVHVALAFLTAIG--DKNHRAH--LALEHMFAPVLDGAVSTLLGVMLAGSEF 1131

Db 302 n-stkgipgcavgvllvqlsmmtllpallvifg 334
Qy 1132 DFIVRYFFAVLAILTVLGVINGLVLPLVLSFFG 1165

RESULT 9
ENTRY BVRCB #type complete
TITLE cyab protein - Bordetella pertussis
ORGANISM #formal_name Bordetella pertussis
DATE 31-Dec-1990 #sequence_revision 31-Dec-1990 #text_change
08-Dec-1994
ACCESSIONS S02386
REFERENCE #authors Glaeser, P.; Sakamoto, H.; Bellalou, J.; Ullmann, A.; Danchin,
A.
#journal EMBO J. (1988) 7:3997-4004
#title Secretion of cyclolysin, the calmodulin-sensitive adenylate
cyclase--haemolysin bifunctional protein of Bordetella
```

```
pertussis.
#cross-references MUID:89091151
#accession S02386
##molecule_type DNA
##residues 1-712 ##label GLA
COMMENT This protein is required for the transport of cyclolysin (or
calmodulin-sensitive adenylate cyclase--hemolysin bifunctional
protein, encoded by cyab) across the cell envelope and for its
release into the external medium. This secretion process is very
similar to that of the E. coli alpha-hemolysin.

GENETICS
#gene cyab
CLASSIFICATION #superfamily hemolysin secretion protein B; malk protein
homology
KEYWORDS ATP binding; cyclolysin transport; membrane protein; P-loop
FEATURE
488-682 #domain malk protein homology #label MK1\
505-513 #region nucleotide-binding motif A (P-loop)\
629-633 #region nucleotide-binding motif B\
511 #binding site ATP (Lys) #status predicted
SUMMARY #length 712 #molecular-weight 77969 #checksum 2892

DB 3; Score 137; Match 27.6%; QryMatch 1.4%; Pred. No. 5.23e-02;
Matches 43; Conservative 39; Mismatches 62; Indels 12; Gaps 10;

Db 166 lviqlflltplffqvmdkvlvnmamelnvlgvflaailfeal-ltgirtylfahts 224
Qy 1014 LLSISWVLACTFLVCAVFLNPWTAGIIVWVLMTVELFGMMGLIGIK--LSAPVP 1069

Db 225 skldvelgarlyahllrplayfgarvgsdvarvr-elehirafitgnavtvlldv-vf 282
Qy 1070 VILASVGIGV-FETVHVALFLTA--IGDKNHRAHLEHMFAPVLDGAVSTLLGVML 1126

Db 283 svv-f-iavmfysvkltlvllaalpcyflilvlvt 316
Qy 1127 AGSEDFIVRYFFAVLAILTVLGVINGLVLPLVLS 1162

RESULT 10
ENTRY S16564 #type complete
TITLE nolH protein - Rhizobium meliloti
ORGANISM #formal_name Rhizobium meliloti
DATE 13-Jan-1995 #sequence_revision 13-Jan-1995 #text_change
13-Jan-1995
ACCESSIONS S16564
REFERENCE #authors Baev, N.; Endre, G.; Petrovics, G.; Banfalvi, Z.; Kondorosi,
A.
#journal Mol. Gen. Genet. (1991) 228:113-124
#title Six nodulation genes of nod box locus 4 in Rhizobium meliloti
are involved in nodulation signal production: nodM codes
for D-glucosamine synthetase.
#cross-references MUID:91360053
#accession S16564
##status preliminary
##molecule_type DNA
##residues 1-215 ##label BAE
##cross-references EMBL:X58632
SUMMARY #length 215 #molecular-weight 23775 #checksum 2405

DB 7; Score 135; Match 23.4%; QryMatch 1.4%; Pred. No. 8.20e-02;
Matches 32; Conservative 47; Mismatches 49; Indels 9; Gaps 8;

Db 1 mfl-nwstrvtitlplvisvgitfaaiyalgftlnmtlmalsisigiliddtiivren 59
```

WATERMAN

(TM)

Release 2.1D John F. Collins, Biocomputing Research Unit.
Copyright (c) 1993, 1994, 1995 University of Edinburgh, U.K.
Distribution rights by IntelliGenetics, Inc.

MPsrch_pp protein - protein database search, using Smith-Waterman algorithm
Run on: Wed Jan 17 17:15:17 1996; MasPar time 16.67 Seconds
499.639 Million cell updates/sec
Tabular output not generated.

Title: >US-08-319-745-4
Description: (1:1311) from US08319745.pep
Perfect Score: 9491
Sequence: 1 MWAPDSEAPSNPRITAAHES.....YRDERHRAPEKRRQFWT 1311

Scoring table: PAM 150
Gap 11

Searched: 53402 seqs, 6354270 residues

Database: a-geneseq18
1 part1
2 part2
3 part3
4 part4
5 part5
6 part6
7 part7
8 part8
9 part9
10 part10

Statistics: Mean 40.660; Variance 210.188; scale 0.193

Pred. No. is the number of results predicted by chance to have a score greater than or equal to the score of the result being printed, and is derived by analysis of the total score distribution.

SUMMARIES

Result No.	Score	Query Match	Length	ID	Description	Pred. No.
1	267	2.8	162	7 R37991	Sequence of a new cyt	3.41e-10
2	159	1.7	3910	7 R38470	ALL-1 protein.	1.27e-02
3	159	1.7	1400	8 R44514	MLL amino acid sequen	1.27e-02
4	150	1.6	3969	9 R52971	Product of the cDNA e	4.94e-02
5	144	1.5	765	1 P92275	Human topoisomerase I	1.21e-01
6	144	1.5	265	3 R12844	HTLV-1 protein expres	1.21e-01
7	137	1.4	740	5 R27530	Plasmodium falciparum	3.37e-01
8	133	1.4	123	9 R51053	Eppstein-Barr nuclear	6.02e-01

9	133	1.4	1023	3 R13319	Partial Human Natural	6.02e-01
10	129	1.4	193	3 P60624	Sequence B encoded by	1.07e+00
11	129	1.4	751	1 P94776	Novel amyloid precurs	1.07e+00
12	129	1.4	179	6 R32082	Plasmid pSCHEC-1414 p	1.07e+00
13	129	1.4	179	3 R14216	HCV detecting peptide	1.07e+00
14	127	1.3	657	6 R29580	FMR-1 gene product.	1.42e+00
15	124	1.3	1141	6 R31961	Human cardiac cgr PDE	2.17e+00
16	123	1.3	566	3 R20181	Sequence encoded by 6	2.50e+00
17	122	1.3	3025	1 P93284	Sequence of clone HIV	2.88e+00
18	120	1.3	301	2 P70867	Sequence of acidic ba	3.80e+00
19	116	1.2	911	3 R15355	Human erythrocyte mem	6.61e+00
20	114	1.2	1284	1 P81187	Sequence encoded by a	8.70e+00
21	113	1.2	517	4 R22904	1-Caldesmon.	9.97e+00
22	111	1.2	720	3 R15381	Pseudomonas SY77-glut	1.31e+01
23	111	1.2	2237	6 R33550	Sequence of the alpha	1.31e+01
24	111	1.2	720	3 R14445	Pseudomonas SY77-glut	1.31e+01
25	111	1.2	720	3 R15382	Pseudomonas SY77-glut	1.31e+01
26	111	1.2	720	3 R15380	Pseudomonas SY77-glut	1.31e+01
27	111	1.2	561	3 P61363	Soybean glycinin A5M4	1.31e+01
28	111	1.2	720	3 R15379	Pseudomonas SY77-glut	1.31e+01
29	110	1.2	1784	1 R05898	Gene product of first	1.50e+01
30	110	1.2	445	3 R14163	Cellular DNA-binding	1.50e+01
31	109	1.1	493	5 R26944	P.falciparum LSA gene	1.71e+01
32	109	1.1	292	3 P60645	Mouse kidney cell Ban	1.71e+01
33	109	1.1	316	5 R26941	P.falciparum LSA-R-NR	1.71e+01
34	108	1.1	3080	1 P93285	Sequence of clone HIV	1.96e+01
35	107	1.1	462	1 R05766	Portion of peptide an	2.24e+01
36	106	1.1	382	7 R39224	Nucleocapsid protein	2.56e+01
37	106	1.1	318	5 R26943	P.falciparum LSA N-te	2.56e+01
38	105	1.1	3210	1 P81770	Deduced sequence enco	2.92e+01
39	105	1.1	312	5 R27361	Sequence of a polypep	2.92e+01
40	105	1.1	532	5 R27362	Sequence of a polypep	2.92e+01
41	105	1.1	558	5 R27363	Sequence of a polypep	2.92e+01
42	104	1.1	482	3 R21409	NADH dehydrogenase su	3.33e+01
43	103	1.1	1026	9 R48993	read S-lyase protein.	3.80e+01
44	103	1.1	307	3 R12785	Caldesmon-like polype	3.80e+01
45	103	1.1	1931	5 R27649	Human calcium channel	3.80e+01

ALIGNMENTS

RESULT 1
ID R37991 standard; Protein; 162 AA.
AC R37991;
DT 29-SEP-1993 (first entry)
DE Sequence of a new cytokine which inhibits induction by gamma
DE interferon of expression of Class II histocompatibility antigens.
KW Cytokine; interferon-gamma antagonist; autoimmune disease therapy;
KW transplants; Class II histocompatibility antigens;
KW gamma interferon; leukaemia line K562.
OS Homo sapiens.
PN W09311232-A.
PD 10-JUN-1993.
PF 02-DEC-1992; F01123.
PR 02-DEC-1991; FR-014908.
PA (INRM) INSERM INST NAT SANTE & RECH MED.
PI Augery-bourget Y, Azzarone B, Boucheix C, Jasmin C;
PI Krief PH;
DR WPI; 93-197051/24.
DR N-PSDB; Q43704.
PT New cytokine as interferon-gamma antagonist - inhibits induction
PT of class II histocompatibility antigens on cell surface by
PT IFN-gamma, for treating and preventing autoimmune disorders
PS Claim 1; Fig 1; 44pp; French.

CC 82% similarity respectively, to the *Drosophila* gene. The third region

KW immunodeficiency; developmental abnormalities; inherited diseases;

1000

Db 171 nekkkkkf 179
 { : : : :

Db 171 nekkkkkf 179
 { : : : :

WATERMAN

(TM)

Release 2.1D John F. Collins, Biocomputing Research Unit.
Copyright (c) 1993, 1994, 1995 University of Edinburgh, U.K.
Distribution rights by IntelliGenetics, Inc.

MPsrch_pp protein - protein database search, using Smith-Waterman algorithm

Run on: Wed Jan 17 17:09:43 1996; MasPar time 28.57 Seconds
703.808 Million cell updates/sec

Tabular output not generated.

Title: >US-08-319-745-4

Description: (1:1311) from US08319745.pep

Perfect Score: 9491

Sequence: 1 MVAPDSEAPSNPRTAAHES.....YRDERDHRAPREKRRQFWT 1311

Scoring table: PAM 150

Gap 11

Searched: 43470 seqs, 15335248 residues

Database: swiss-prot31

1 part1

2 part2

3 part3

4 part4

5 part5

6 part6

7 part7

8 part8

Statistics: Mean 57.597; Variance 137.878; scale 0.418

Pred. No. is the number of results predicted by chance to have a score greater than or equal to the score of the result being printed, and is derived by analysis of the total score distribution.

SUMMARIES

Result No.	Score	Query Match	Length	ID	Description	Pred. No.
1	4527	47.7	1286	5	PATC DROME MEMBRANE PROTEIN PATC	0.00e+00
2	342	3.6	382	6	RDP HUMAN RD PROTEIN.	7.34e-31
3	339	3.6	1106	7	STC DROME SHUTTLE CRAFT PROTEIN	2.10e-30
4	329	3.5	375	6	RDP MOUSE RD PROTEIN (W1623).	6.91e-29
5	295	3.1	471	6	RUI7 XENIA U1 SMALL NUCLEAR RIBO	8.42e-24
6	255	2.7	614	6	RUI7 HUMAN U1 SMALL NUCLEAR RIBO	5.48e-18
7	255	2.7	448	6	RUI7 DROME U1 SMALL NUCLEAR RIBO	5.48e-18
8	251	2.6	185	7	T2 MOUSE OCTAPEPTIDE-REPEAT PR	2.03e-17
9	225	2.4	633	8	YL53 CAEL HYPOTHETICAL 70.7 KD	8.78e-14
10	202	2.1	494	7	SR75_HUMAN PRE-MRNA SPLICING FAC	1.13e-10

11	198	2.1	483	7	VE2 HPV14	E2 PROTEIN.	3.81e-10
12	194	2.0	244	8	YL53 CAEL	HYPOTHETICAL 29.0 KD	1.27e-09
13	190	2.0	475	7	UZAF MOUSE	SPLICING FACTOR UZAF	4.22e-09
14	190	2.0	475	7	UZAF HUMAN	SPLICING FACTOR UZAF	4.22e-09
15	186	2.0	330	6	RLX3 STAAU	RLX PROTEIN.	1.39e-08
16	182	1.9	502	7	VE2 HPV25	E2 PROTEIN.	4.51e-08
17	182	1.9	197	7	TRSF DROME	FEMALE-SPECIFIC TRANS	4.51e-08
18	181	1.9	221	7	SC35 CHICK	SPLICING FACTOR SC35	6.04e-08
19	181	1.9	221	7	SC35 HUMAN	SPLICING FACTOR SC35	6.04e-08
20	174	1.8	429	7	TO34 YEAST	TOM34 PROTEIN.	4.61e-07
21	173	1.8	233	8	YOD2 CAEL	HYPOTHETICAL 26.8 KD	6.14e-07
22	171	1.8	1523	7	SON HUMAN	SON PROTEIN (SON3).	1.09e-06
23	169	1.8	349	7	SR55 DROME	SERINE-ARGININE PROTE	1.92e-06
24	167	1.8	1549	7	TRHY SHEEP	TRICHOHYALIN.	3.39e-06
25	167	1.8	695	8	XE7 HUMAN	PROTEIN XE7.	3.39e-06
26	167	1.8	252	8	VPHE NPVAC	POLYHEDRAL ENVELOPE P	3.39e-06
27	166	1.7	493	7	VE2 HPV19	E2 PROTEIN.	4.49e-06
28	165	1.7	568	2	DISC DROME	DISCONNECTED PROTEIN.	5.95e-06
29	165	1.7	164	8	X16 HUMAN	PRE-MRNA SPLICING FAC	5.95e-06
30	163	1.7	346	8	INPI CAEL	HYPOTHETICAL 42.2 KD	1.04e-05
31	162	1.7	1061	6	RNE ECOLI	RIBONUCLEASE E (EC 3.	1.82e-05
32	161	1.7	767	7	TOPI MOUSE	DNA TOPOISOMERASE I (1.82e-05
33	160	1.7	767	7	TOPI CRIGR	DNA TOPOISOMERASE I (2.40e-05
34	159	1.7	57	4	HSP1 DROMA	SPERM PROTEIN P1 (C	3.16e-05
35	159	1.7	3969	4	HRX HUMAN	ZINC FINGER PROTEIN H	3.16e-05
36	158	1.7	503	8	YOW5 CAEL	HYPOTHETICAL 57.1 KD	4.17e-05
37	155	1.6	1023	3	GLT DROME	GLUTACTIN PRECURSOR.	9.47e-05
38	154	1.6	514	7	VE2 HPV5B	E2 PROTEIN.	1.24e-04
39	152	1.6	241	8	YOB7 CAEL	HYPOTHETICAL 28.5 KD	2.13e-04
40	152	1.6	240	7	UZAG HUMAN	SPLICING FACTOR UZAF	2.13e-04
41	150	1.6	264	7	TRA2 DROME	TRANSFORMER-2 SEX-DET	3.65e-04
42	150	1.6	1407	7	TRHY RABIT	TRICHOHYALIN.	3.65e-04
43	149	1.6	461	7	VE2 HPV09	E2 PROTEIN.	4.77e-04
44	147	1.5	1898	7	TRHY HUMAN	TRICHOHYALIN.	8.11e-04
45	146	1.5	488	7	VE2 HPV49	E2 PROTEIN.	1.06e-03

ALIGNMENTS

RESULT	1	ID	PATC DROME	STANDARD;	PRT;	1286 AA.
AC	P18502;					
DT	01-NOV-1990 (REL. 16, CREATED)					
DT	01-NOV-1990 (REL. 16, LAST SEQUENCE UPDATE)					
DT	01-FEB-1994 (REL. 28, LAST ANNOTATION UPDATE)					
DE	MEMBRANE PROTEIN PATCHED.					
GN	PTC.					
OS	DROSOPHILA MELANOGASTER (FRUIT FLY).					
OC	EUKARYOTA; METAZOA; ARTHROPODA; INSECTA; DIPTERA.					
RN	[1]					
RP	SEQUENCE FROM N.A.					
RM	90058658					
RA	HOOPER J.E., SCOTT M.P.;					
RL	CELL 59:751-765(1989).					
RN	[2]					
RP	SEQUENCE FROM N.A.					
RM	90015164					
RA	AKANO Y., GUERRERO I., HIDALGO A., TAYLOR A., WHITTLE J.R.S.,					
RL	INGHAM P.W.;					
CC	NATURE 341:508-513(1989).					
CC	-!- FUNCTION: SEGMENTATION POLARITY PROTEIN. EXACT FUNCTION NOT					
CC	KNOWN. PTC PROBABLY PARTICIPATES IN CELL INTERACTIONS THAT					
CC	ESTABLISH PATTERN WITHIN THE SEGMENT.					
CC	-!- SUBCELLULAR LOCATION: INTEGRAL MEMBRANE PROTEIN.					

ALL INFORMATION CONTAINED HEREIN IS UNCLASSIFIED

— —
• •
• •
——
• •
——
——
——
——
——
——

THE
OFFICE OF THE ATTORNEY GENERAL
STATE OF TEXAS

THE
OFFICE OF THE ATTORNEY GENERAL
STATE OF TEXAS

W A P S R E A (TM)

Release 2.1D John F. Collins, Biocomputing Research Unit.
Copyright (c) 1993, 1994, 1995 University of Edinburgh, U.K.
Distribution rights by Intelligenetics, Inc.

MPerch_pp protein - protein database search, using Smith-Waterman algorithm

Run on: Wed Jan 17 17:12:10 1996; MasPar time 43.92 Seconds
711.918 Million cell updates/sec

Tabular output not generated.

Title: >US-08-319-745-4
Description: (1:1311) from US08319745.pep
Perfect Score: 9491
Sequence: 1 MWAPDSEAPNPRITAAHES.....YDERDHRASPREKQRFWT 1311

Scoring table: PAM 150
Gap 11

Searched: 78488 seqs, 23849247 residues

Database: pir45
1 ann1
2 ann2
3 ann3
4 unann1
5 unann2
6 unann3
7 unann4
8 unann5
9 unann6
10 unann7
11 unrev1
12 unrev2

Statistics: Mean 55.100; Variance 168.754; scale 0.327

Pred. No. is the number of results predicted by chance to have a score greater than or equal to the score of the result being printed, and is derived by analysis of the total score distribution.

SUMMARIES

Result No.	Score	Query Match	Length	ID	Description	Pred. No.
1	4527	47.7	1286	9	A33468 probable membrane pr	0.00e+00
2	4372	46.1	1299	9	S06119 membrane protein pat	0.00e+00
3	342	3.6	380	12	S36789 gene RD protein - hu	5.32e-24
4	342	3.6	382	9	A33640 class III histocompa	5.32e-24
5	342	3.6	325	9	JH0189 arginine/aspartate-r	5.32e-24
6	329	3.5	375	10	A40112 MHC-region RD-repeat	1.93e-22

7	300	3.2	1170	11	S52525	hypothetical protein	5.31e-19
8	295	3.1	471	9	S02016	U1 snRNP 70K protein	2.05e-18
9	260	2.7	378	10	S04336	U1 snRNP 70K protein	2.27e-14
10	255	2.7	301	10	S03048	U1 snRNP 70K protein	8.37e-14
11	255	2.7	437	12	S52484	ribonucleoprotein 68	8.37e-14
12	255	2.7	448	9	A36311	70K U1 small nuclear	8.37e-14
13	255	2.7	614	9	A25707	70K small nuclear r1	8.37e-14
14	255	2.7	437	12	S41225	70K protein - human	8.37e-14
15	239	2.5	303	8	S28147	U1 snRNP 70K protein	5.26e-12
16	225	2.4	633	9	S44795	F09C8.3 protein - Ca	1.85e-10
17	202	2.1	494	10	A48133	pre-mRNA splicing SR	5.60e-08
18	198	2.1	483	11	S36470	envelope protein - h	1.48e-07
19	197	2.1	783	10	A55817	cyclin-dependent kin	1.89e-07
20	194	2.0	244	9	S44822	F44E2.3 protein - Ca	3.89e-07
21	192	2.0	492	10	S19938	splicing factor U2AF	6.30e-07
22	192	2.0	492	12	S22646	splicing factor U2AF	6.30e-07
23	190	2.0	475	10	S20250	splicing factor U2AF	1.02e-06
24	188	2.0	484	9	A40988	54K arginine-rich nu	1.64e-06
25	186	2.0	330	6	S28102	rlx protein - Staphy	2.63e-06
26	182	1.9	197	9	A29648	female-specific tran	6.77e-06
27	182	1.9	502	6	S36494	envelope protein - h	6.77e-06
28	181	1.9	221	10	A42634	splicing factor SC35	8.57e-06
29	181	1.9	416	9	A48249	pre-mRNA splicing fa	8.57e-06
30	181	1.9	221	9	B42701	PR264 protein - chic	8.57e-06
31	181	1.9	221	9	S17327	PR264 protein - chic	8.57e-06
32	181	1.9	221	10	S17328	PR264 protein - huma	8.57e-06
33	181	1.9	221	10	A42701	PR264 protein - huma	8.57e-06
34	174	1.8	429	8	S45459	hypothetical protein	4.38e-05
35	173	1.8	233	9	S44882	2C262.2 protein - Ca	5.51e-05
36	171	1.8	483	10	PN0099	Son3 protein - human	8.73e-05
37	169	1.8	350	9	A40459	nuclear phosphoprote	1.38e-04
38	169	1.8	350	9	S14620	RS55 protein - fruit	1.38e-04
39	167	1.8	1549	10	A40691	trichohyalin (EF han	2.18e-04
40	167	1.8	1549	10	S32633	trichohyalin - sheep	2.18e-04
41	167	1.8	252	6	C43679	ORF3 protein - Autog	2.18e-04
42	167	1.8	1203	9	S26650	DNA-binding protein	2.18e-04
43	166	1.7	493	6	S36488	envelope protein - h	2.73e-04
44	165	1.7	164	12	S14016	X16 protein - mouse	3.43e-04
45	165	1.7	568	9	S15008	gene disco protein -	3.43e-04

ALIGNMENTS

RESULT ENTRY TITLE	1	A33468	#type complete
ORGANISM		probable membrane protein patched - fruit fly (Drosophila melanogaster)	
DATE		20-Dec-1989	#sequence_revision 20-Dec-1989
ACCESSIONS		A33468	
REFERENCE		Hooper, J.E.; Scott, M.P.	
#authors		Cell (1989) 59:751-765	
#journal		The Drosophila patched gene encodes a putative membrane protein required for segmental patterning.	
#title		protein required for segmental patterning.	
#cross-references		MUID:90058658	
#accession		A33468	
#status		preliminary; not compared with conceptual translation	
#molecule_type		mRNA	
#residues		1-1286	#label H00
#cross-references		GB:M28418; GB:M28999	
#note		nucleotide sequence is not given	
KEYWORDS		membrane protein	

[illegible]

Db	899	yaampfyhlglcdteqtkltighirdsvkyegf.glpypsgipfifweymtlrslam	958
Qy	894	YTPQPFYLSGLSDTSAIKTLIRSDVCLCKYEAGLPNPPSGICFLFWEQYLYLRTSLLL	953
Db	959	ilacvllaalvlslllsyaaavlsvlsvlaela-qifgaamtllgkilaipavillls	1017
Qy	954	ALACALAAVFIAMVLLINNAWAVLTIA-LATVILQLLGVALLGVKLSAMPVLLVIA	1012
Db	1018	vqmmclfnvlielgfmtesvgnqrivqlsmqmslqplvhqmltsqvavfmlstsfefvi	1077
Qy	1013	TORGUETHVHLCGEVTSICCKRRASIAIESVIAVPVHGALAAALAAWSLAASCEGEVA	1072
Db	1078	rhfcwlllvclvgaacnslvipillamvsgaeelyplehdriestpslpvrssksqk	1137
Qy	1073	RILFURLLLDVFELGLDGLFFIVILSLFGPAAEVRPTEHPERLSTSPKSCPIHPRKSS	1132
Db	1138	syvvqgsrsrsgcsqkshhkhkdlndpsltttiteepgskwsnsaiq	1185
Qy	1133	SSSGGDKSRTS--KSAPRC---APSLTTITERPSWHSASHSVQ	1174
RESULT	2		
ENTRY	S06119	#type complete	
TITLE	membrane protein patched - fruit fly (Drosophila melanogaster)		
ORGANISM	#formal_name Drosophila melanogaster		
DATE	30-Sep-1991	#sequence_revision 30-Sep-1991	#text_change
ACCESSIONS	S06119	09-Sep-1994	
REFERENCE	S06119		
#authors	Nakano, Y.; Guerrero, I.; Hidalgo, A.; Taylor, A.; Whittle, J.R.S.; Ingham, P.W.		
#journal	Nature (1989) 341:508-513		
#title	A protein with several possible membrane-spanning domains encoded by the Drosophila segment polarity gene patched.		
#cross-references	UID:90015164		
#accession	S06119	not compared with conceptual translation	
#status		not compared with conceptual translation	
#molecule_type	DNA		
#residues	1-1299	##label NAK	
GENETICS			
#gene	ptc		
#map_position	2 4D3-D4		
KEYWORDS	glycoprotein; transmembrane protein		
FEATURE			
74-92	#domain transmembrane	#status predicted	#label TM01\
427-448	#domain transmembrane	#status predicted	#label TM02\
456-503	#domain transmembrane	#status predicted	#label TM03\
529-555	#domain transmembrane	#status predicted	#label TM04\
557-585	#domain transmembrane	#status predicted	#label TM05\
677-699	#domain transmembrane	#status predicted	#label TM06\
967-1017	#domain transmembrane	#status predicted	#label TM07\
1019-1047	#domain transmembrane	#status predicted	#label TM08\
1061-1086	#domain transmembrane	#status predicted	#label TM09\
1093-1121	#domain transmembrane	#status predicted	#label TM10\
142,298,335,398,807,861,1194,1271	#binding_site carbohydrate (Asn)	(covalent)	#status predicted

```

#length 1299 #molecular-weight 144091 #checksum 7740
UNWARY
DB 9; Score 4372; Match 53.0%; QryMatch 46.1%; Pred. No. 0.00e+00;
Matches 636; Conservative 259; Mismatches 260; Indels 47; Gaps 27;

```



```

complex.
##cross-references MUID:90126228
#accession A33640
##status preliminary
##molecule_type DNA
##residues 1-382 ##label SPE
##cross-references GB:M3275; GB:M30115; GB:M32276; GB:M33230; GB:M33231
##note the authors translated the codon AAG or residue 95 as Ala
SUMMARY      #length 382   #molecular-weight 43441    #checksum 8129

DB 9; Score     342; Match 57.7%; QryMatch 3.6%; Pred.No. 5.3/e-24;
Matches        45; Conservative 15; Mismatches 16; Indels 2; Gaps 2;

Db 180 sasprsrstsdghrnrddrdreidrdrdrrdrdrdrdrdrdrdrdrdrdrdrdrdrdrdr 239
QY 1231 TPSDRKRSRRSYHYVRRRDREDDRDRDREDRDDRDQRDRDRDRDRDRDRSRERDR-R 1289
       :: | :|:: | :|| || |:||||:||:||:||:||:||:||:||:||:||:||:||
Db 240 ddrdgpfrrdsfpperr 257
QY 1290 DRYRDERDHASP-REKR 1306
       || | :| :| :| :|

ACCESSIONS JH0189          #type complete
REFERENCE   arginine/aspartate-rich 37.3K protein - human RD protein
            #formal name Homo sapiens #common name man
DATE        31-Dec-1991 #sequence_revision 31-Dec-1991 #text_change
           24-Feb-1995

AUTHORS Surowsy, C.S.; Hoganson, G.; Gosink, J.; Strunk, K.; Spritz, R.A.

JOURNAL Gene (1990) 90:299-302
TITLE The human RD protein is closely related to nuclear RNA-binding proteins and has been highly conserved.
CROSS-REFERENCES MUID:90382680
#accession JH0189
#molecule_type mRNA
#residues 1-325 ##label SUR
##cross-references EMBL:X16105
COMMENT This protein consists of alternating basic and acidic amino acid, primarily arginine and aspartic acid, and contains a 'ribonucleoprotein sequence domain'. This protein is closely related to nuclear RNA-binding proteins.
CLASSIFICATION #superfamily ribonucleoprotein repeat homology
FEATURE
195-282 #domain ribonucleoprotein #label RNP\
208-267 #domain ribonucleoprotein repeat homology #label RRM3
SUMMARY      #length 325   #molecular-weight 37216    #checksum 1492

DB 9; Score     342; Match 57.7%; QryMatch 3.6%; Pred.No. 5.3/e-24;
Matches        45; Conservative 15; Mismatches 16; Indels 2; Gaps 2;

Db 124 sasprsrstsdghrnrddrdreidrdrdrdrdrdrdrdrdrdrdrdrdrdrdrdrdrdr 183
QY 1231 TPSDRKRSRRSYHYVRRRDREDDRDRDREDRDDRDQRDRDRDRDRDRDRSRERDR-R 1289
       :: | :|:: | :|| || |:||||:||:||:||:||:||:||::~||| :|:| |
Db 184 ddrdgpfrrdsfpperr 201
QY 1290 DRYRDERDHASP-REKR 1306
       || | :| :| :| :|

```

RESULT	6
ENTRY	A40112 #type complete
TITLE	MHC-region RD-repeat protein - mouse
ORGANISM	#formal name Mus musculus #common name house mouse
DATE	20-Mar-1992 #sequence_revision 20-Mar-1992 #text_change 24-Feb-1995
ACCESSIONS	A40112
REFERENCE	A40112
#authors	Levi-Strauss, M.; Carroll, M.C.; Steinmetz, M.; Meo, T.
#journal	Science (1988) 240:201-204
#title	A previously undetected MHC gene with an unusual periodic structure.
#cross-references	MUID:88178091
#accession	A40112
#status	preliminary
#molecule type mRNA	
#residues_1-375 ##label LEV	
#cross-references GB:M21332	
CLASSIFICATION	#superfamily ribonucleoprotein repeat homology
FEATURE	
267-326	#domain ribonucleoprotein repeat homology #label RRM3
SUMMARY	#length 375 #molecular-weight 42555 #checksum 8746
DB 10; Score	329; Match 57.5%; QryMatch 3.5%; Pred. No. 1.93e-22;
Matches	46; Conservative 15; Mismatches 16; Indels 3; Gaps 3
Dbb	179 easprsrdrshdrrdkrdkddrdrdrdrdrdkdkdkdrdrdrkdrrdrdr 238
Qy	1231 rTSDRSRRSYHYDRRDDEDDREDDRDRDREDRDRDRDRDRERSR-R 1289
Ddb	239 drdr-erd-regfrdsf 256
Qy	1290 DRYDERDHRAPEKQRF 1309
RESULT	7
ENTRY	S52525 #type complete
TITLE	hypothetical protein - yeast (<i>Saccharomyces cerevisiae</i>)
ORGANISM	#formal name Saccharomycetes cerevisiae
DATE	08-May-1995 #sequence_revision 08-May-1995 #text_change 08-May-1995
ACCESSIONS	S52525
REFERENCE	S52519
#authors	Badcock, K.; Churcher, C.
#submission	submitted to the EMBL Data Library, February 1995
#accession	S52525
#status	preliminary
#residues_1-1170 ##label BAD	
#cross-references EMBL:Z48483	
SUMMARY	#length 1170 #molecular-weight 132644 #checksum 5191
DB 11; Score	300; Match 28.7%; QryMatch 3.2%; Pred. No. 5.31e-19;
Matches	52; Conservative 64; Mismatches 56; Indels 9; Gaps 8
Dbb	979 ldmfays-pfyiffvqvtl-gpltklkigsailiffsavflmqiresllalvlvtmi 1036
Qy	929 LPNPPSGIPFF-LFWEQLYLRYSLLAL-ACALAAVFIAVMVLIIAMAVLVUTAIATL 986
Dbb	1037 ivdigalmalgisnavslmllicvglwefcvhvrstfvvpsetkkdnarsrvysl 1096
Qy	987 VIQLGGLWAGKLSPAPVILVAIARGVHETVHLCLGF-VTSIGCKRRRAS--L-AL 1042
Dbb	1097 ntigewikgltkfkgvcwlfafasqskfdvfyrmwftliivaalhfflpallself 1156

identification of an snRNP-associated kinase activity that phosphorylates arginine/serine rich domains typical of splicing factors.

```
#length 437 #molecular-weight 51556 #checksum 9783
```

DB 12; Score 255; Match 47.6%; QryMatch 2.7%; Pred. No. 8.37e-14; Matches 30; Conservative 13; Mismatches 18; Indels 2; Gaps 2;

Db 219 yderpgpsplphrdrdrdrerer-tersrerdkererrrrsdrtrrrrsrdkeerrrr 277

[illegible]

Db 278 rer 280

100

QY 1303 REK 1305

RESULT	15
--------	----

ENTRY	S28147	#type complete
-------	--------	----------------

TITLE	U1 snRNP 70K protein - Arabidopsis thaliana
<p> 1 2 3 4 5 6 7 8 9 10 11 12 13 14 15 16 17 18 19 20 21 22 23 24 25 26 27 28 29 30 31 32 33 34 35 36 37 38 39 40 41 42 43 44 45 46 47 48 49 50 51 52 53 54 55 56 57 58 59 60 61 62 63 64 65 66 67 68 69 70 71 72 73 74 75 76 77 78 79 80 81 82 83 84 85 86 87 88 89 90 91 92 93 94 95 96 97 98 99 100 101 102 103 104 105 106 107 108 109 110 111 112 113 114 115 116 117 118 119 120 121 122 123 124 125 126 127 128 129 130 131 132 133 134 135 136 137 138 139 140 141 142 143 144 145 146 147 148 149 150 151 152 153 154 155 156 157 158 159 160 161 162 163 164 165 166 167 168 169 170 171 172 173 174 175 176 177 178 179 180 181 182 183 184 185 186 187 188 189 190 191 192 193 194 195 196 197 198 199 200 201 202 203 204 205 206 207 208 209 210 211 212 213 214 215 216 217 218 219 220 221 222 223 224 225 226 227 228 229 230 231 232 233 234 235 236 237 238 239 240 241 242 243 244 245 246 247 248 249 250 251 252 253 254 255 256 257 258 259 260 261 262 263 264 265 266 267 268 269 270 271 272 273 274 275 276 277 278 279 280 281 282 283 284 285 286 287 288 289 290 291 292 293 294 295 296 297 298 299 300 301 302 303 304 305 306 307 308 309 310 311 312 313 314 315 316 317 318 319 320 321 322 323 324 325 326 327 328 329 330 331 332 333 334 335 336 337 338 339 340 341 342 343 344 345 346 347 348 349 350 351 352 353 354 355 356 357 358 359 360 361 362 363 364 365 366 367 368 369 370 371 372 373 374 375 376 377 378 379 380 381 382 383 384 385 386 387 388 389 390 391 392 393 394 395 396 397 398 399 400 401 402 403 404 405 406 407 408 409 410 411 412 413 414 415 416 417 418 419 420 421 422 423 424 425 426 427 428 429 430 431 432 433 434 435 436 437 438 439 440 441 442 443 444 445 446 447 448 449 450 451 452 453 454 455 456 457 458 459 460 461 462 463 464 465 466 467 468 469 470 471 472 473 474 475 476 477 478 479 480 481 482 483 484 485 486 487 488 489 490 491 492 493 494 495 496 497 498 499 500 501 502 503 504 505 506 507 508 509 510 511 512 513 514 515 516 517 518 519 520 521 522 523 524 525 526 527 528 529 530 531 532 533 534 535 536 537 538 539 540 541 542 543 544 545 546 547 548 549 550 551 552 553 554 555 556 557 558 559 560 561 562 563 564 565 566 567 568 569 570 571 572 573 574 575 576 577 578 579 580 581 582 583 584 585 586 587 588 589 590 591 592 593 594 595 596 597 598 599 600 601 602 603 604 605 606 607 608 609 610 611 612 613 614 615 616 617 618 619 620 621 622 623 624 625 626 627 628 629 630 631 632 633 634 635 636 637 638 639 640 641 642 643 644 645 646 647 648 649 650 651 652 653 654 655 656 657 658 659 660 661 662 663 664 665 666 667 668 669 670 671 672 673 674 675 676 677 678 679 680 681 682 683 684 685 686 687 688 689 690 691 692 693 694 695 696 697 698 699 700 701 702 703 704 705 706 707 708 709 710 711 712 713 714 715 716 717 718 719 720 721 722 723 724 725 726 727 728 729 730 731 732 733 734 735 736 737 738 739 740 741 742 743 744 745 746 747 748 749 750 751 752 753 754 755 756 757 758 759 760 761 762 763 764 765 766 767 768 769 770 771 772 773 774 775 776 777 778 779 780 781 782 783 784 785 786 787 788 789 790 791 792 793 794 795 796 797 798 799 800 801 802 803 804 805 806 807 808 809 810 811 812 813 814 815 816 817 818 819 820 821 822 823 824 825 826 827 828 829 830 831 832 833 834 835 836 837 838 839 840 841 842 843 844 845 846 847 848 849 850 851 852 853 854 855 856 857 858 859 860 861 862 863 864 865 866 867 868 869 870 871 872 873 874 875 876 877 878 879 880 881 882 883 884 885 886 887 888 889 890 891 892 893 894 895 896 897 898 899 900 901 902 903 904 905 906 907 908 909 910 911 912 913 914 915 916 917 918 919 920 921 922 923 924 925 926 927 928 929 930 931 932 933 934 935 936 937 938 939 940 941 942 943 944 945 946 947 948 949 950 951 952 953 954 955 956 957 958 959 960 961 962 963 964 965 966 967 968 969 970 971 972 973 974 975 976 977 978 979 980 981 982 983 984 985 986 987 988 989 990 991 992 993 994 995 996 997 998 999 1000 1001 1002 1003 1004 1005 1006 1007 1008 1009 1010 1011 1012 1013 1014 1015 1016 1017 1018 1019 1020 1021 1022 1023 1024 1025 1026 1027 1</p>	

[illegible]

ORGANISM	#formal_name	Arabidopsis thaliana	#common_name	mouse-ear cress
----------	--------------	----------------------	--------------	--------------------

```
DATE      #sequence_revision 20-Feb-1995 #text_change
20-Feb-1995 12-May-1995
```

ACCESSIONS S28147

REFERENCE
S28147

authors
Reddy,

#journal Biochim. Biophys. Acta (1992) 1171:88-92

Cloning of the cDNA for U1 small nuclear

particle 70K protein from *Arabidopsis thaliana*.

accession S28147

```
##status preliminary
```

##molecule type mRNA
1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65
66
67
68
69
70
71
72
73
74
75
76
77
78
79
80
81
82
83
84
85
86
87
88
89
90
91
92
93
94
95
96
97
98
99
100

```
##residues 1-1* 1-303 ##label RED
```

CLASSIFICATION #superfamily ribonucleoprotein repeat homology

FEATURE

33-100

```

SUMMARY
#length 303 #molecular-weight 36430 #checksum 969

```

DB 8; Score 239; Match 43.6%; QrvMatch 2.5%; Pred. No. 5.26e-12;

Matches	34;	Conservative	13;	Mismatches	27;	Indels	4;	Gaps	4;
----------------	------------	---------------------	------------	-------------------	------------	---------------	-----------	-------------	-----------

1. **Introduction**

Db 167 erersrelsh-eaprrersdrpredkhhrrdrdgaardrdrdsrrdrdrtrdradrrdr 225

[illegible]

Ov 1234 DRKSRSYHYDRRRRDEDRDRERDRDRD-RDRDRDRDRDRERSR-ERDRDR 1291

[x] 1 2 3 4 5 6 7 8 9 10 11 12 13 14 15 16 17 18 19 20 21 22 23 24 25 26 27 28 29 30 31 32 33 34 35 36 37 38 39 40 41 42 43 44 45 46 47 48 49 50 51 52 53 54 55 56 57 58 59 60 61 62 63 64 65 66 67 68 69 70 71 72 73 74 75 76 77 78 79 80 81 82 83 84 85 86 87 88 89 90 91 92 93 94 95 96 97 98 99 100 101 102 103 104 105 106 107 108 109 110 111 112 113 114 115 116 117 118 119 120 121 122 123 124 125 126 127 128 129 130 131 132 133 134 135 136 137 138 139 140 141 142 143 144 145 146 147 148 149 150 151 152 153 154 155 156 157 158 159 160 161 162 163 164 165 166 167 168 169 170 171 172 173 174 175 176 177 178 179 180 181 182 183 184 185 186 187 188 189 190 191 192 193 194 195 196 197 198 199 200 201 202 203 204 205 206 207 208 209 210 211 212 213 214 215 216 217 218 219 220 221 222 223 224 225 226 227 228 229 230 231 232 233 234 235 236 237 238 239 240 241 242 243 244 245 246 247 248 249 250 251 252 253 254 255 256 257 258 259 260 261 262 263 264 265 266 267 268 269 270 271 272 273 274 275 276 277 278 279 280 281 282 283 284 285 286 287 288 289 290 291 292 293 294 295 296 297 298 299 300 301 302 303 304 305 306 307 308 309 310 311 312 313 314 315 316 317 318 319 320 321 322 323 324 325 326 327 328 329 330 331 332 333 334 335 336 337 338 339 340 341 342 343 344 345 346 347 348 349 350 351 352 353 354 355 356 357 358 359 360 361 362 363 364 365 366 367 368 369 370 371 372 373 374 375 376 377 378 379 380 381 382 383 384 385 386 387 388 389 390 391 392 393 394 395 396 397 398 399 400 401 402 403 404 405 406 407 408 409 410 411 412 413 414 415 416 417 418 419 420 421 422 423 424 425 426 427 428 429 430 431 432 433 434 435 436 437 438 439 440 441 442 443 444 445 446 447 448 449 450 451 452 453 454 455 456 457 458 459 460 461 462 463 464 465 466 467 468 469 470 471 472 473 474 475 476 477 478 479 480 481 482 483 484 485 486 487 488 489 490 491 492 493 494 495 496 497 498 499 500 501 502 503 504 505 506 507 508 509 510 511 512 513 514 515 516 517 518 519 520 521 522 523 524 525 526 527 528 529 530 531 532 533 534 535 536 537 538 539 540 541 542 543 544 545 546 547 548 549 550 551 552 553 554 555 556 557 558 559 560 561 562 563 564 565 566 567 568 569 570 571 572 573 574 575 576 577 578 579 580 581 582 583 584 585 586 587 588 589 590 591 592 593 594 595 596 597 598 599 600 601 602 603 604 605 606 607 608 609 610 611 612 613 614 615 616 617 618 619 620 621 622 623 624 625 626 627 628 629 630 631 632 633 634 635 636 637 638 639 640 641 642 643 644 645 646 647 648 649 650 651 652 653 654 655 656 657 658 659 660 661 662 663 664 665 666 667 668 669 670 671 672 673 674 675 676 677 678 679 680 681 682 683 684 685 686 687 688 689 690 691 692 693 694 695 696 697 698 699 700 701 702 703 704 705 706 707 708 709 710 711 712 713 714 715 716 717 718 719 720 721 722 723 724 725 726 727 728 729 730 731 732 733 734 735 736 737 738 739 740 741 742 743 744 745 746 747 748 749 750 751 752 753 754 755 756 757 758 759 760 761 762 763 764 765 766 767 768 769 770 771 772 773 774 775 776 777 778 779 780 781 782 783 784 785 786 787 788 789 790 791 792 793 794 795 796 797 798 799 800 801 802 803 804 805 806 807 808 809 810 811 812 813 814 815 816 817 818 819 820 821 822 823 824 825 826 827 828 829 830 831 832 833 834 835 836 837 838 839 840 841 842 843 844 845 846 847 848 849 850 851 852 853 854 855 856 857 858 859 860 861 862 863 864 865 866 867 868 869 870 871 872 873 874 875 876 877 878 879 880 881 882 883 884 885 886 887 888 889 890 891 892 893 894 895 896 897 898 899 900 901 902 903 904 905 906 907 908 909 910 911 912 913 914 915 916 917 918 919 920 921 922 923 924 925 926 927 928 929 930 931 932 933 934 935 936 937 938 939 940 941 942 943 944 945 946 947 948 949 950 951 952 953 954 955 956 957 958 959 960 961 962 963 964 965 966 967 968 969 970 971 972 973 974 975 976 977 978 979 980 981 982 983 984 985 986 987 988 989 990 991 992 993 994 995 996 997 998 999 1000 1001 1002 1003 1004 1005 1006 1007 1008 1009 1010 1011 1012 1013 1014 1015 1016 1017 1018 1019 1020 1021 1022 1023 1024 1025 1026 1027 1028 1029 1030 1031 1032 1033 1034 1035 1036 1037 1038 1039

Db 226 drqrdrtsrhdrrrk 243

1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21	22	23	24	25	26	27	28	29	30	31	32	33	34	35	36	37	38	39	40	41	42	43	44	45	46	47	48	49	50	51	52	53	54	55	56	57	58	59	60	61	62	63	64	65	66	67	68	69	70	71	72	73	74	75	76	77	78	79	80	81	82	83	84	85	86	87	88	89	90	91	92	93	94	95	96	97	98	99	100
---	---	---	---	---	---	---	---	---	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	-----

QV 1292 YRD-ERDHRASPREKRQR 1308

2000

Search completed: Wed

Job time : 169 secs.

PR 26-MAY-1992; US-899651.
PA (BECT) BECTON DICKINSON CO.
PI Shank DD, Spears PA;
DR WPI; 93-378844/48.
PT New oligo:nucleotide probes specific for Mycobacteria - used for
PT detection and amplification of Mycobacteria nucleic acid In
PS samples
PS Claim 3; Page 14; 23pp; English.
CC Oligonucleotide probe MK14-A consists of nucleotides 5-95 of MK14
CC (Q51735). It hybridized to all spp. of mycobacteria tested, but
CC cross reacted to a few non-mycobacterial spp. The probe may
CC be useful as an initial screen for mycobacterial infection.
CC See also Q51735-45 and Q51747-59.
SQ Sequence 91 BP; 5 A; 17 C; 15 G; 4 T;

DB 9; Score 43; Match 2.0%; QryMatch 1.0%; Pred.No. 1.33e-10;
Matches 1; Conservative 46; Mismatches 4; Indels 0; Gaps 0;

Ddb 11 svshyyvvhvhhshvhhvhvsvvvhvhhvhhvhvhhvhyrvyvc 61
:: :::::::::::::::::::: :
Qy 227 CGAGGCTCTTCGAATCTCGAATAAGCGTCAACAGAGGCCCTCGC 277

RESULT 7

ID Q43704 standard; DNA; 756 BP.

AC Q43704;

DT 29-SEP-1993 (first entry)

DE Sequence encoding a new cytokine which inhibits induction by gamma

DE interferon of expression of Class II histocompatibility antigens.

KW Cytokine; interferon-gamma antagonist; autoimmune disease therapy;

KW transplants; Class II histocompatibility antigens;

KW gamma interferon; leukaemia line K562; ss.

OS Homo sapiens.

FS Key

OH Location/Qualifiers

FT Key

FT CDS 40..528

FT /*tag= a

FP W09311232-A.

PN 10-JUN-1993.

PD 02-DEC-1992; F01123.

PR 02-DEC-1991; FR-014908.

PP (INRM) INSERM INST NAT SANTE & RECH MED.

PA Augery-bourget Y, Azzarone B, Brecheix C, Jasmin C;

PI Brief PH;

DR WPI; 93-197051/24.

DR P-PSDB; R37991.

PT New cytokine as interferon-gamma antagonist - inhibits induction

PT of class II histocompatibility antigens on cell surface by

PT IFN-gamma, for treating and preventing auto-immune disorders

PS Claim 4; Fig 1; 44pp; French.

CC The new cytokine specifically inhibits induction by gamma interferon

CC of expression of Class II histocompatibility antigens on cell

CC surfaces. It is naturally secreted by cells which are difficult or

CC impossible to induce with gamma-interferon, e.g. the leukaemia line

CC K562, from which it can be isolated. To isolate DNA, a K562 gene

CC bank was constructed in lambda gt10 phage, used to transform Y1090

CC bacteria and these screened for reactivity with MAbs specific for

CC the cytokine.Positive colonies were purified,their inserts isolated

CC with EcoRI and subcloned into pBlSCR KS. Recombinant plasmids were

CC used to transform HB101 cells and the insert from one positive clone

CC sequenced. This insert was recloned in vector CDM8 and recombinants

CC used to transform COS cells. The transformants expressed functional

CC recombinant cytokine.

SQ Sequence 756 BP; 254 A; 136 C; 241 G; 125 T

DB	7;	Score	37;	Match	68.0%;	QryMatch	0.8%;	Pred.	No. 3.59e-07;
Matches	70;	Conservative	0;	Mismatches	33;	Indels	0;	Gaps	0;
Db	116	atcgggaacgcagcgtgacggaagagacagagaccgtgaccgagagcagagcagag 175							
Qy	3940	ATCGTCGAGAGGATCCGGATGAGATAGGATCGAGACCGCTGAAAGGACAGAGATCGCG 3999							
Db	176	aacgagatcggaacgagagcagcgagcgagcgagaccgagagagaqa 218							
Qy	4000	ACAGGGATCGGGATAGGGATCGTGCACCGGACAGGGATAGGCA 4042							
RESULT	8								
ID	Q58032	standard; DNA; 4752 BP.							
AC	Q58032;								
DT	01-AUG-1994	(first entry)							
DE	DE Tyrosinase promoter.								
KW	Tyrosinase; cytokine; tumor therapy; gene therapy; melanoma;								
KW	B16; tumor-specific promoter; polymerase chain reaction; PCR;								
KW	B16; primer; ds.								
OS	Mus sp.								
PN	W09404196-A.								
PD	03-MAR-1994.								
PF	16-AUG-1993; G01730.								
PR	14-AUG-1992; GB-017270.								
PR	27-FEB-1993; GB-004024.								
PA	(IMCR) IMPERIAL CANCER RES TECHNOLOGY.								
PI	Hart IR, Vile RG;								
DI	WPI; 94-082848/10.								
PT	DNA construct encoding cytokine for expression in tumour cells -								
PT	for treating melanoma, pancreatic, breast, colon, prostate cancer								
PS	Disclosure; Page 67-70; 107pp; English.								
CC	Cytokine genes are expressed in tumor cells, especially melanoma								
CC	cells, under the control of the tumor-specific tyrosinase gene								
CC	promoter, for gene therapy. A DNA fragment from the 5' end of								
CC	the tyrosinase gene was generated by PCR from genomic DNA of the								
CC	B16 melanoma line using the primers given in Q58041-46.								
SQ	Sequence 4752 BP; 1550 A; 854 C; 883 G; 1465 T;								
DB	10;	Score	33;	Match	64.6%;	QryMatch	0.7%;	Pred.	No. 5.52e-05;
Matches	73;	Conservative	0;	Mismatches	40;	Indels	0;	Gaps	0;
Db	1799	gaaggag 1858							
Qy	3957	GATCAAGATAGGGATCCAGCCGCTGAAAGGGACAGAGATCCGCACAGCGGATCGGATAGG 4016							
Db	1859	gag 1911							
Qy	4017	GATCGTGACCGGGACAGGATAGGATAGAGAACGATCGAGAGAACGACAG 4069							
RESULT	9								
ID	Q58005	standard; DNA; 3461 BP.							
AC	Q58005;								
DT	14-SEP-1994	(first entry)							
DE	Sequence of plasmid pTHN1 showing the translation elongation factor								
DE	(tefl) promoter and coding sequence.								
KW	promoter; translation elongation factor; pTHN1; ss.								
OS	Trichoderma reesei.								
FT	Key	Location/Qualifiers							
FT	promoter	1..1234							
FT	/*tag= a								
FT	CDS	1235..1237							
FT	/*tag= b								

FT /label= start codon
PN W09404673-A.
PD 03-MAR-1994.
PR 19-AUG-1993; F10330.
PR 19-AUG-1992; US-932485.
PA (AIKO-) AIKO OY AB.
PI Ilmen MH, Nakari TH, Nevalainen KMH, Onnela M, Penttilae ME;
DR WPI; 94-083192/10.
PT Cloning promoters active in a desired environmental condition -
PT used partic. for expression of genes in Trichoderma fungal hosts
PT in glucose-contg. medium
PS Claim 15; Figure 1A; 120pp; English.
CC An assay was undertaken to isolate Trichoderma reesei genes which
CC are strongly expressed on glucose. The cDNAs of clones cDNA33,
CC cDNA1, cDNA10, cDNA12 and cDNA 15 were used as probes to isolate
CC corresp. genes and promoters from a Trichoderma chromosomal lambda-
CC bank. The promoters and either the 5' parts of the chromosomal genes
CC or the whole genes were subcloned into pSP73 vector yielding the
CC plasmids pTHN1, pEA33, pTHN3, pEA10, pEA12 and pEA155, corresp. to
CC the clones cDNA33, cDNA1, cDNA10, cDNA12 and cDNA15, respectively.
CC Sequences were obtd. from the 5' ends of the genes and from the
CC promoters using primers designed from previously obtd. sequences.
CC The sequences of the isolated promoters and genes or parts of them
CC (either obtd. from cDNA or chromosomal DNA) are shown in Q58005 for
CC cDNA33, Q58006 for cDNA1, Q58007 for cDNA10, Q58008 for cDNA12,
CC and Q58009 for cDNA15. Based on sequence similarity to known
CC sequences in a protein data bank the clone cDNA33 could be
CC identified as a translation elongation factor, TEF1-alpha.
SQ Sequence 3461 BP; 850 A; 1044 C; 860 G; 697 T;

DB 10; Score 33; Match 82.4%; QryMatch 0.7%; Pred. No. 5.52e-05;
Matches 42; Conservative 0; Mismatches 9; Indels 0; Gaps 0;

Db 608 gatcgagaggatcactacagagagagcagcgagcgagggagggatcgagat 658
||||||| ||||||| ||||||| ||||||| ||||||| ||||||| ||||||| |||
QY 3993 GATCGGACGAGGATCGGGATAGGATCTCTGACGGGACGAGATAGGAT 4043

RESULT 10
ID Q23445 standard; DNA; 3699 BP.
AC Q23445;
DT 17-AUG-1992 (first entry)
DE Vector pCpE.
KW Mouse; heavy chain; cloning; ss.
OS Rattus rattus.
PN W09203918-A.
PD 19-MAR-1992.
PF 28-AUG-1991; U06185.
PR 29-AUG-1990; US-574748.
PR 31-AUG-1990; US-575962.
PA (GENP-) GENPHARM INT INC.
PI Lonberg N, Kay R;
DR WPI; 92-113962/14.
PT Immunoglobulin trans:genes - for prodn. of heterologous
PT non-rearranged and/or rearranged Ig chains
PS Example 14; Page 81; 172pp; English.
CC The vector sequence was obtd. by PCR amplification of the
CC immunoglobulin heavy chain 3' enhancer (Patterson, et al., (1990)
CC Nature, 344: 165-168) from rat liver DNA using the PCR primers
CC oligo-44 and oligo-45 (see Q23446,7). The amplified product
CC was digested by BamHI and SphI and cloned into a pUC derived
CC plasmid contg. a polylinker (pN03). The resulting plasmid,
CC pRE3, was digested with BamHI and HindIII and the insert contg.
CC the rat Ig heavy chain 3' enhancer cloned into pCp1b. The resulting

CC plasmid, pCpE contains several unique restriction sites into which
CC sequences can be cloned and subsequently excised together with the
CC 3' enhancer by NotI digestion, allowing for cloning of very large
CC DNA sequences.
CC See also Q23419-50, Q22417-30.
SQ Sequence 3699 BP; 882 A; 995 C; 899 G; 923 T;

DB 3; Score 30; Match 63.9%; QryMatch 0.7%; Pred. No. 2.07e-03;
Matches 69; Conservative 0; Mismatches 39; Indels 0; Gaps 0;

Db 122 ctgtctctgtctctctgtctctgtctgtctgtctgtctctctctctctctct 181
||||||| ||||||| ||||||| ||||||| ||||||| ||||||| ||||||| |||
Cp 4069 CTGCTCTGCTTCTCTCGATCGTTCCTATCCCTATCCCTGTCGCCGTCACGATCCCTATCC 4010

Db 182 gtctctctgtctctctgtctctgtctctgtctctgtctctctctctctct 229
|| ||||| || ||||||| || ||||||| || ||||||| || ||||||| || |||||
Cp 4009 CGATCCCTGTCCGATCTCTGCTCCCTTTTCAGGCTCGATCCCTATCT 3962

RESULT 11
ID Q31726 standard; DNA; 1577 BP.
AC Q31726;
DT 04-APR-1993 (first entry)
DE Rat IgH 3'-enhancer.
KW Mouse; cross hybridisation; B cell specific; target; ss.
OS Rattus rattus.
PN W09221762-A.
PD 10-DEC-1992.
PF 03-JUN-1992; SE0375.
PR 07-JUN-1991; SE-001740.
PA (PETT/) PETERSSON S.
PI Pettersson S;
DR WPI; 92-433660/52.
PT Mouse immunoglobulin H 3' enhancer - which cross-hybridises with
PT rat 3' enhancer and is used to target tissue-specific gene
PT expression
PS Claim 3; Fig 3B 13; 33pp; English.
CC A mouse liver library in bacteriophage was screened using as a probe
CC an AccI-BglI subfragment from the core of the rat 3' enhancer
CC (nucleotides 432-748). Two overlapping gps. of clones (lambda M2 and
CC lambda M3) were isolated. Phage lambda M3 extends 3' of lambda M2.
CC The sequence of the mouse IgH 3' enhancer was determined and aligned
CC with that of the rat. Hybridisations of subclones of the mouse IgH
CC 3' enhancer to phage lambda M2 showed that the 3' enhancers of mouse
CC and rat were present in the genome in opposite orientations. The
CC enhancers are B cell specific and may be used to target tissue
CC specific expression of genes, so may be useful therapeutically, e.g.
CC in targeting prodn. of proteins and in hybridoma technology. The
CC enhancer may be used to enhance expression of genes in host cells, in
CC vivo or in vitro, partic. in certain lymphoid cell lines and in
CC transgenic animals. It may be used for the prodn. of monoclonal
CC antibodies. See also Q31725.
SQ Sequence 1577 BP; 406 A; 402 C; 456 G; 313 T;

DB 5; Score 30; Match 63.9%; QryMatch 0.7%; Pred. No. 2.07e-03;
Matches 69; Conservative 0; Mismatches 39; Indels 0; Gaps 0;

Db 837 agagagacagacagacagacagacagacagacagacagacagacagagagac 896
||||||| ||||||| ||||||| ||||||| ||||||| ||||||| ||||||| |||
QY 3962 AGATAGGATCGAGACCCCTGAAAGGGACAGAGATCCGGACAGGATCGGGATCGGATCG 4021

Db 897 agagagacacacacagacagacagacagacagacagacagacagacagacag 944
|| ||||| || ||||||| || ||||||| || ||||||| || ||||||| || |||||
QY 4022 TGACCGGACAGGGATAGGGATAGAGAACGATCGAGAGAACGACAG 4069

```
RESULT 12
ID N71302 standard; DNA; 3871 BP.
AC N71302;
DT 30-APR-1991 (first entry)
DE HSV-1 gB and surrounding regions.
KW Vaccine; prophylaxis; treatment; Herpes Simplex Virus-1;
OS glycoprotein; gB; ss.
FH Key
  Location/Qualifiers
FT misc_RNA
  1..375
FT /tag= a
FT /note= "5' extra sequences beginning with the XhoI
FT site"
FT CAAT_signal
  406..410
FT /tag= b
FT /number= 1
FT CAAT_signal
  443..448
FT /tag= c
FT /number= 2
FT TATA_signal
  476..479
FT /tag= d
FT misc_RNA
  501..789
FT /tag= e
FT /label= mRNA start sequence
FT /note= "501 is possible start site"
FT misc_RNA
  504..789
FT /tag= f
FT /label= mRNA start sequence
FT /note= "504 is possible start site"
FT misc_RNA
  506..789
FT /tag= g
FT /label= mRNA start sequence
FT /note= "506 is possible start site"
FT CDS
  790..3498
FT /tag= h
FT /label= HSV-1 gB
FT /note= "includes N-terminal hydrophobic leader and
FT a membrane-spanning sequence, a C-terminal
FT ionic sequence, and 9 N-linked
FT saccharide-addition sites"
FT 3'UTR
  3499..3549
FT /tag= i
FT polyA_signal
  3518..3525
FT /tag= j
FT polyA_site
  3549..3549
FT /tag= k
FT misc_RNA
  3549..3997
FT /tag= l
FT /note= "3' nonessential sequences to the BamHI
FT site"
PN US4642333-A.
PD 10-FEB-1987.
PF 20-JUN-1984; 622496.
PR 16-SEP-1983; US-532996.
PR 20-JUN-1985; US-622496.
PA (PERS/) PERSON S.
PI Person S;
DR WPI; 87-056354/08.
PT Amino acid chain of glyco:protein B of HSV-1 and 2 - prepd. as
PT recombinant and used for vaccines for herpes simplex virus types 1
PT and 2.
PS Example; Table 1; 16pp; English.
```

```
CC !NOTE! This sequence has been indexed as represented in the
CC specification, except that bases 'E' have been replaced by 'N'.
CC The features have been indexed according to the legend of table 1 on
CC column 19/20 and the Sequence Summary of column 5 (sic). Note that
CC the base numbering of the features does not correspond to the
CC the sequence numbering below.
CC For another DNA sequence of HSB-1 gB see N71303 (P71135),
CC and for HSV-2 gB see N71399 (P71136).
CC A pure non-glycosylated amino acid (AA) chain comprising a sequence
CC corresponding to that occurring in glycoprotein B of HSV-1 or HSV-2
CC virus which is antigenic to HSV-1 of HSV-2, which contains no more
CC than 750 AA residues, and which includes AA residues 135-649
CC inclusive is claimed. It can be used to produce vaccines for
CC prophylaxis and treatment of HSV-1 and HSV-2.
SQ Sequence 3871 BP; 743 A; 1402 C; 544 G; 754 T;

DB 2; Score 29; Match 39.6%; QryMatch 0.7%; Pred. No. 6.68e-03;
Matches 53; Conservative 31; Mismatches 49; Indels 1; Gaps 1;

Db 2817 bcaccatcacccatcacgbbtcacacbcgcacbccacbccatattcbbcbbbbcb 2876
:||||| ||||| ||||| ||||| ||||| ||||| ||||| ||||| |||||
Cp 4042 TCCCTATCCCTGTCGGGTACCGATCCCTATCCCGATCCCTGTCGGCATCTCTGCTT 3983
:||||| ||||| ||||| ||||| ||||| ||||| ||||| |||||
Db 2877 bobctttctcabbgatbbgcacctbqggcgcggtcb-bcaaggtbbtgatbbcat 2935
:||||| ||||| ||||| ||||| ||||| ||||| ||||| |||||
Cp 3982 TCACGGTCTCGATCCCTATCTTCATCCGGATCCCTTCGACGATCATAGTAGAGAA 3923
:||||| ||||| ||||| ||||| ||||| ||||| ||||| |||||
Db 2936 cbtbbcggtatcbb 2949
:||||| |||||
Cp 3922 CGTCGGATTTCTCT 3909

RESULT 13
ID Q44179 standard; DNA; 3699 BP.
AC Q44179;
DT 10-NOV-1993 (first entry)
DE Plasmid pCPE for cloning large Ig transgenes.
KW Immunoglobulin; rat Ig heavy chain 3' enhancer; cloning vector;
KW heavy chain minilocus transgene; ss.
OS Synthetic.
FH Key Location/Qualifiers
FT enhancer
  1..665
FT /tag= a
FT /note= "rat Ig heavy chain 3' enhancer inserted
FT between the BamHI and HindIII sites of
FT the pCPE polylinker. The 8 Others in this
FT region correspond to nucleotides which were
FT illegible in the printed specification"
PN W0931227-A.
PD 24-JUN-1993.
PF 17-DEC-1992; U10983.
PR 17-DEC-1991; US-810279.
PR 18-MAR-1992; US-853408.
PR 23-JUN-1992; US-904068.
PA (GENP-) GENPHARM INT INC.
PI Kay RM, Lonberg N;
DR WPI; 93-214169/26.
PT Transgenic non-human animals contg. immunoglobulin heavy chain
PT trans gene - used to produce useful antibodies by isotype
PT switching
PS Example 12; Page 90; 196pp; English.
CC Plasmid pCPE1a was derived from pBR322 by insertion of an EcoRI-StyI
CC linker. The plasmid contains a NotI restriction site downstream
CC (relative to the ampicillin resistance gene) of a strong
```

CC transcription termination signal derived from the trpA gene.
CC Potential toxicity of the plasmid is reduced by being of low copy
CC number (due to retaining the pBR322 copy number control region) and
CC by eliminating readthrough transcription from the Amp^r gene (due to
CC the strong terminator). Plasmid pGP1b was derived from pGP1a by
CC insertion of a short polylinker region, flanked by NotI sites.
CC Oligonucleotides "oligo-44" and "oligo-45" (Q44177 and Q44178,
CC respectively) were used to amplify the immunoglobulin heavy chain
CC 3' enhancer from rat liver DNA by the polymerase chain reaction
CC technique. The amplified product was subcloned and an insert
CC containing the enhancer was cloned into BamHI/HindIII digested
CC pGP1b. The resulting plasmid pCPE (i.e. Q44179) contains several
CC unique restriction sites into which large transgene sequences can be
CC cloned and subsequently excised together with the 3' enhancer.
SQ Sequence 3699 BP; 882 A; 992 G; 897 C; 920 T;

DB 7; Score 29; Match 63.0%; QryMatch 0.7%; Pred. No. 6.68e-03;
Matches 68; Conservative 0; Mismatches 40; Indels 0; Gaps 0;

Db 122 ctgtctctgtctctgtctctgtctctgtctgtgtgtgtctctctctctctctctctct 181
||||| ||||| ||||| ||||| ||||| ||||| ||||| ||||| ||||| ||||| |||||

Cp 4069 CTGTCGTTCTCGATCGTCTCTATCCGATCCCTGTCGGGTCAGATCCCTATCC 4010

Db 182 gtctctgtctctgtctctgtctctgtctctgtctctgtctctctctctctctctct 229
||||| ||||| ||||| ||||| ||||| ||||| ||||| ||||| ||||| ||||| |||||

Cp 4009 CGATCCGTCGGGATCTCTGTGTCCTTCAGGCTTCGATCCCTATCT 3962

RESULT 14

ID Q57506 standard; DNA; 700 BP.

AC Q57506;

DT 09-JUL-1994 (first entry)

DE Rat GAP-43 promoter.

KW GAP-43; internal regulatory protein; IRP; recombinant; neuron;

KW neuronal growth; promoter; *es*.

OS Rattus sp.

FH Key Location/Qualifiers

FT promoter 1..600

FT /*tag= a

FT /note= "strong potential for H-DNA conformation"

FT misc_binding 82..89

FT /*tag= b

FT /note= "consensus Pit-1 binding site"

FT misc_signal 519

FT /*tag= c

FT /note= "transcriptional start site"

FT misc_signal 545

FT /*tag= d

FT /note= "transcriptional start site"

FT misc_signal 546

FT /*tag= e

FT /note= "transcriptional start site"

FT misc_signal 549

FT /*tag= f

FT /note= "transcriptional start site"

FT misc_signal 550

FT /*tag= g

FT /note= "transcriptional start site"

FT CDS 601..700

FT /*tag= h

FT /note= "5' end of GAP-43 gene"

PN AU9347435-A.

PD 23-DEC-1993.

PF 17-SEP-1993; 047435.

PR 22-DEC-1989; US-465635.

PR 02-JUL-1990; US-546453.

PA (FEDE/) FEDEROFF H J.

PA (FISH/) FISHMAN M C.

PA (STRI/) STRITTMATTER SM.

PA (VALE/) VALENZUELA D.

PA (ZUBE/) ZUBER M X.

PI Federoff HJ, Fishman MC, Strittmatter SM, Valenzuela D;

PI Zuber MX;

DR WPI; 94-049278/07.

PT Recombinant mammalian GAP-43 protein - used to monitor and

PT regulate neuronal growth in animals, pref. humans

PS Disclosure; Fig 14; 156pp; English.

CC The nucleotide sequence of the rat GAP-43 gene promoter region is

CC given in Q57506.

SQ Sequence 700 BP; 230 A; 87 C; 278 G; 105 T;

DB 9; Score 27; Match 65.5%; QryMatch 0.6%; Pred. No. 6.58e-02;
Matches 57; Conservative 0; Mismatches 30; Indels 0; Gaps 0;

Db 280 ggga 339
||||| ||||| ||||| ||||| ||||| ||||| ||||| ||||| ||||| ||||| |||||

Qy 3985 GGGACAGAGATCCGACAGGATCGGGATAGGATCTGACCGGACAGGATAGGATA 4044

Db 340 ga 366
||||| ||||| ||||| ||||| ||||| ||||| ||||| ||||| ||||| |||||

Qy 4045 GAGAAGCATCGACAGACGACAGCAGGC 4071

RESULT 15

ID Q56643 standard; DNA; 3586 BP.

AC Q56643;

DT 16-SEP-1994 (first entry)

DE Human tyrosinase gene.

KW Tyrosinase; albino; albinism; detection; diagnosis;

KW treatment; prevention; *es*.

OS Homo sapiens.

FH Key Location/Qualifiers

FT CDS 987..1805

FT /*tag= a

FT /product= Tyrosinase.

FT /note= "CDS excludes termination codon."

PN J06000100-A.

PD 11-JAN-1994.

PF 18-JUN-1992; 185885.

PR 18-JUN-1992; JP-185885.

PA (DAIL) DAICEL CHEM IND LTD.

DR WPI; 94-077141/10.

DR P-PSDB; R48368.

PT A DNA sequence which hybridises to the human tyrosinase gene -

PT used for the detection, diagnosis, prevention and treatment of

PT human albinism

PS Disclosure; Page 5-7; 8pp; Japanese.

CC A DNA sequence having at least 12 successive bases and which

CC hybridises upstream and/or downstream of the 310th base from the

CC transcription start site of the human tyrosinase gene can be

CC used for the detection, diagnosis, prevention and treatment of human

CC albinism.

SQ Sequence 3586 BP; 1127 A; 636 C; 760 G; 1063 T;

DB 10; Score 27; Match 65.2%; QryMatch 0.6%; Pred. No. 6.58e-02;
Matches 58; Conservative 0; Mismatches 31; Indels 0; Gaps 0;

Db 177 gagagagga 236

Jan 17 18:06

US-08-319-745-3.jpg

17

3981 GAAAGGCACAGATCCGCACAGGGATCGGGATAGCGGATCCTGACCCGGGACACGGGATAGG 4040

D_b 237 gagagagaaagagagagagagagagagag 265

4041 GATAGACAACGATCGAGAGAACGAGACAG 4069 QY

Search completed: Wed Jan 17 18:13:03 1996

Job time : 251 secs.

MA J-X; CHAO J; CHAO L
DEP. BIOCHEMISTRY, MOLECULAR BIOLOGY, MED. UNIV. S.C., 171 ASHLEY AVENUE,
CHARLESTON, S.C. 29425, USA.
BIOCHEMISTRY 31 (44). 1992. 10922-10928. CODEN: BICHA
Full Journal Title: Biochemistry
Language: ENGLISH

We have cloned and determined the nucleotide sequence of a novel kallikrein-like mRNA, designated rKlk10*, from rat submandibular gland and kidney with the aid of the polymerase chain reaction (PCR). This cDNA contains 737 base pairs comprising the sequence encoding a mature protein of 235 amino acid residues, partial zymogen peptide, and 3' noncoding sequence. Sequence comparisons showed that rKlk10 mRNA shares 87 and 88% sequence identity with rat tissue kallikrein at nucleic acid and amino acid levels, respectively. It encodes a 26 428-Da acidic protein whose derived amino acid sequence matches completely with the partial amino acid sequence of a kallikrein-like enzyme designated as T-kininogenase, K10 protein, or antigen- γ . purified from rat submandibular gland [Xiong et al. (1990) J. Biol. Chem. 265, 2822-2827; Gutman et al. (1991) Eur. J. Biochem. 784, 1-5; Berg et al. (1991) Biochem. J. 280, 19-25]. The protein encoded by rKlk10 retains the key amino acid residues determining kallikrein cleavage specificity. Northern blot analysis with an rKlk10-specific oligonucleotide probe showed that its mRNA level in the submandibular gland is decreased dramatically by administration of the β -agonist isoproterenol. Tissue-specific expression of rKlk10 was analyzed by Northern blotting and Southern blotting of PCR-amplified cDNA, which showed that rKlk10 is expressed at high levels in the submandibular gland and low levels in the kidney but not in seven other tissues including prostate, liver, heart, adrenal gland, testes, pituitary, and pancreas. rKlk10 cDNAs cloned from the kidney and submandibular gland show sequence identity. Specific expression of rKlk10 in the kidney in addition to the submandibular gland indicates that rKlk10 may be involved in the regulation of renal function. The PCR-based cloning strategy provided an efficient and reliable way to potentially identify all expressed kallikrein-related genes in rat submandibular gland.

10/7/33 (Item 6 from file: 55)
DIALOG(R) File 55:BIOSIS PREVIEWS(R)
(c) 1996 BIOSIS. All rts. reserv.

7883667 BIOSIS Number: 40084667

PCR-BASED CLONING STRATEGY FOR G PROTEIN-COUPLED RECEPTORS
PARMENTIER M; LIBERT F; LEFORT A; PERRET J; GERARD C; MAENHAUT C; VAN
SANDE J; MOLLEREAU C; EGGERICKX D; ET AL

I.R.I.B.H.N., ULB CAMPUS ERASME, FREE UNIV. BRUSSELS, BRUSSELS, BELG.
SYMPOSIUM ON GENE REGULATION AND SIGNALLING IN ENDOCRINE SYSTEMS HELD AT
THE 20TH ANNUAL MEETING OF THE KEYSTONE SYMPOSIA ON MOLECULAR AND CELLULAR
BIOLOGY, STEAMBOAT SPRINGS, COLORADO, USA, JANUARY 19-25, 1991. J CELL
BIOCHEM SUPPL 0 (15 PART B). 1991. 262. CODEN: JCBSD
Language: ENGLIS

File 154:MEDLINE(R) 1985-1996/Jan W4
(c) format only 1996 Knight-Ridder Info
File 55:BIOSIS PREVIEWS(R) 1985-1995/Jan W1
(c) 1996 BIOSIS

08/319,745

search cloning by PCR

Set Items Description

?e au=scott m

Ref	Items	Index-term
E1	1	AU=SCOTT LV
E2	10	AU=SCOTT LW
E3	338	*AU=SCOTT M
E4	30	AU=SCOTT M A
E5	3	AU=SCOTT M B
E6	48	AU=SCOTT M C
E7	43	AU=SCOTT M D
E8	3	AU=SCOTT M D S
E9	79	AU=SCOTT M E
E10	5	AU=SCOTT M F
E11	104	AU=SCOTT M G
E12	3	AU=SCOTT M G B

Enter P or PAGE for more

?p

Ref	Items	Index-term
E13	28	AU=SCOTT M H
E14	1	AU=SCOTT M I H
E15	56	AU=SCOTT M J
E16	2	AU=SCOTT M J III
E17	2	AU=SCOTT M J JR
E18	5	AU=SCOTT M J L
E19	18	AU=SCOTT M K
E20	83	AU=SCOTT M L
E21	7	AU=SCOTT M M
E22	2	AU=SCOTT M N
E23	15	AU=SCOTT M O
E24	96	AU=SCOTT M P

Enter P or PAGE for more

?p

Ref	Items	Index-term
E25	4	AU=SCOTT M R
E26	10	AU=SCOTT M R D
E27	11	AU=SCOTT M S
E28	11	AU=SCOTT M T
E29	1	AU=SCOTT M-O
E30	16	AU=SCOTT MA
E31	1	AU=SCOTT MB
E32	34	AU=SCOTT MC
E33	27	AU=SCOTT MD
E34	41	AU=SCOTT ME

E35 64 AU=SCOTT MG
E36 24 AU=SCOTT MH

Enter P or PAGE for more

?p

Ref	Items	Index-term
E37	2	AU=SCOTT MI
E38	3	AU=SCOTT MILLAR RN
E39	31	AU=SCOTT MJ
E40	8	AU=SCOTT MJ JR
E41	4	AU=SCOTT MJ 3D
E42	3	AU=SCOTT MJ 3RD
E43	6	AU=SCOTT MK
E44	44	AU=SCOTT ML
E45	4	AU=SCOTT MM
E46	10	AU=SCOTT MO
E47	1	AU=SCOTT MOTOR NEURON DIS RES GROUP
E48	63	AU=SCOTT MP

Enter P or PAGE for more

?s e3 or e24 or e48

338 AU=SCOTT M
96 AU=SCOTT M P
63 AU=SCOTT MP

S1 497 AU="SCOTT M" OR AU="SCOTT M P" OR AU="SCOTT MP"
?s s1 and patched

497 S1
388 PATCHED
S2 6 S1 AND PATCHED

?rd

...completed examining records
S3 4 RD (unique items)
?t s3/6/1-4

3/6/1 (Item 1 from file: 154)
08984026 94299026
patched overexpression causes loss of wingless expression in Drosophila embryos.

3/6/2 (Item 2 from file: 154)
07151658 90058658
The Drosophila patched gene encodes a putative membrane protein required for segmental patterning.

3/6/3 (Item 1 from file: 55)
7236094 BIOSIS Number: 38016615
GENES THAT CONTROL PATTERN FORMATION DURING DEVELOPMENT

3/6/4 (Item 2 from file: 55)

6834732 BIOSIS Number: 37029111

THE DROSOPHILA SEGMENT POLARITY GENE PATCHED ENCODES A MEMBRANE PROTEIN
?s patched

S4 388 PATCHED
?s s4 and (human or mouse or mosquito or butterfly or beetle)

388 S4
5018157 HUMAN
318853 MOUSE
8288 MOSQUITO
2310 BUTTERFLY
7440 BEETLE

S5 157 S4 AND (HUMAN OR MOUSE OR MOSQUITO OR BUTTERFLY OR
BEETLE)
?s s5 and (gene? or clone? or DNA?)

Processing

157 S5
1429785 GENE?
156776 CLONE?
538652 DNA?

S6 14 S5 AND (GENE? OR CLONE? OR DNA?)
?rd

...completed examining records

S7 11 RD (unique items)
?t s7/6/1-11

7/6/1 (Item 1 from file: 154)
09365985 95295985

Human recombinant IGF-I induces the functional expression of AMPA/kainate receptors in cerebellar granule cells.

7/6/2 (Item 2 from file: 154)
09183708 95113708

Surface distribution and partition during freeze-fracture of CD8 antigens on human lymphocytes and on epithelial transfected cells.

7/6/3 (Item 3 from file: 154)
07784014 91303014

Expression and capping of a proliferation-associated surface membrane p34 kDa antigen on different human hematopoietic cell lines.

7/6/4 (Item 4 from file: 154)
07656880 91175880

Lateral diffusion of nerve growth factor receptor: modulation by ligand-binding and cell-associated factors.

7/6/5 (Item 5 from file: 154)
07151658 90058658

The Drosophila patched gene encodes a putative membrane protein required for segmental patterning.

7/6/6 (Item 6 from file: 154)
06796384 89098384
Inhibition of human immunodeficiency virus (HIV-1) replication by
synthetic oligo-RNA derivatives.

7/6/7 (Item 7 from file: 154)
05484758 85100758
A kinetic study of membrane immunoglobulin capping by flow cytometry.

7/6/8 (Item 8 from file: 154)
05424908 85040908
Popliteal vein pseudoaneurysm: a case report.

7/6/9 (Item 1 from file: 55)
10525940 BIOSIS Number: 96125940
INTEGRIN-MEDIATED NEURITE OUTGROWTH IN NEUROBLASTOMA CELLS DEPENDS ON THE
ACTIVATION OF POTASSIUM CHANNELS

7/6/10 (Item 2 from file: 55)
5778744 BIOSIS Number: 83041051
A METHOD OF MEASURING HUMAN AORTIC VOLUME PULSE WAVE VELOCITY BY
ELECTRICAL IMPEDANCE PLETHYSMOGRAPHY AND ITS CLINICAL APPLICATION

7/6/11 (Item 3 from file: 55)
5407358 BIOSIS Number: 82052161
CONSIDERATIONS ON THE TROPHIC NICHES OF TAWNY OWL STRIX-ALUCO AND BARN
OWL TYTO-ALBA
?t s7/7/5

7/7/5 (Item 5 from file: 154)
DIALOG(R)File 154:MEDLINE(R)
(c) format only 1996 Knight-Ridder Info. All rts. reserv.

07151658 90058658
The Drosophila patched gene encodes a putative membrane protein required
for segmental patterning.
Hooper JE; Scott MP
Howard Hughes Medical Institute, University of Colorado, Boulder
80309-0347.

Cell (UNITED STATES) Nov 17 1989, 59 (4) p751-65, ISSN 0092-8674
Journal Code: CQ4
Contract/Grant No.: HD24584; F32HD06784
Languages: ENGLISH
Document type: JOURNAL ARTICLE

The patched (ptc) gene is one of several segment polarity genes required
for correct patterning within every segment of Drosophila. The absence of
ptc gene function causes a transformation of the fate of cells in the
middle part of each segment so that they form pattern elements
characteristic of cells positioned around the segment border. Analysis of
the mutant phenotype demonstrates that both segment and parasegment borders

are included in the duplicated pattern of ptc mutants. We have cloned the ptc gene and deduced that the product is a 1286 amino acid protein with at least seven putative transmembrane alpha helices. ptc RNA is expressed in embryos in broad stripes of segmental periodicity that later split into two stripes per segment primordium. The pattern of expression does not directly predict the transformation seen in ptc mutant embryos, suggesting that ptc participates in cell interactions that establish pattern within the segment.

?s drosophila and (gal or galactosidase)

133 DROSPHILA

7514 GAL

16555 GALACTOSIDASE

S8 4 DROSPHILA AND (GAL OR GALACTOSIDASE)

?s drosophila and (gal or galactosidase)

35618 DROSOPHILA

7514 GAL

16555 GALACTOSIDASE

S9 502 DROSOPHILA AND (GAL OR GALACTOSIDASE)

?s s9 and embryo?

502 S9

163692 EMBRYO?

S10 210 S9 AND EMBRYO?

?s s10a nd develop?

S11 0 S10A ND DEVELOP?

?s s10 and develop?

210 S10

993699 DEVELOP?

S12 125 S10 AND DEVELOP?

?ds

Set	Items	Description
S1	497	AU="SCOTT M" OR AU="SCOTT M P" OR AU="SCOTT MP"
S2	6	S1 AND PATCHED
S3	4	RD (unique items)
S4	388	PATCHED
S5	157	S4 AND (HUMAN OR MOUSE OR MOSQUITO OR BUTTERFLY OR BEETLE)
S6	14	S5 AND (GENE? OR CLONE? OR DNA?)
S7	11	RD (unique items)
S8	4	DROSPHILA AND (GAL OR GALACTOSIDASE)
S9	502	DROSOPHILA AND (GAL OR GALACTOSIDASE)
S10	210	S9 AND EMBRYO?
S11	0	S10A ND DEVELOP?
S12	125	S10 AND DEVELOP?

?s drosophila(10n) (gal or galactosidase)

35618 DROSOPHILA

7514 GAL

16555 GALACTOSIDASE

S13 181 DROSOPHILA(10N) (GAL OR GALACTOSIDASE)

?s s13 and develop? and embryo?

181 S13

993699 DEVELOP?

163692 EMBRYO?

S14 40 S13 AND DEVELOP? AND EMBRYO?

?rd

...completed examining records

S15 36 RD (unique items)

?t s15/5/1-36

15/5/1 (Item 1 from file: 154)

DIALOG(R)File 154:MEDLINE(R)

(c) format only 1996 Knight-Ridder Info. All rts. reserv.

09502347 96023947

Identification of fat-cell enhancer activity in *Drosophila melanogaster* using P-element enhancer traps.

Hoshizaki DK; Lunz R; Ghosh M; Johnson W

University of Illinois College of Medicine at Chicago, Department of Biochemistry 60612, USA.

Genome (CANADA) Jun 1995, 38 (3) p497-506, ISSN 0831-2796

Journal Code: FNP

Languages: ENGLISH

Document type: JOURNAL ARTICLE

JOURNAL ANNOUNCEMENT: 9601

Subfile: INDEX MEDICUS

To identify genes important in fat-cell metabolism and development, we have screened *Drosophila* stocks carrying an engineered transposable element that can reveal the presence of nearby enhancer elements. We have identified those "enhancer-trap lines" that contain transposable P elements integrated near fat-cell specific enhancer elements. We anticipate that the genes associated with these enhancers will provide information concerning fat-cell function and serve as target genes for studying fat-cell specific gene expression. Furthermore, the identification of enhancer-trap lines active in the developing fat cell should provide an entry point into the molecular and genetic analysis of early fat-cell development. Analysis of two lines has revealed that the transcription factors svp, a steroid-hormone receptor, and Kr, a zinc-finger protein, are present in the fat body; these factors are likely to be involved in fat-cell gene expression. In two other lines, beta-galactosidase was detected in a subset of adipothelial cells that may be the precursors to the adult fat cell. And finally, in a single line transgene activity is present in the progenitor cells of the embryonic fat body. The genes associated with these enhancer-trap lines may be involved in fat-cell development.

Tags: Animal; Support, Non-U.S. Gov't

Descriptors: **Drosophila melanogaster*--Physiology--PH; *Enhancer Elements (Genetics)--Genetics--GE; *Fat Body--Physiology--PH; beta-Galactosidase --Metabolism--ME; *Drosophila melanogaster*--Embryology--EM; *Drosophila melanogaster*--Genetics--GE; DNA-Binding Proteins; Fat Body--Cytology--CY; Fat Body--Embryology--EM; Genes, Insect; Larva--Growth and Development--GD; Larva--Genetics--GE; Mesoderm--Cytology--CY; Mesoderm--Physiology--PH; Receptors, Steroid; Transcription Factors--Genetics--GE; Zinc Fingers

CAS Registry No.: 0 (seven-up protein); 0 (DNA-Binding Proteins); 0 (Receptors, Steroid); 0 (Transcription Factors)

Enzyme No.: EC 3.2.1.23 (beta-Galactosidase)

Gene Symbol: Kr; svp

15/5/2 (Item 2 from file: 154)
DIALOG(R) File 154:MEDLINE(R)
(c) format only 1996 Knight-Ridder Info. All rts. reserv.

09348962 95278962

An improved method for chemical devitellinization of X-gal stained Drosophila embryos.

Singh A; Kango M; Sinha P

Drosophila Stock Center, School of Life Science, Indore, India.

Indian J Exp Biol (INDIA) Feb 1995, 33 (2) p150-2, ISSN 0019-5189

Journal Code: GIZ

Languages: ENGLISH

Document type: JOURNAL ARTICLE

JOURNAL ANNOUNCEMENT: 9509

Subfile: INDEX MEDICUS

In Drosophila developmental biological studies, X-gal staining is commonly employed to study the spatio-temporal expression of the lacZ reporter gene in the transformed flies or their embryos. Study of the lacZ pattern in embryos often suffers from the lack of an efficient and high yielding technique for devitellinization of X-gal stained embryos. Devitellinization techniques employed during antibody staining, in situ hybridization or embryonic cuticular preparations generally do not give satisfactory results when used for similar purpose in X-gal stained embryos. This results in the flaky appearance of the blue stain. We present here an improved chemical devitellinization technique which gives a high yield of devitellinized embryos and a better resolution of the X-gal staining pattern.

Tags: Animal; Comparative Study; Support, Non-U.S. Gov't

Descriptors: *Drosophila--Drug Effects--DE; *Galactosides; *Indoles; *Vitelline Membrane--Drug Effects--DE; Drosophila--Embryology--EM; Embryo, Non-Mammalian--Drug Effects--DE; Stains and Staining

CAS Registry No.: 0 (Galactosides); 0 (Indoles); 7240-90-6 (5-bromo-4-chloro-3-indolyl beta-galactoside)

15/5/3 (Item 3 from file: 154)
DIALOG(R) File 154:MEDLINE(R)
(c) format only 1996 Knight-Ridder Info. All rts. reserv.

09337699 95267699

ovo, a Drosophila gene required for ovarian development, is specifically expressed in the germline and shares most of its coding sequences with shavenbaby, a gene involved in embryo patterning.

Mevel-Ninio M; Terracol R; Salles C; Vincent A; Payre F

Centre de Genetique Moleculaire du C.N.R.S, Gif sur Yvette, France.

Mech Dev (IRELAND) Jan 1995, 49 (1-2) p83-95, ISSN 0925-4773

Journal Code: AXF

Languages: ENGLISH

Document type: JOURNAL ARTICLE

JOURNAL ANNOUNCEMENT: 9508

Subfile: INDEX MEDICUS

Genetic analyses of Drosophila oogenesis have revealed the central role of ovo, a gene required for differentiation of the female germline. A number of recessive ovo mutations also affect the shavenbaby (svb) function required for late embryo patterning, suggesting a tight structural link between ovo and svb. By using various genomic probes for in situ hybridization to wild type and mutant embryos, we show that ovo indeed shares most of its coding sequences with svb. svb expression is detected

early in the presumptive head region and later in each segment. It requires control elements located upstream of the ovo genomic region. ovo expresses abundant maternal RNAs which are uniformly distributed in early cleavage embryos. A fraction that lacks an alternative ovo-specific protein coding region (ORF 2b) is detected in pole cells. Expression of an ovo-specific lacZ reporter gene (ovoB) shows that ovo encodes a nuclear protein present in the germline of both sexes. Zygotic ovoB expression is first detected in embryos at around stage 17 and persists up to the adult stage. Our data show that the germline specific expression of ovo in females correlates with its function in oogenesis. This expression, however, is also observed in males in which ovo is not required.

Tags: Animal; Comparative Study; Female; Male; Support, Non-U.S. Gov't

Descriptors: *Drosophila--Embryology--EM; *Gene Expression Regulation, Developmental--Physiology--PH; *Ovary--Embryology--EM; beta-Galactosidase--Genetics--GE; Amino Acid Sequence; Base Sequence; Drosophila--Genetics--GE; DNA, Recombinant; Embryo, Non-Mammalian; Exons; Genetic Code; Molecular Sequence Data; Ovum--Metabolism--ME; Peptide Chain Initiation; Sequence Homology, Nucleic Acid; Transcription, Genetic

CAS Registry No.: 0 (DNA, Recombinant)

Enzyme No.: EC 3.2.1.23 (beta-Galactosidase)

Gene Symbol: ovo; svb; ovoB; lacZ

15/5/4 (Item 4 from file: 154)

DIALOG(R) File 154:MEDLINE(R)

(c) format only 1996 Knight-Ridder Info. All rts. reserv.

09241451 95171451

Protein kinase A and hedgehog signaling in Drosophila limb development.

Jiang J; Struhl G

Howard Hughes Medical Institute, Department of Genetics and Development Columbia University College of Physicians and Surgeons, New York, New York 10032.

Cell (UNITED STATES) Feb 24 1995, 80 (4) p563-72, ISSN 0092-8674

Journal Code: CQ4

Languages: ENGLISH

Document type: JOURNAL ARTICLE

JOURNAL ANNOUNCEMENT: 9506

Subfile: INDEX MEDICUS

The Drosophila hedgehog (hh) gene encodes a secreted protein involved in organizing growth and patterning in many developmental processes. Hh appears to act by inducing the localized expression of at least two other signaling molecules, decapentaplegic (dpp) and wingless (wg), which then govern cell proliferation and patterning in surrounding tissue. Here, we demonstrate that cyclic AMP (cAMP)-dependent protein kinase A (PKA) is essential during limb development to prevent inappropriate dpp and wg expression. We also show that a constitutively active form of PKA can prevent inappropriate dpp and wg expression, but does not interfere with their normal induction by hh. We propose that the basal activity of PKA imposes a block on the transcription of dpp and wg and that hh exerts its organizing influence by alleviating this block.

Tags: Animal; Support, Non-U.S. Gov't

Descriptors: *Cyclic AMP-Dependent Protein Kinases--Metabolism--ME; *Drosophila--Embryology--EM; *Gene Expression Regulation; *Proteins--Metabolism--ME; *Signal Transduction; beta-Galactosidase--Analysis--AN; Animals, Transgenic; Biological Markers; Cell Communication; Cyclic AMP-Dependent Protein Kinases--Genetics--GE; Down-Regulation (Physiology); Embryonic Induction; Extremities--Embryology--EM; Heat; Insect Hormones

--Biosynthesis--BI; Insect Hormones--Genetics--GE; Membrane Proteins
--Genetics--GE; Proteins--Genetics--GE; Proto-Oncogene Proteins
--Biosynthesis--BI; Stress; Transcription, Genetic; Wing--Embryology--EM
CAS Registry No.: 0 (dpp protein, Drosophila); 0 (patched protein); 0
(Biological Markers); 0 (Insect Hormones); 0 (Membrane Proteins); 0
(Proteins); 0 (Proto-Oncogene Proteins); 117758-26-6 (wingless protein,
Drosophila); 149291-21-4 (hedgehog protein, Drosophila)
Enzyme No.: EC 2.7.10.- (Cyclic AMP-Dependent Protein Kinases); EC
3.2.1.23 (beta-Galactosidase)

15/5/5 (Item 5 from file: 154)
DIALOG(R)File 154:MEDLINE(R)
(c) format only 1996 Knight-Ridder Info. All rts. reserv.

09235092 95165092

Expression of the ligand-binding nicotinic acetylcholine receptor subunit
D alpha 2 in the Drosophila central nervous system.

Jonas PE; Phannavong B; Schuster R; Schroder C; Gundelfinger ED

Center for Molecular Neurobiology, University of Hamburg, Germany.

J Neurobiol (UNITED STATES) Dec 1994, 25 (12) p1494-508, ISSN
0022-3034 Journal Code: JAM

Languages: ENGLISH

Document type: JOURNAL ARTICLE

JOURNAL ANNOUNCEMENT: 9505

Subfile: INDEX MEDICUS

The D alpha 2 gene encodes a ligand-binding subunit of nicotinic
acetylcholine receptors (nAChRs) from Drosophila melanogaster. We have
studied the distribution of D alpha 2 transcripts and protein by in situ
hybridization and immunohistochemistry, respectively, as well as the
regulation of D alpha 2 gene expression in vivo using D alpha 2 promoter
fragments fused to the Escherichia coli lacZ gene. Transcripts and protein
from the D alpha 2 gene were detected exclusively in the central nervous
system. Both in late embryos and adults D alpha 2-like immunoreactivity is
widely but not uniformly distributed in the synaptic neuropil, suggesting
that the D alpha 2 protein is a subunit of a synaptic nicotinic receptor.
Its distribution resembles that of ALS and ARD proteins, two other nAChR
subunits of the fly. Five different D alpha 2-lacZ fusion gene constructs
were introduced into the Drosophila genome by P-element-mediated gene
transfer to identify functional elements of the D alpha 2 promoter. All
constructs produce a basic lacZ expression pattern that is compatible with
the distribution of D alpha 2 transcripts and protein. A 880 bp upstream
fragment harbors the cis elements for the expression of a weak but specific
basic D alpha 2 pattern. The next 350 bp further upstream significantly
enhance beta-galactosidase expression without influencing the pattern of
expression. Between 1.7 and 7.3 kb upstream of the transcription start site
one or more elements that are required for D alpha 2 expression in optic
lobe tangential cells are located.

Tags: Animal; Support, Non-U.S. Gov't

Descriptors: *beta-Galactosidase--Analysis--AN; *Brain Chemistry;
*Drosophila--Genetics--GE; *Gene Expression Regulation, Developmental;
*Promoter Regions (Genetics)--Genetics--GE; *Receptors, Nicotinic
--Ultrastructure--UL; Base Sequence; Central Nervous System--Cytology--CY;
Drosophila--Chemistry--CH; Drosophila--Embryology--EM; In Situ
Hybridization; Introns; Molecular Sequence Data; Receptors, Nicotinic
--Analysis--AN; Receptors, Nicotinic--Chemistry--CH; Receptors, Nicotinic
--Genetics--GE

CAS Registry No.: 0 (Receptors, Nicotinic)

Enzyme No.: EC 3.2.1.23 (beta-Galactosidase)

15/5/6 (Item 6 from file: 154)

DIALOG(R)File 154:MEDLINE(R)

(c) format only 1996 Knight-Ridder Info. All rts. reserv.

09233585 95163585

Ultrabithorax protein is necessary but not sufficient for full activation of decapentaplegic expression in the visceral mesoderm.

Sun B; Hursh DA; Jackson D; Beachy PA

Howard Hughes Medical Institute, Department of Molecular Biology and Genetics, Johns Hopkins University School of Medicine, Baltimore, MD 21205.

EMBO J (ENGLAND) Feb 1 1995, 14 (3) p520-35, ISSN 0261-4189

Journal Code: EMB

Languages: ENGLISH

Document type: JOURNAL ARTICLE

JOURNAL ANNOUNCEMENT: 9505

Subfile: INDEX MEDICUS

To elucidate the mechanisms by which homeotic selector (HOM) genes specify the unique features of Drosophila segments, we have analyzed the regulation of decapentaplegic (dpp), a transforming growth factor (TGF)-beta superfamily member, and have found that the Ultrabithorax (Ubx) HOM protein directly activates dpp expression in parasegment 7 (PS7) of the embryonic visceral mesoderm. Other factors are also required, including one that appears to act through homeodomain protein binding sites and may be encoded by extradenticle (exd). The exd protein binds in a highly co-operative manner to regulatory sequences mediating PS7-specific dpp expression, consistent with a genetic requirement for exd function in normal visceral mesoderm expression of dpp. A second mechanism contributing to PS7 expression of dpp appears not to require Ubx protein directly, and involves a general visceral mesoderm enhancer coupled to a spatially specific repression element. Thus, even in an apparently simple case where visceral mesoderm expression of the dpp target gene mirrors that of the Ubx HOM protein, full activation by Ubx protein requires at least one additional factor. In addition, a distinct regulatory mode not directly involving Ubx protein also appears to contribute to PS7-specific expression.

Tags: Animal

Descriptors: *Drosophila--Embryology--EM; *DNA-Binding Proteins
--Metabolism--ME; *Gene Expression Regulation, Developmental; *Homeodomain
Proteins--Metabolism--ME; *Insect Hormones--Biosynthesis--BI; beta-Galactos
idase--Genetics--GE; Base Sequence; Binding Sites; Carrier Proteins; DNA
--Metabolism--ME; Genes, Insect; Genes, Reporter; Insect Hormones--Genetics
--GE; Mesoderm--Metabolism--ME; Models, Genetic; Molecular Sequence Data;
Regulatory Sequences, Nucleic Acid--Genetics--GE; Sequence Analysis, DNA;
Transcription Factors--Metabolism--ME; Transforming Growth Factor beta
--Biosynthesis--BI; Viscera--Embryology--EM; Viscera--Metabolism--ME

CAS Registry No.: 0 (dpp protein, Drosophila); 0 (extradenticle
protein); 0 (ultrabithorax protein); 0 (Carrier Proteins); 0
(DNA-Binding Proteins); 0 (Homeodomain Proteins); 0 (Insect Hormones);
0 (Transcription Factors); 0 (Transforming Growth Factor beta);
9007-49-2 (DNA)

Enzyme No.: EC 3.2.1.23 (beta-Galactosidase)

Gene Symbol: ubx; dpp

15/5/7 (Item 7 from file: 154)

DIALOG(R)File 154:MEDLINE(R)

(c) format only 1996 Knight-Ridder Info. All rts. reserv.

08971560 94286560

Tau-beta-galactosidase, an axon-targeted fusion protein.

Callahan CA; Thomas JB

Molecular Neurobiology Laboratory, Salk Institute for Biological Studies,
San Diego, CA 92186.

Proc Natl Acad Sci U S A (UNITED STATES) Jun 21 1994, 91 (13) p5972-6,
ISSN 0027-8424 Journal Code: PV3

Contract/Grant No.: GM07198, GM, NIGMS

Languages: ENGLISH

Document type: JOURNAL ARTICLE

JOURNAL ANNOUNCEMENT: 9409

Subfile: INDEX MEDICUS

The most commonly used enzymatic reporter molecule, Escherichia coli beta-galactosidase (beta-gal; beta-D-galactoside galactohydrolase, EC 3.2.1.23), fails to readily diffuse into axons; consequently, the morphologies of beta-gal-labeled neurons cannot directly be determined. For analysis of neuronal pathfinding and synaptic connectivity, this information is essential. We have constructed an axon-targeted beta-gal reporter by fusing the cDNA encoding the bovine microtubule-binding protein, tau, to lacZ, the E. coli gene encoding beta-gal. This reporter labels cell bodies and axons when expressed by developing and adult Drosophila neurons. It also reveals the entire cellular extent of nonneuronal cells such as muscle fibers and glia. To generate neuronal markers for studies of Drosophila neural development, we constructed a tau-beta-gal enhancer-trap transposon. From 1500 independent lines generated by mobilization of this transposon, we have isolated a set of useful markers for specific subsets of neurons, glia, and muscles. Since the tau cDNA-lacZ reporter utilizes bovine tau, it may also effectively target beta-gal in vertebrate neurons and prove to be a useful reagent for the analysis of vertebrate nervous systems.

Tags: Animal; Support, Non-U.S. Gov't; Support, U.S. Gov't, P.H.S.

Descriptors: *beta-Galactosidase--Biosynthesis--BI; *tau Proteins
--Biosynthesis--BI; *Axons--Physiology--PH; *Neurons--Metabolism--ME;
*Recombinant Fusion Proteins--Biosynthesis--BI; beta-Galactosidase
--Analysis--AN; tau Proteins--Analysis--AN; Cattle; Drosophila--Embryology
--EM; Drosophila--Physiology--PH; Embryo, Non-Mammalian--Physiology--PH;
Escherichia coli--Enzymology--EN; Immunohistochemistry; Neurons--Physiology
--PH; Organ Specificity; Recombinant Fusion Proteins--Analysis--AN;
Restriction Mapping

CAS Registry No.: 0 (tau Proteins); 0 (Recombinant Fusion Proteins)

Enzyme No.: EC 3.2.1.23 (beta-Galactosidase)

15/5/8 (Item 8 from file: 154)

DIALOG(R)File 154:MEDLINE(R)

(c) format only 1996 Knight-Ridder Info. All rts. reserv.

08872694 94187694

Multiple cis-acting targeting sequences are required for orb mRNA
localization during Drosophila oogenesis.

Lantz V; Schedl P

Department of Biology, Washington University, St. Louis, Missouri 63130.

Mol Cell Biol (UNITED STATES) Apr 1994, 14 (4) p2235-42, ISSN
0270-7306 Journal Code: NGY

Languages: ENGLISH

Document type: JOURNAL ARTICLE

JOURNAL ANNOUNCEMENT: 9406

Subfile: INDEX MEDICUS

The targeting of positional information to specific regions of the oocyte or early embryo is one of the key processes in establishing anterior-posterior and dorsal-ventral polarity. In many developmental systems, this is accomplished by localization of mRNAs. The germ line-specific *Drosophila orb* gene plays a critical role in defining both axes of the developing oocyte, and its mRNA is localized in a complex pattern during oogenesis. We have identified a 280-bp sequence from the orb 3' untranslated region capable of reproducing this complex localization pattern. Furthermore, we have found that multiple cis-acting elements appear to be required for proper targeting of orb mRNA.

Tags: Animal; Female; Support, U.S. Gov't, P.H.S.

Descriptors: **Drosophila melanogaster*--Physiology--PH; *Gene Expression; *Oocytes--Physiology--PH; *Oogenesis; *RNA, Messenger--Biosynthesis--BI; beta-Galactosidase--Biosynthesis--BI; *Drosophila melanogaster*--Genetics--GE; Embryo, Non-Mammalian--Physiology--PH; Heat-Shock Proteins--Biosynthesis--BI; Ovary--Physiology--PH; Plasmids; Promoter Regions (Genetics); Recombinant Proteins--Biosynthesis--BI; Restriction Mapping; RNA, Messenger--Analysis--AN; Translation, Genetic

CAS Registry No.: 0 (Heat-Shock Proteins); 0 (Plasmids); 0 (Recombinant Proteins); 0 (RNA, Messenger)

Enzyme No.: EC 3.2.1.23 (beta-Galactosidase)

Gene Symbol: orb

15/5/9 (Item 9 from file: 154)

DIALOG(R) File 154:MEDLINE(R)

(c) format only 1996 Knight-Ridder Info. All rts. reserv.

08717745 94032745

A double staining technique using 5-bromo,4-chloro-3-indolyl-beta-D-galactopyranoside (X-gal) and immunoperoxidase in whole *Drosophila* embryos.

Kobayashi S; Okada M

Institute of Biological Sciences, University of Tsukuba, Ibaraki, Japan.

Biotech Histochem (UNITED STATES) Jul 1993, 68 (4) p237-9, ISSN 1052-0295 Journal Code: A29

Languages: ENGLISH

Document type: JOURNAL ARTICLE

JOURNAL ANNOUNCEMENT: 9402

Subfile: INDEX MEDICUS

We have developed a double staining method using 5-bromo,4-chloro-3-indolyl-beta-D-galactopyranoside X-gal and immunoperoxidase for whole *Drosophila* embryos. The dechorionated embryos are fixed in heptane saturated with 4% formaldehyde, then in heptane and 50% methanol. Fixed embryos are devitellinized with a tungsten needle and processed for immunoperoxidase staining immediately prior to peroxidase color development. The embryos are stained with X-gal, then peroxidase staining is resumed. This procedure enables us to observe cells stained with both X-gal and a specific antibody in whole embryos.

Tags: Animal; Support, Non-U.S. Gov't

Descriptors: *Embryo, Non-Mammalian--Ultrastructure--UL; *Drosophila*; Galactosidases--Immunology--IM; Galactosides; Immunoenzyme Techniques; Indoles; Paraffin

CAS Registry No.: 0 (Galactosides); 0 (Indoles); 7240-90-6 (5-bromo-4-chloro-3-indolyl beta-galactoside); 8002-74-2 (Paraffin)

Enzyme No.: EC 3.2.1.- (Galactosidases)

15/5/10 (Item 10 from file: 154)
DIALOG(R) File 154:MEDLINE(R)
(c) format only 1996 Knight-Ridder Info. All rts. reserv.

08434171 93144171

The mouse Enhancer trap locus 1 (Etl-1): a novel mammalian gene related to Drosophila and yeast transcriptional regulator genes.

Soininen R; Schoor M; Henseling U; Tepe C; Kisters-Woike B; Rossant J; Gossler A

Max-Delbrück-Laboratorium, Max-Planck-Gesellschaft, Köln, FRG.

Mech Dev (IRELAND) Nov 1992, 39 (1-2) p111-23, ISSN 0925-4773

Journal Code: AXF

Contract/Grant No.: 5R01HD2533403, HD, NICHD

Languages: ENGLISH

Document type: JOURNAL ARTICLE

JOURNAL ANNOUNCEMENT: 9305

Subfile: INDEX MEDICUS

A novel mouse gene, Enhancer trap locus 1 (Etl-1), was identified in close proximity to a lacZ enhancer trap integration in the mouse genome showing a specific beta-galactosidase staining pattern during development. In situ analysis revealed a widespread but not ubiquitous expression of Etl-1 throughout development with particularly high levels in the central nervous system and epithelial cells. The amino acid sequence of the Etl-1 protein deduced from the cDNA shows strong similarity, over a stretch of 500 amino acids, to the Drosophila brahma protein involved in the regulation of homeotic genes and to the yeast transcriptional activator protein SNF2/SWI2 as well as to the RAD54 protein and the recently described helicase-related yeast proteins STH1 and MOT1. Etl-1 is the first mammalian member of this group of proteins that are implicated in gene regulation and/or influencing chromatin structure. The homology to the regulatory proteins SNF2/SWI2 and brahma and the expression pattern during embryogenesis suggest that Etl-1 protein might be involved in gene regulating pathways during mouse development.

Tags: Animal; Comparative Study; Support, Non-U.S. Gov't; Support, U.S. Gov't, P.H.S.

Descriptors: *Enhancer Elements (Genetics); *Genes, Structural; *Mice --Genetics--GE; *Proteins--Genetics--GE; beta-Galactosidase--Biosynthesis --BI; beta-Galactosidase--Genetics--GE; Amino Acid Sequence; Base Sequence ; Drosophila melanogaster--Genetics--GE; DNA--Genetics--GE; Fetal Development--Genetics--GE; Genetic Techniques; Mice--Embryology--EM; Molecular Sequence Data; Open Reading Frames; Organ Specificity; Proteins --Biosynthesis--BI; Recombinant Fusion Proteins--Biosynthesis--BI; Recombinant Fusion Proteins--Genetics--GE; RNA, Messenger--Analysis--AN; Saccharomyces cerevisiae--Genetics--GE; Sequence Alignment; Sequence Homology, Amino Acid; Transcription Factors--Biosynthesis--BI; Transcription Factors--Genetics--GE

CAS Registry No.: 0 (Etl-1 protein); 0 (Proteins); 0 (Recombinant Fusion Proteins); 0 (RNA, Messenger); 0 (Transcription Factors); 9007-49-2 (DNA)

Enzyme No.: EC 3.2.1.23 (beta-Galactosidase)

Gene Symbol: ETL-1; LACZ; SNF2; SW12; MOT1; STH1; brm; neo

15/5/11 (Item 11 from file: 154)
DIALOG(R) File 154:MEDLINE(R)
(c) format only 1996 Knight-Ridder Info. All rts. reserv.

08432018 93142018

A wingless-dependent polar coordinate system in *Drosophila* imaginal discs
[see comments]

Couso JP; Bate M; Martinez-Arias A

Department of Zoology, University of Cambridge, United Kingdom.

Science (UNITED STATES) Jan 22 1993, 259 (5094) p484-9, ISSN

0036-8075 Journal Code: UJ7

Comment in Science 1993 Jan 22;259(5094):471-2

Languages: ENGLISH

Document type: JOURNAL ARTICLE

JOURNAL ANNOUNCEMENT: 9304

Subfile: INDEX MEDICUS

The patterning of the imaginal discs in *Drosophila melanogaster* is a progressive process that, like the patterning of the larval epidermis during embryogenesis, requires the activity of segment polarity genes. One segment polarity gene, wingless, encodes a homolog of the mouse proto-oncogene Wnt-1 and plays a prominent role in the patterning of the larval epidermis and the imaginal discs. However, whereas the function of wingless in the embryo is initially associated with a pattern of stripes along the anteroposterior axis that are part of a Cartesian coordinate system, it is shown here that during imaginal development wingless is associated with a pattern of sectors that provide references for a polar coordinate system homologous to that postulated in a well-known model for the regeneration of insect and vertebrate limbs.

Tags: Animal; Comparative Study; Support, Non-U.S. Gov't

Descriptors: **Drosophila melanogaster*--Genetics--GE; beta-Galactosidase--Genetics--GE; *Drosophila melanogaster*--Embryology--EM; *Drosophila melanogaster*--Growth and Development--GD; Embryo, Non-Mammalian--Cytology--CY; Embryo, Non-Mammalian--Physiology--PH; Gene Expression; Larva; Mice; Phenotype; Protein-Tyrosine Kinase--Genetics--GE; Proto-Oncogene Proteins--Genetics--GE; Proto-Oncogenes; Sequence Homology, Nucleic Acid; Wing

CAS Registry No.: 0 (proto-oncogene protein int-1); 0 (Proto-Oncogene Proteins)

Enzyme No.: EC 2.7.1.112 (Protein-Tyrosine Kinase); EC 3.2.1.23 (beta-Galactosidase)

Gene Symbol: lacZ; wg; en; Wnt-1

15/5/12 (Item 12 from file: 154)

DIALOG(R) File 154:MEDLINE(R)

(c) format only 1996 Knight-Ridder Info. All rts. reserv.

08391146 93101146

Apical secretion and association of the *Drosophila* yellow gene product with developing larval cuticle structures during embryogenesis.

Kornezos A; Chia W

Drosophila Neurobiology Laboratory, National University of Singapore, Kent Ridge Crescent.

Mol Gen Genet (GERMANY) Nov 1992, 235 (2-3) p397-405, ISSN 0026-8925

Journal Code: NGP

Languages: ENGLISH

Document type: JOURNAL ARTICLE

JOURNAL ANNOUNCEMENT: 9303

Subfile: INDEX MEDICUS

The yellow (y) gene of *Drosophila melanogaster* is required for the pigmentation of larval and adult cuticle structures. The deduced y protein sequence includes two putative N-linked glycosylation sites and a putative

signal peptide, suggesting that it might be a secreted molecule. Consistent with the characteristics of a secreted protein, our in vitro translation studies using RNA synthesised from the y cDNA demonstrate that the nascent y polypeptide is a preprotein that cotranslationally translocates into the endoplasmic reticulum (ER) membrane and becomes glycosylated. The N-terminal peptide is cleaved from the preprotein between the two alanine residues at positions 21 and 22, to release the final product into the lumen of the ER. Antibodies raised against the y polypeptide detect the protein starting at 13 h post-fertilization in epidermal cells and in the cuticle structures secreted by them that later become pigmented; in addition, yellow protein is detected in the cuticle structures associated with Keilin's organs. The embryonic beta-galactosidase staining pattern of a transgene, bearing a construct in which expression of the lacZ gene is driven by the y promoter, is also described and is similar to that of the y protein. Our results indicate that the y gene product is an apically secreted protein which becomes an immobilised structural component of the pigmented cuticle.

Tags: Animal; Support, Non-U.S. Gov't

Descriptors: *Drosophila melanogaster--Genetics--GE; *Embryo, Non-Mammalian--Physiology--PH; *Insect Hormones--Genetics--GE; beta-Galactosidase--Genetics--GE; beta-Galactosidase--Metabolism--ME; Amino Acid Sequence; Base Sequence; Drosophila melanogaster--Embryology--EM; DNA--Genetics--GE; Endoplasmic Reticulum--Metabolism--ME; Fertilization; Insect Hormones--Analysis--AN; Insect Hormones--Biosynthesis--BI; Larva; Molecular Sequence Data; Oligodeoxyribonucleotides; Polymerase Chain Reaction--Methods--MT; Recombinant Fusion Proteins--Metabolism--ME; Restriction Mapping; Translation, Genetic

CAS Registry No.: 0 (yellow locus protein, Drosophila); 0 (Insect Hormones); 0 (Oligodeoxyribonucleotides); 0 (Recombinant Fusion Proteins); 9007-49-2 (DNA)

Enzyme No.: EC 3.2.1.23 (beta-Galactosidase)

15/5/13 (Item 13 from file: 154)

DIALOG(R) File 154:MEDLINE(R)

(c) format only 1996 Knight-Ridder Info. All rts. reserv.

08219775 92357775

dOct2, a Drosophila Oct transcription factor that functions in yeast.

Prakash K; Fang XD; Engelberg D; Behal A; Parker CS

Division of Chemistry, California Institute of Technology, Pasadena 91125.

Proc Natl Acad Sci U S A (UNITED STATES) Aug 1 1992, 89 (15) p7080-4, ISSN 0027-8424 Journal Code: PV3

Contract/Grant No.: GM42671, GM, NIGMS

Languages: ENGLISH

Document type: JOURNAL ARTICLE

JOURNAL ANNOUNCEMENT: 9211

Subfile: INDEX MEDICUS

Oct factors are members of the POU family of transcription factors that are shown to play important roles during development in mammals. Here we report the cDNA cloning and expression of a Drosophila Oct transcription factor. Whole mount in situ hybridization experiments revealed that the spatial expression patterns of this gene during embryonic development have not yet been observed for any other gene. In early embryogenesis, its transcripts are transiently expressed as a wide uniform band from 20% to 40% of the egg length, very similar to that of gap genes. This pattern progressively resolves into a series of narrower stripes followed by

expression in 14 stripes. Subsequently, transcripts from this gene are expressed in the central nervous system and the brain. When expressed in the yeast *Saccharomyces cerevisiae*, this *Drosophila* factor functions as a strong, octamer-dependent activator of transcription. Our data strongly suggest possible functions for the Oct factor in pattern formation in *Drosophila* that might transcend the boundaries of genetically defined segmentation genes.

Tags: Animal; Comparative Study; Human; Support, Non-U.S. Gov't; Support, U.S. Gov't, P.H.S.

Descriptors: **Drosophila*--Genetics--GE; *DNA-Binding Proteins--Metabolism--ME; **Saccharomyces cerevisiae*--Genetics--GE; *Transcription Factors--Metabolism--ME; beta-Galactosidase--Genetics--GE; beta-Galactosidase--Isolation and Purification--IP; beta-Galactosidase--Metabolism--ME; Amino Acid Sequence; Base Sequence; Cloning, Molecular; *Drosophila*--Metabolism--ME; DNA--Genetics--GE; Embryo, Non-Mammalian; Gene Library; Molecular Sequence Data; Plasmids; Protein Conformation; Recombinant Fusion Proteins--Isolation and Purification--IP; Recombinant Fusion Proteins--Metabolism--ME; Sequence Homology, Nucleic Acid; Transcription Factors--Genetics--GE; Transcription Factors--Isolation and Purification--IP

Molecular Sequence Databank No.: GENBANK/M93149

CAS Registry No.: 0 (transcription factor OTF-2); 0 (DNA-Binding Proteins); 0 (Plasmids); 0 (Recombinant Fusion Proteins); 0 (Transcription Factors); 9007-49-2 (DNA)

Enzyme No.: EC 3.2.1.23 (beta-Galactosidase)

15/5/14 (Item 14 from file: 154)

DIALOG(R)File 154:MEDLINE(R)

(c) format only 1996 Knight-Ridder Info. All rts. reserv.

08209352 92347352

Individual dorsal morphogen binding sites mediate activation and repression in the *Drosophila* embryo.

Jiang J; Rushlow CA; Zhou Q; Small S; Levine M

Department of Biological Sciences, Fairchild Center, Columbia University, New York, NY 10027.

EMBO J (ENGLAND) Aug 1992, 11 (8) p3147-54, ISSN 0261-4189

Journal Code: EMB

Contract/Grant No.: GM 46638, GM, NIGMS

Languages: ENGLISH

Document type: JOURNAL ARTICLE

JOURNAL ANNOUNCEMENT: 9211

Subfile: INDEX MEDICUS

The dorsal (dl) morphogen gradient is responsible for initiating the differentiation of the mesoderm, neuroectoderm and dorsal ectoderm in the *Drosophila* embryo. dl encodes a sequence-specific DNA binding protein that belongs to the Rel family of transcription factors. Previous studies have shown that dl activates the mesoderm determinant twist (twi); here we use a combination of site-directed mutagenesis and P-transformation assays to demonstrate that it also functions as a direct transcriptional repressor of a second target gene, *zerknüllt* (zen). By exchanging dl binding sites between the promoters we show that activator sites from twi can mediate repression when placed in the context of the zen promoter, and that repressor sites from zen can mediate activation in the context of the twi promoter. This represents the first demonstration that common binding sites for any DNA binding protein can mediate both activation and repression in a developing embryo. Evidence is also presented that the affinities of dl binding sites are important for the efficiency of repression, but are not

the sole determinants of the threshold response to the dl gradient.

Tags: Animal; Support, U.S. Gov't, P.H.S.

Descriptors: *Drosophila--Genetics--GE; *DNA-Binding Proteins--Genetics--GE; *Nuclear Proteins--Genetics--GE; beta-Galactosidase--Genetics--GE; beta-Galactosidase--Metabolism--ME; Base Sequence; Binding Sites; Cloning, Molecular; Drosophila--Embryology--EM; DNA Insertion Elements; DNA-Binding Proteins--Metabolism--ME; Embryo, Non-Mammalian--Physiology--PH; Heat-Shock Proteins--Genetics--GE; Heat-Shock Proteins--Metabolism--ME; Molecular Sequence Data; Morphogenesis--Genetics--GE; Mutagenesis, Site-Directed; Nuclear Proteins--Metabolism--ME; Oligodeoxyribonucleotides; Promoter Regions (Genetics); Recombinant Fusion Proteins--Metabolism--ME; Transformation, Genetic

CAS Registry No.: 0 (dorsal protein, Drosophila); 0 (DNA Insertion Elements); 0 (DNA-Binding Proteins); 0 (Heat-Shock Proteins); 0 (Nuclear Proteins); 0 (Oligodeoxyribonucleotides); 0 (Recombinant Fusion Proteins)

Enzyme No.: EC 3.2.1.23 (beta-Galactosidase)

Gene Symbol: dl; zen; lacZ

15/5/15 (Item 15 from file: 154)

DIALOG(R)File 154:MEDLINE(R)

(c) format only 1996 Knight-Ridder Info. All rts. reserv.

08090796 92228796

Dorsal-ventral patterning in Drosophila: DNA binding of snail protein to the single-minded gene.

Kasai Y; Nambu JR; Lieberman PM; Crews ST

Department of Biology, University of California, Los Angeles 90024.

Proc Natl Acad Sci U S A (UNITED STATES) Apr 15 1992, 89 (8) p3414-8,

ISSN 0027-8424 Journal Code: PV3

Contract/Grant No.: T 32 CA09056, CA, NCI; R01 HD25251, HD, NICHD

Languages: ENGLISH

Document type: JOURNAL ARTICLE

JOURNAL ANNOUNCEMENT: 9207

Subfile: INDEX MEDICUS

The Drosophila snail gene is required for proper mesodermal development. Genetic studies suggest that it functions by repressing adjacent ectodermal gene expression including that of the single-minded (sim) gene. The snail gene encodes a protein with a zinc-finger motif, and here we report that the snail gene product is a sequence-specific DNA binding protein. The snail protein recognizes a 14-base-pair consensus sequence that is found nine times in a 2.8-kilobase sim regulatory region. These results provide evidence for the direct control of sim transcription by snail.

Tags: Animal; Support, Non-U.S. Gov't; Support, U.S. Gov't, P.H.S.

Descriptors: *Drosophila--Genetics--GE; *DNA-Binding Proteins--Genetics--GE; *Genes; *Zinc Fingers--Genetics--GE; beta-Galactosidase--Genetics--GE; beta-Galactosidase--Metabolism--ME; Alleles; Base Sequence; Blastoderm--Physiology--PH; Drosophila--Embryology--EM; DNA-Binding Proteins--Metabolism--ME; Embryo, Non-Mammalian--Physiology--PH; Molecular Sequence Data; Oligodeoxyribonucleotides; Oncogene Proteins, Viral--Genetics--GE; Promoter Regions (Genetics)

CAS Registry No.: 0 (Adenovirus Early Proteins); 0 (DNA-Binding Proteins); 0 (Oligodeoxyribonucleotides); 0 (Oncogene Proteins, Viral)

Enzyme No.: EC 3.2.1.23 (beta-Galactosidase)

Gene Symbol: sim; Sna

15/5/16 (Item 16 from file: 154)
DIALOG(R) File 154:MEDLINE(R)
(c) format only 1996 Knight-Ridder Info. All rts. reserv.

08073453 92211453

Functional assay of a putative Drosophila sodium channel gene in homozygous deficiency neurons.

Germeraad S; O'Dowd D; Aldrich RW

Department of Biological Sciences, San Jose State University, CA 95192.

J Neurogenet (ENGLAND) Feb 1992, 8 (1) p1-16, ISSN 0167-7063

Journal Code: JKE

Contract/Grant No.: NS27501, NS, NINDS; NS23294, NS, NINDS

Languages: ENGLISH

Document type: JOURNAL ARTICLE

JOURNAL ANNOUNCEMENT: 9207

Subfile: INDEX MEDICUS

Using voltage-clamp techniques, we have examined embryonic sodium currents in neurons deficient for a gene located at 60E5/6 that shares extensive amino acid similarity with vertebrate sodium channel genes. These neurons expressed sodium currents similar to wildtype, supporting the hypothesis that para, and not the gene at 60E5/6, is the primary sodium channel gene expressed in embryonic neurons. A simple marking procedure allowing positive identification of the genotypes of cultured Drosophila embryos obtained from heterozygous parents was used to recognize cultures homozygous for deficiencies. The morphological development of both neurons and myotubes in these cultures was similar to wildtype, making it feasible to compare the properties of normal diploid cells and cells completely lacking a putative sodium channel gene.

Tags: Animal; Support, Non-U.S. Gov't; Support, U.S. Gov't, P.H.S.

Descriptors: *Drosophila--Genetics--GE; *Neurons--Physiology--PH; *Sodium Channels--Physiology--PH; beta-Galactosidase--Analysis--AN; beta-Galactosidase--Genetics--GE; Cells, Cultured; Chromosome Deletion; Chromosome Mapping; Drosophila--Embryology--EM; Drosophila--Physiology--PH; Homozygote; Ion Channel Gating--Genetics--GE; Larva; Mutation--Genetics--GE; Phenotype; Sequence Homology, Nucleic Acid

CAS Registry No.: 0 (Sodium Channels)

Enzyme No.: EC 3.2.1.23 (beta-Galactosidase)

15/5/17 (Item 17 from file: 154)
DIALOG(R) File 154:MEDLINE(R)
(c) format only 1996 Knight-Ridder Info. All rts. reserv.

08026858 92164858

Regulatory interactions and role in cell type specification of the Malpighian tubules by the cut, Kruppel, and caudal genes of Drosophila.

Liu S; Jack J

Program in Molecular Biology, Sloan-Kettering Institute, New York, New York.

✓ Dev Biol (UNITED STATES) Mar 1992, 150 (1) p133-43, ISSN 0012-1606

Journal Code: E7T

Languages: ENGLISH

Document type: JOURNAL ARTICLE

JOURNAL ANNOUNCEMENT: 9206

Subfile: INDEX MEDICUS

Kruppel and caudal genes are both required for normal segmentation of the embryo, and the developmental regulatory gene cut is necessary for the normal specification of external sensory organs. These three genes are also

expressed in the Malpighian tubules before and during differentiation. Two of the genes, Kruppel and cut, are known to be required for development of the tubules. We report that the absence of maternal and zygotic caudal function reduces their normal growth and elongation. Normal Kruppel function, which is known to be required for caudal expression, is also required for cut expression, while cut and caudal are expressed independently of each other. Cell type transformations of Malpighian tubules were studied by examining the effects of mutations on the expression of markers specific to Malpighian tubules, hindgut, or midgut of normal embryos. Loss of Kruppel activity confers hindgut characteristics on those cells that normally form the Malpighian tubules with all markers tested. Loss of cut function alters the expression of some markers but not others. The pathway of tissue specific gene regulation, apparently, branches beyond Kruppel to form at least a cut and a caudal branch.

Tags: Animal; Comparative Study; Support, U.S. Gov't, Non-P.H.S.

Descriptors: *Cell Differentiation--Genetics--GE; *Drosophila--Genetics--GE; *Gene Expression Regulation, Enzymologic--Genetics--GE; *Malpighian Tubules--Embryology--EM; beta-Galactosidase--Analysis--AN; Drosophila--Embryology--EM; DNA--Analysis--AN; Malpighian Tubules--Metabolism--ME; Mutation

CAS Registry No.: 9007-49-2 (DNA)

Enzyme No.: EC 3.2.1.23 (beta-Galactosidase)

Gene Symbol: cut; Kruppel; caudal

15/5/18 (Item 18 from file: 154)

DIALOG(R) File 154:MEDLINE(R)

(c) format only 1996 Knight-Ridder Info. All rts. reserv.

08018168 92156168

A single locus encodes both phenylalanine hydroxylase and tryptophan hydroxylase activities in Drosophila.

Neckameyer WS; White K

Department of Biology, Brandeis University, Waltham, Massachusetts 02254.

J Biol Chem (UNITED STATES) Feb 25 1992, 267 (6) p4199-206, ISSN

0021-9258 Journal Code: HIV

Contract/Grant No.: NS23510, NS, NINDS; RRO 4671, RR, NCRR

Languages: ENGLISH

Document type: JOURNAL ARTICLE

JOURNAL ANNOUNCEMENT: 9205

Subfile: INDEX MEDICUS

We have used a full-length clone encoding rabbit tryptophan hydroxylase (TRH) to isolate the Drosophila homologue (DTPH). Southern analysis of Drosophila genomic DNA reveals a pattern indicative of a single gene. The single transcript is expressed in adult head and body mRNA but is also detected in mRNA from early embryos. The embryonic transcript is ubiquitously expressed and appears to concentrate in yolk granules. In situ hybridization of TRH-homologous antisense RNA probe to sectioned tissue from third instar larvae demonstrated the presence of this transcript in fat body and cuticular tissue. Developmental immunoblot analysis using antibodies raised against a beta-galactosidase-Drosophila fusion protein revealed a 45-kDa embryonic protein also detected in female abdomens and a 50-kDa protein found in larval and adult stages. Immunocytochemical analysis of the Drosophila protein in the larval central nervous system showed that it appeared to be present in both serotonin- and catecholamine-containing neurons. A nonfusion protein generated in Escherichia coli hydroxylates both tryptophan and phenylalanine. We propose that there are only two aromatic amino acid hydroxylase genes in

Drosophila: one encoding tyrosine hydroxylase, DTH, and DTPH, a gene encoding both tryptophan and phenylalanine hydroxylase activities.

Tags: Animal; Female; Male; Support, U.S. Gov't, P.H.S.

Descriptors: *Drosophila melanogaster--Genetics--GE; *Phenylalanine Hydroxylase--Genetics--GE; *Tryptophan Hydroxylase--Genetics--GE; beta-Galactosidase--Metabolism--ME; Amino Acid Sequence; Base Sequence; Blotting, Western; Chromosome Mapping; Drosophila melanogaster--Enzymology--EN; DNA--Genetics--GE; Electrophoresis, Polyacrylamide Gel; Gene Expression; Immunohistochemistry; Molecular Sequence Data; Nucleic Acid Hybridization; Phenylalanine Hydroxylase--Metabolism--ME; Recombinant Fusion Proteins--Metabolism--ME; Restriction Mapping; RNA--Genetics--GE; Tryptophan Hydroxylase--Metabolism--ME

Molecular Sequence Databank No.: GENBANK/M81833; GENBANK/X59129; GENBANK/X59130; GENBANK/L01671; GENBANK/L01672; GENBANK/L01673; GENBANK/L01674; GENBANK/L01675; GENBANK/L01676; GENBANK/L01677

CAS Registry No.: 0 (Recombinant Fusion Proteins); 0 (RNA); 9007-49-2 (DNA)

Enzyme No.: EC 1.14.16.1 (Phenylalanine Hydroxylase); EC 1.14.16.4 (Tryptophan Hydroxylase); EC 3.2.1.23 (beta-Galactosidase)

Gene Symbol: DTH; DTPH

15/5/19 (Item 19 from file: 154)

DIALOG(R) File 154:MEDLINE(R)

(c) format only 1996 Knight-Ridder Info. All rts. reserv.

07828894 91347894

An extensive 3' cis-regulatory region directs the imaginal disk expression of decapentaplegic, a member of the TGF-beta family in Drosophila.

Blackman RK; Sanicola M; Raftery LA; Gillevet T; Gelbart WM

Department of Cellular and Developmental Biology, Harvard University, Cambridge, MA 02138-2097.

Development (ENGLAND) Mar 1991, 111 (3) p657-66, ISSN 0950-1991

Journal Code: ECW

Languages: ENGLISH

Document type: JOURNAL ARTICLE

JOURNAL ANNOUNCEMENT: 9112

Subfile: INDEX MEDICUS

The decapentaplegic (dpp) gene in Drosophila melanogaster encodes a TGF-beta-like signalling molecule that is expressed in a complex and changing pattern during development. One of dpp's contributions is to proximal-distal outgrowth of the adult appendages, structures derived from the larval imaginal disks. Appendage specific mutations of dpp fall in a 20 kb interval 3' to the known dpp transcripts. Here, we directly test the hypothesis that these mutations define an extended 3' cis-regulatory region. By analysis of germ-line transformants expressing a reporter gene, we show that sequences from this portion of the gene, termed the dppdisk region, are capable of directing expression comparable to that defined by RNA in situ hybridization. We localize two intervals of the dppdisk region that appear to account for much of the dpp spatial pattern in imaginal disks and discuss the positions of these important elements in terms of the genetics of dpp. Finally, we provide evidence to suggest that one of our constructs expresses beta-galactosidase in the early imaginal disk primordia in the embryo, at approximately the time when they are set aside from surrounding larval epidermal tissues. Thus, dpp may be involved directly in the determination of the imaginal disks.

Tags: Animal; Female; Male; Support, Non-U.S. Gov't; Support, U.S. Gov't,

P.H.S.

Descriptors: *Drosophila--Genetics--GE; *DNA; *Gene Expression--Genetics--GE; *Genes, Structural--Physiology--PH; *Regulatory Sequences, Nucleic Acid--Physiology--PH; beta-Galactosidase--Genetics--GE; Animals, Transgenic; Drosophila--Embryology--EM; Drosophila--Ultrastructure--UL; Enhancer Elements (Genetics)--Genetics--GE; Microscopy, Electron; Transforming Growth Factor beta--Genetics--GE

CAS Registry No.: 0 (Transforming Growth Factor beta); 9007-49-2 (DNA)
Enzyme No.: EC 3.2.1.23 (beta-Galactosidase)

15/5/20 (Item 20 from file: 154)

DIALOG(R) File 154:MEDLINE(R)

(c) format only 1996 Knight-Ridder Info. All rts. reserv.

07811452 91330452

Generating lineage-specific markers to study Drosophila development.

Perrimon N; Noll E; McCall K; Brand A

Department of Genetics, Harvard Medical School, Boston, MA 02115.

Dev Genet (UNITED STATES) 1991, 12 (3) p238-52, ISSN 0192-253X

Journal Code: DEG

Languages: ENGLISH

Document type: JOURNAL ARTICLE

JOURNAL ANNOUNCEMENT: 9111

Subfile: INDEX MEDICUS

To generate cell- and tissue-specific expression patterns of the reporter gene lacZ in Drosophila, we have generated and characterized 1,426 independent insertion strains using four different P-element constructs. These four transposons carry a lacZ gene driven either by the weak promoter of the P-element transposase gene or by partial promoters from the even-skipped, fushi-tarazu, or engrailed genes. The tissue-specific patterns of beta-galactosidase expression that we are able to generate depend on the promoter utilized. We describe in detail 13 strains that can be used to follow specific cell lineages and demonstrate their utility in analyzing the phenotypes of developmental mutants. Insertion strains generated with P-elements that carry various sequences upstream of the lacZ gene exhibit an increased variety of expression patterns that can be used to study Drosophila development.

Tags: Animal; Female; Male; Support, Non-U.S. Gov't

Descriptors: *Drosophila--Genetics--GE; *Genetic Markers--Genetics--GE; beta-Galactosidase--Genetics--GE; Cloning, Molecular; Drosophila--Embryology--EM; DNA Insertion Elements; Gene Expression; Immunoenzyme Techniques; Mutation; Organ Specificity--Genetics--GE; Promoter Regions (Genetics)

CAS Registry No.: 0 (DNA Insertion Elements); 0 (Genetic Markers)

Enzyme No.: EC 3.2.1.23 (beta-Galactosidase)

15/5/21 (Item 21 from file: 154)

DIALOG(R) File 154:MEDLINE(R)

(c) format only 1996 Knight-Ridder Info. All rts. reserv.

07546324 91065324

Regulatory elements of the bithorax complex that control expression along the anterior-posterior axis.

Simon J; Peifer M; Bender W; O'Connor M

Department of Biological Chemistry and Molecular Pharmacology, Harvard Medical School, Boston, MA 02115.

EMBO J (ENGLAND) Dec 1990, 9 (12) p3945-56, ISSN 0261-4189
Journal Code: EMB
Languages: ENGLISH
Document type: JOURNAL ARTICLE
JOURNAL ANNOUNCEMENT: 9103
Subfile: INDEX MEDICUS

The *Drosophila bithorax* complex (BX-C) controls segmental development by selectively deploying three protein products, Ubx, abd-A and Abd-B, within specific segments along the body axis. Expression of these products within any one segment (or, more accurately, parasegment) is affected by mutations clustered in a particular region of the BX-C. The regulatory regions defined by this genetic analysis span 20-50 kb and there is one region for each segmental unit. Here we describe regulatory elements from several of these regions, identified by fusion to a Ubx-lacZ gene and analysis in germline transformants. A small DNA fragment from the abx region programs expression with an anterior boundary in the second thoracic segment (parasegment 5). This anterior limit is appropriate, since the abx region normally controls Ubx in parasegment 5. Other regulatory regions of the BX-C that control development of parasegments 6, 7 or 8 contain similar regulatory elements that program expression with anterior limits in parasegments 6, 7 or 8, respectively. These experiments define a class of BX-C regulatory elements that control expression along the anterior-posterior axis. The early appearance of the lacZ patterns in embryos suggests a role for these elements in the initial activation of expression from the BX-C.

Tags: Animal; Female; Male; Support, Non-U.S. Gov't; Support, U.S. Gov't, P.H.S.

Descriptors: *Drosophila--Genetics--GE; *Genes, Regulator; beta-Galactosidase--Genetics--GE; beta-Galactosidase--Metabolism--ME; Crosses, Genetic; Drosophila--Anatomy and Histology--AH; Drosophila--Embryology--EM; Embryo, Non-Mammalian--Physiology--PH; Enhancer Elements (Genetics); Genes, Homeobox; Genetic Vectors; Recombinant Fusion Proteins--Biosynthesis--BI; Thorax--Anatomy and Histology--AH

CAS Registry No.: 0 (Recombinant Fusion Proteins)

Enzyme No.: EC 3.2.1.23 (beta-Galactosidase)

Gene Symbol: Ubx

15/5/22 (Item 22 from file: 154)

DIALOG(R) File 154:MEDLINE(R)

(c) format only 1996 Knight-Ridder Info. All rts. reserv.

07527017 91046017

Evidence for positive and negative regulation of the Hox-3.1 gene.

Bieberich CJ; Utset MF; Awgulewitsch A; Ruddle FH

Department of Biology, Yale University, New Haven, CT 06511.

Proc Natl Acad Sci U S A (UNITED STATES) Nov 1990, 87 (21) p8462-6,

ISSN 0027-8424 Journal Code: PV3

Contract/Grant No.: GM 009966, GM, NIGMS; GM 43334-01, GM, NIGMS

Languages: ENGLISH

Document type: JOURNAL ARTICLE

JOURNAL ANNOUNCEMENT: 9102

Subfile: INDEX MEDICUS

The region-specific patterns of expression of mouse homeobox genes are considered important for establishing the embryonic body plan. A 5-kilobase (kb) DNA fragment from the Hox-3.1 locus that is sufficient to confer region-specific expression to a beta-galactosidase reporter gene in transgenic mouse embryos has been defined. The observed reporter gene

expression pattern closely parallels endogenous Hox-3.1 expression in 8- to 9.5-day postcoitum (p.c.) embryos. At 10.5 days p.c. and later, the pattern of beta-galactosidase activity diverges from the Hox-3.1 pattern, and an inappropriately high level of reporter gene expression is observed in posterior spinal ganglia. Inclusion of an additional 2 kb of upstream sequences is sufficient to suppress this aberrant expression in the developing spinal ganglia. Together, these results show that the control of early Hox-3.1 expression is complex, involving at least one positively acting and one negatively acting element.

Tags: Animal; Support, Non-U.S. Gov't; Support, U.S. Gov't, P.H.S.

Descriptors: *Gene Expression Regulation; *Genes, Homeobox; beta-Galactosidase--Genetics--GE; Drosophila--Genetics--GE; Embryo--Enzymology--EN; Escherichia coli--Enzymology--EN; Escherichia coli--Genetics--GE; Mice; Mice, Transgenic; Restriction Mapping
Enzyme No.: EC 3.2.1.23 (beta-Galactosidase)
Gene Symbol: Hox-3.1; lacZ

15/5/23 (Item 23 from file: 154)

DIALOG(R) File 154:MEDLINE(R)

(c) format only 1996 Knight-Ridder Info. All rts. reserv.

07387149 90294149

P-element-mediated enhancer detection allows rapid identification of developmentally regulated genes and cell specific markers in Drosophila.

Bellen HJ; Wilson C; Gibson G; Grossniklaus U; Pearson RK; O'Kane C; Gehring WJ

Dept. of Cell Biology, Biozentrum, Univ. of Basel, Switzerland.

J Physiol (Paris) (FRANCE) 1990, 84 (1) p33-41, ISSN 0021-7948
Journal Code: JRB

Languages: ENGLISH

Document type: JOURNAL ARTICLE

JOURNAL ANNOUNCEMENT: 9010

Subfile: INDEX MEDICUS

We have employed a new technique in Drosophila that allows in vivo detection of genomic regulatory elements using a beta-galactosidase reporter gene. A translational fusion of the reporter gene to the P-transposase gene, which is encoded by the P-transposon of Drosophila, places the expression of beta-galactosidase under the control of the weak P-transposase promoter. Flies carrying single insertions of this P-element construct at different locations in the Drosophila genome frequently stain for beta-galactosidase activity in a temporally and spatially restricted fashion in embryos, larvae and adult ovaries, reflecting the influence of nearby genomic regulatory elements on the P-transposase promoter. This technique is a powerful tool as it can be used to produce very many different cell markers and to isolate developmentally regulated genes in Drosophila. We discuss the implications of our results and the applications of the technique to further the study of Drosophila development.

Tags: Animal; Support, Non-U.S. Gov't

Descriptors: *Biological Markers--Analysis--AN; *Drosophila melanogaster--Genetics--GE; *DNA Insertion Elements; *Enhancer Elements (Genetics); *Regulatory Sequences, Nucleic Acid; beta-Galactosidase--Analysis--AN; Drosophila melanogaster--Embryology--EM; Embryo, Non-Mammalian--Analysis--AN; Gene Expression Regulation; Oogenesis--Genetics--GE

CAS Registry No.: 0 (Biological Markers); 0 (DNA Insertion Elements)

Enzyme No.: EC 3.2.1.23 (beta-Galactosidase)

15/5/24 (Item 24 from file: 154)
DIALOG(R) File 154:MEDLINE(R)
(c) format only 1996 Knight-Ridder Info. All rts. reserv.

07201680 90108680

P-element-mediated enhancer detection: an efficient method for isolating and characterizing developmentally regulated genes in *Drosophila*.

Wilson C; Pearson RK; Bellen HJ; O'Kane CJ; Grossniklaus U; Gehring WJ
Department of Cell Biology, Biozentrum, University of Basel, Switzerland.
Genes Dev (UNITED STATES) Sep 1989, 3 (9) p1301-13, ISSN 0890-9369

Journal Code: FN3

Languages: ENGLISH

Document type: JOURNAL ARTICLE

JOURNAL ANNOUNCEMENT: 9004

Subfile: INDEX MEDICUS

We describe a new approach for identifying and studying genes involved in *Drosophila* development. Single copies of an enhancer detector transposon, P[1ArB], have been introduced into flies at many different genomic locations. The beta-galactosidase reporter gene in this construct is influenced by a wide range of genomic transcriptional regulatory elements in its vicinity. Our results suggest that a significant proportion of these regulatory sequences are control elements of nearby *Drosophila* genes. These genes need not be disrupted for their regulatory elements to be identified by P[1ArB]. The P[1ArB] transposon has been designed to facilitate both rapid cloning and deletion analysis of genomic sequences into which it inserts. Therefore, the enhancer detection system is an efficient method of screening for genes primarily on the basis of their expression pattern and then rapidly analyzing those of particular interest at the molecular and genetic levels.

Tags: Animal; Support, Non-U.S. Gov't

Descriptors: **Drosophila melanogaster*--Genetics--GE; *DNA Insertion Elements; *Gene Expression Regulation; *Regulatory Sequences, Nucleic Acid ; beta-Galactosidase--Analysis--AN; *Drosophila melanogaster*--Embryology--EM ; Embryo, Non-Mammalian--Analysis--AN; Fetal Development; Genetic Vectors; Recombinant Fusion Proteins--Analysis--AN

CAS Registry No.: 0 (DNA Insertion Elements); 0 (Recombinant Fusion Proteins)

Enzyme No.: EC 3.2.1.23 (beta-Galactosidase)

15/5/25 (Item 25 from file: 154)
DIALOG(R) File 154:MEDLINE(R)
(c) format only 1996 Knight-Ridder Info. All rts. reserv.

07201679 90108679

P-element-mediated enhancer detection: a versatile method to study development in *Drosophila*.

Bellen HJ; O'Kane CJ; Wilson C; Grossniklaus U; Pearson RK; Gehring WJ
Department of Cell Biology, Biozentrum, University of Basel, Switzerland.
Genes Dev (UNITED STATES) Sep 1989, 3 (9) p1288-300, ISSN 0890-9369

Journal Code: FN3

Languages: ENGLISH

Document type: JOURNAL ARTICLE

JOURNAL ANNOUNCEMENT: 9004

Subfile: INDEX MEDICUS

We generated and characterized greater than 500 *Drosophila* strains that carry single copies of a novel P-element enhancer detector. In the majority of the strains, the beta-galactosidase reporter gene in the P-transposon

responds to nearby transcriptional regulatory sequences in the genome. A remarkable diversity of spatially and temporally regulated staining patterns is observed in embryos carrying different insertions. We selected numerous strains as markers for different embryonic organs, tissues, and cells. Many of these strains should allow the study of complex developmental processes, such as nervous system development, which have not been convenient to analyze previously. Also, we present genetic evidence that some of the detected regulatory elements control nearby *Drosophila* genes. In light of our results, we discuss the diversity and complexity of cis-acting regulatory elements in the genome and the general applications of the enhancer detector method for the study of *Drosophila* development.

Tags: Animal; Support, Non-U.S. Gov't

Descriptors: **Drosophila melanogaster*--Genetics--GE; *DNA Insertion Elements; *Enhancer Elements (Genetics); beta-Galactosidase--Analysis--AN; Biological Markers--Analysis--AN; *Drosophila melanogaster*--Embryology--EM; Embryo, Non-Mammalian--Analysis--AN; Gene Expression Regulation; Genetic Vectors; Recombinant Fusion Proteins--Analysis--AN

CAS Registry No.: 0 (Biological Markers); 0 (DNA Insertion Elements); 0 (Recombinant Fusion Proteins)

Enzyme No.: EC 3.2.1.23 (beta-Galactosidase)

15/5/26 (Item 26 from file: 154)

DIALOG(R) File 154:MEDLINE(R)

(c) format only 1996 Knight-Ridder Info. All rts. reserv.

07201678 90108678

Searching for pattern and mutation in the *Drosophila* genome with a P-lacZ vector.

Bier E; Vaessin H; Shepherd S; Lee K; McCall K; Barbel S; Ackerman L; Carretto R; Uemura T; Grell E; et al

Howard Hughes Medical Institute, San Francisco, California.

Genes Dev (UNITED STATES) Sep 1989, 3 (9) p1273-87, ISSN 0890-9369
Journal Code: FN3

Languages: ENGLISH

Document type: JOURNAL ARTICLE

JOURNAL ANNOUNCEMENT: 9004

Subfile: INDEX MEDICUS

A P-element vector has been constructed and used to generate lines of flies with single autosomal P-element insertions. The lines were analyzed in two ways: (1) the identification of cis-acting patterning information within the *Drosophila* genome, as revealed by a lacZ reporter gene within the P element, and (2) the isolation of lethal mutations. We examined 3768 independent lines for the expression of lacZ in embryos and looked among these lines for lethal mutations affecting embryonic neurogenesis. This type of screen appears to be an effective way to find new loci that may play a role in the development of the *Drosophila* nervous system.

Tags: Animal; Support, Non-U.S. Gov't

Descriptors: **Drosophila melanogaster*--Genetics--GE; *DNA Insertion Elements; beta-Galactosidase--Analysis--AN; *Drosophila melanogaster*--Embryology--EM; Embryo, Non-Mammalian--Analysis--AN; Evolution; Gene Expression Regulation; Genes, Lethal; Genetic Vectors; Mutation; Nervous System--Embryology--EM; Recombinant Fusion Proteins--Analysis--AN

CAS Registry No.: 0 (DNA Insertion Elements); 0 (Recombinant Fusion Proteins)

Enzyme No.: EC 3.2.1.23 (beta-Galactosidase)

15/5/27 (Item 27 from file: 154)
DIALOG(R)File 154:MEDLINE(R)
(c) format only 1996 Knight-Ridder Info. All rts. reserv.

07082425 89384425

Regulatory elements involved in the tissue-specific expression of the yellow gene of Drosophila.

Martin M; Meng YB; Chia W

Department of Biochemistry, School of Medical Sciences, University of Bristol, UK.

Mol Gen Genet (GERMANY, WEST) Jul 1989, 218 (1) p118-26, ISSN 0026-8925 Journal Code: NGP

Languages: ENGLISH

Document type: JOURNAL ARTICLE

JOURNAL ANNOUNCEMENT: 8912

Subfile: INDEX MEDICUS

We have assessed the DNA sequence requirements for the correct spatial pattern and phenotypic expression of y in the late embryo/larvae. The wild-type larval phenotype requires both the regions between -294 bp and -92 bp and a portion of the intron; the sequence element(s) located within the intron can act in a position independent manner to effect the wild-type larval phenotype. The larval expression pattern was examined by tissue experiments in situ and by staining germline transformants derived from various y/lacZ fusion constructs. The larval expression of y is restricted to the mouthparts, microsetae and anal plates. While the -495 bp to +194 bp region alone cannot effect a wild-type larval expression pattern, this region in conjunction with the intron appears to be sufficient to drive beta-gal expression in an essentially wild-type pattern. Our data further suggest that the -294 bp to -92 bp region contains elements which specify the larval pattern and that the element(s) in the intron normally act to enhance the level of expression necessary for the wild-type larval phenotype. We also present a phenotypic analysis of the adult cuticle structures of germline transformants derived from a variety of deletion and rearrangement constructs of the y gene. This analysis has revealed several new features associated with the regulation of y expression.

Tags: Animal; Support, Non-U.S. Gov't

Descriptors: *Drosophila--Genetics--GE; *Insect Hormones--Genetics--GE; *Regulatory Sequences, Nucleic Acid; beta-Galactosidase--Metabolism--ME; Base Sequence; Cloning, Molecular; Drosophila--Growth and Development--GD; DNA Insertion Elements; Gene Expression Regulation; Insect Hormones --Biosynthesis--BI; Introns; Lac Operon; Nucleic Acid Hybridization; Phenotype; Promoter Regions (Genetics); Restriction Mapping; Transcription, Genetic; Transformation, Genetic

CAS Registry No.: 0 (yellow locus protein, Drosophila); 0 (DNA Insertion Elements); 0 (Insect Hormones)

Enzyme No.: EC 3.2.1.23 (beta-Galactosidase)

15/5/28 (Item 28 from file: 154)
DIALOG(R)File 154:MEDLINE(R)
(c) format only 1996 Knight-Ridder Info. All rts. reserv.

06894899 89196899

Developmental expression of the Drosophila zeste gene and localization of zeste protein on polytene chromosomes.

Pirrotta V; Bickel S; Mariani C

Department of Cell Biology, Baylor College of Medicine, Houston, Texas 77030.

Genes Dev (UNITED STATES) Dec 1988, 2 (12B) p1839-50, ISSN 0890-9369
Journal Code: FN3
Contract/Grant No.: GM-34630
Languages: ENGLISH
Document type: JOURNAL ARTICLE
JOURNAL ANNOUNCEMENT: 8907
Subfile: INDEX MEDICUS

The expression of the zeste gene varies through the life cycle of the fly. Its transcription is most abundant in maternal RNA, declines to very low levels during larval growth, but rises again in late third instar larvae and pupae. Using transposons containing a zeste-lacZ gene, we found a corresponding variation in the tissue distribution of zeste from stage to stage. Nearly ubiquitous expression of the zeste-lacZ gene is found in late embryos and first instar larvae, but disappears almost completely except in brain and gonads by third instar larva. Shortly before pupation expression rises again in imaginal discs, Malpighian tubules, and salivary glands and again becomes nearly ubiquitous in pupae. zeste continues to be expressed in adult brain and gonads. We constructed flies carrying a zeste gene controlled by the heat shock promoter and studied the distribution of zeste protein in their polytene chromosomes as well as those of wild-type flies. Using affinity-purified anti-zeste antibodies, we find that wild-type salivary gland chromosomes contain about 60 strong bands of zeste immunofluorescence at specific cytological locations. After heat induction of larvae containing the hs-zeste gene, many hundreds of bands appear. These results suggest the involvement of zeste in the expression of a wide variety of genes at different developmental stages.

Tags: Animal; Support, Non-U.S. Gov't; Support, U.S. Gov't, Non-P.H.S.; Support, U.S. Gov't, P.H.S.

Descriptors: *Drosophila--Genetics--GE; *Gene Expression Regulation; *Genes, Structural; *Insect Hormones--Genetics--GE; beta-Galactosidase--Genetics--GE; Cloning, Molecular; Drosophila--Growth and Development--GD; DNA Insertion Elements; Fluorescent Antibody Technique; Heat-Shock Proteins--Genetics--GE; Larva--Genetics--GE; Mutation; Pupa--Genetics--GE; Salivary Glands--Cytology--CY

CAS Registry No.: 0 (DNA Insertion Elements); 0 (Heat-Shock Proteins); 0 (Insect Hormones)
Enzyme No.: EC 3.2.1.23 (beta-Galactosidase)

15/5/29 (Item 29 from file: 154)

DIALOG(R) File 154:MEDLINE(R)

(c) format only 1996 Knight-Ridder Info. All rts. reserv.

06809103 89111103

Expression of an engrailed-like gene during development of the early embryonic chick nervous system.

Gardner CA; Darnell DK; Poole SJ; Ordahl CP; Barald KF

Department of Anatomy and Cell Biology, University of Michigan, Ann Arbor 48109.

J Neurosci Res (UNITED STATES) Oct-Dec 1988, 21 (2-4) p426-37, ISSN 0360-4012 Journal Code: KAC

Languages: ENGLISH

Document type: JOURNAL ARTICLE

JOURNAL ANNOUNCEMENT: 8905

Subfile: INDEX MEDICUS

The engrailed gene has been identified in Drosophila as an important developmental gene involved in the control of segmentation. Here we describe the embryonic expression of a chicken gene, ChickEn (Darnell et

al.: J Cell Biol 103(5):311a, 1986), which contains homology to the Drosophila engrailed gene. Northern blots of early chick embryo tissue poly(A)+ RNA resulted in hybridization to at least three bands expressed predominantly in the brain/head region when probed with ChickEn genomic fragments. Eight cDNA clones generated from embryonic day 6 (stage 29-30) chick brain poly(A)+ RNA are identical in their nucleotide sequence with the ChickEn genomic clone. In situ hybridization to sections of 4-day (stage 24) embryos indicated that ChickEn transcripts were concentrated in the posterior mesencephalon and anterior metencephalon. In cultures of chick cranial neural crest cells (eight to nine somites; stage 9) ChickEn transcripts were localized in a subset (approx. 8%) of cells examined after 2 days in culture. A mouse monoclonal antibody, inv-4D9D4, made by Coleman and Kornberg recognizes the engrailed-like homeo domain of the engrailed and invected proteins (Martin-Blanco, Coleman, and Kornberg, personal communication). Patel, Coleman, Kornberg and Goodman (unpublished) have shown that this antibody binds to the hindbrain of 2-day-old chick embryos. We have confirmed these results and shown that this antibody binds to the same region of 4-day (stage 24) chick brains that in situ hybridization showed contained ChickEn transcripts. This antibody also recognizes a homeo domain-containing ChickEn peptide expressed as a beta-galactosidase fusion protein in Drosophila cell culture. We have not detected ChickEn protein in any tissue prior to eight to nine somites (stage 9). These results delineate the major expression pattern of the ChickEn gene during early (prior to stage 30) embryonic development in the chick.

Tags: Animal; Support, Non-U.S. Gov't; Support, U.S. Gov't, Non-P.H.S.; Support, U.S. Gov't, P.H.S.

Descriptors: *Fetal Development; *Gene Expression Regulation; *Genes, Structural; *Neural Crest--Metabolism--ME; Chick Embryo; Chromosome Mapping; Nucleic Acid Hybridization; RNA--Metabolism--ME

CAS Registry No.: 0 (RNA)

15/5/30 (Item 30 from file: 154)

DIALOG(R) File 154:MEDLINE(R)

(c) format only 1996 Knight-Ridder Info. All rts. reserv.

06805994 89107994

Control elements of the P2 promoter of the Antennapedia gene.

Boulet AM; Scott MP

Department of Molecular, Cellular, and Developmental Biology, University of Colorado, Boulder 80302.

Genes Dev (UNITED STATES) Dec 1988, 2 (12A) p1600-14, ISSN 0890-9369

Journal Code: FN3

Contract/Grant No.: 18163

Languages: ENGLISH

Document type: JOURNAL ARTICLE

JOURNAL ANNOUNCEMENT: 8905

Subfile: INDEX MEDICUS

Antennapedia (Antp), a homeotic gene of Drosophila required for proper differentiation of the thorax of the fly, is expressed in complex spatial patterns during development. The gene is greater than 100 kb long and has two independently regulated promoters. To characterize cis-acting regulatory elements responsible for the expression pattern, fusions of the Antp promoter 2 cap site and upstream sequences to an Adh-lacZ gene were introduced into flies. A 10-kb sequence directs beta-galactosidase production in a pattern that closely resembles the endogenous P2 pattern. Transcription from the 10-kb fusions is regulated by three genes that regulate Antp transcription. Control elements, including a target of action

of homeo-domain-containing proteins, were mapped by deleting parts of the 10-kb sequence.

Tags: Animal; Support, Non-U.S. Gov't; Support, U.S. Gov't, P.H.S.

Descriptors: *Drosophila--Genetics--GE; *Genes, Homeobox; *Promoter Regions (Genetics); *Regulatory Sequences, Nucleic Acid; beta-Galactosidase--Biosynthesis--BI; beta-Galactosidase--Genetics--GE; Cloning, Molecular; Drosophila--Embryology--EM; Escherichia coli--Genetics--GE; Mutation; Plasmids; Recombinant Fusion Proteins--Biosynthesis--BI; Recombinant Fusion Proteins--Genetics--GE; Restriction Mapping; Transformation, Genetic
CAS Registry No.: 0 (Plasmids); 0 (Recombinant Fusion Proteins)
Enzyme No.: EC 3.2.1.23 (beta-Galactosidase)

15/5/31 (Item 31 from file: 154)

DIALOG(R) File 154:MEDLINE(R)

(c) format only 1996 Knight-Ridder Info. All rts. reserv.

06551887 88196887

Spatial and temporal expression of the period gene in Drosophila melanogaster.

Liu X; Lorenz L; Yu QN; Hall JC; Rosbash M

Department of Biology, Brandeis University, Waltham, Massachusetts 02254.

Genes Dev (UNITED STATES) Feb 1988, 2 (2) p228-38, ISSN 0890-9369

Journal Code: FN3

Contract/Grant No.: GM33205

Languages: ENGLISH

Document type: JOURNAL ARTICLE

JOURNAL ANNOUNCEMENT: 8808

Subfile: INDEX MEDICUS

The temporal and spatial expression of the period gene of Drosophila melanogaster has been analyzed by examining the expression of a per beta-galactosidase fusion gene in transformants and by in situ hybridization experiments with wild-type flies. Several strains of Drosophila melanogaster, transformed with the fusion gene, have been generated. The gene is active in mid-late embryos in the midline of the nervous system. Thereafter, beta-galactosidase activity is undetectable until the pupal stage when the prothoracic gland-corpora allata and the optic lobes are beta-galactosidase positive. In adults a surprisingly large number of tissues stain positively, including antennae, proboscis, eyes, optic lobes, cells of the central brain, cells of the thoracic ganglia, gut, Malpighian tubules, and ovarian follicle cells. The temporal pattern of expression agrees well with previous estimates made from developmental Northern blots with RNA extracted from wild-type animals. We suggest that many of the tissues that express the per gene contain their own intrinsic oscillator activity.

Tags: Animal; Support, U.S. Gov't, P.H.S.

Descriptors: *Drosophila melanogaster--Genetics--GE; *Gene Expression Regulation; beta-Galactosidase--Genetics--GE; Circadian Rhythm; Cloning, Molecular; Drosophila melanogaster--Growth and Development--GD; Drosophila melanogaster--Physiology--PH; Mutation; Transformation, Genetic

Enzyme No.: EC 3.2.1.23 (beta-Galactosidase)

15/5/32 (Item 32 from file: 154)

DIALOG(R) File 154:MEDLINE(R)

(c) format only 1996 Knight-Ridder Info. All rts. reserv.

06452440 88097440

Detection in situ of genomic regulatory elements in Drosophila.

O'Kane CJ; Gehring WJ

Department of Cell Biology, University of Basel, Switzerland.

Proc Natl Acad Sci U S A (UNITED STATES) Dec 1987, 84 (24) p9123-7,

ISSN 0027-8424 Journal Code: PV3

Languages: ENGLISH

Document type: JOURNAL ARTICLE

JOURNAL ANNOUNCEMENT: 8804

Subfile: INDEX MEDICUS

We have developed an approach for the in situ detection of genomic elements that regulate transcription in Drosophila melanogaster. The approach is analogous to a powerful method of bacterial genetics, the random generation of operon fusions, that enables the isolation and characterization of genes simply by knowing or postulating their pattern of expression; it is not necessary initially to screen for mutant phenotypes. To apply this approach to Drosophila, we have used the expression of the lacZ gene of Escherichia coli from the P-element promoter in germ-line transformant flies to screen for chromosomal elements that can act at a distance to stimulate expression from this apparently weak promoter. Of 49 transformed fly lines obtained, approximately 70% show some type of spatially regulated expression of the lacZ gene in embryos; many of these express lacZ specifically in the nervous system. The P-lacZ fusion gene is, therefore, an efficient tool for the recovery of elements that may regulate gene expression in Drosophila and for the generation of a wide variety of cell-type-specific markers.

Tags: Animal; Support, Non-U.S. Gov't

Descriptors: *Drosophila melanogaster--Genetics--GE; *DNA Insertion Elements; *Gene Expression Regulation; *Promoter Regions (Genetics); *Regulatory Sequences, Nucleic Acid; beta-Galactosidase--Diagnostic Use--DU; Drosophila melanogaster--Embryology--EM; DNA, Recombinant; Nervous System--Physiology--PH; Tissue Distribution

CAS Registry No.: 0 (DNA Insertion Elements); 0 (DNA, Recombinant)

Enzyme No.: EC 3.2.1.23 (beta-Galactosidase)

15/5/33 (Item 33 from file: 154)

DIALOG(R) File 154:MEDLINE(R)

(c) format only 1996 Knight-Ridder Info. All rts. reserv.

06313209 87287209

Borders of parasegments in Drosophila embryos are delimited by the fushi tarazu and even-skipped genes.

Lawrence PA; Johnston P; Macdonald P; Struhl G

Nature (ENGLAND) Jul 30-Aug 5 1987, 328 (6129) p440-2, ISSN 0028-0836

Journal Code: NSC

Languages: ENGLISH

Document type: JOURNAL ARTICLE

JOURNAL ANNOUNCEMENT: 8711

Subfile: INDEX MEDICUS

One of the earliest molecular signs of segmentation in Drosophila embryos is the striped expression of some pair-rule genes during the blastoderm stage. Two of these genes, fushi-tarazu (ftz) and even-skipped (eve) are expressed during this stage in complementary patterns of seven stripes which develop and disappear in concert. Here, we map the cells expressing each of these two pair-rule genes with respect to the 14 stripes of cells expressing the engrailed gene. We find that both ftz and eve generate stripes which have sharp boundaries at the anterior margin, but fade away posteriorly. The anterior boundaries correspond cell by cell with the

anterior boundaries of expression of the engrailed gene. We therefore suggest that a key function of early ftz and eve gene activity is the formation of a sharp stable boundary at the anterior margin of each stripe. These boundary lines, rather than the narrowing zonal stripes, would delimit the anterior boundaries of engrailed and other homoeotic genes and thereby subdivide the embryo into parasegments.

Tags: Animal; Support, Non-U.S. Gov't; Support, U.S. Gov't, P.H.S.

Descriptors: *Drosophila--Genetics--GE; *Genes, Homeobox; beta-Galactosidase--Genetics--GE; Blastoderm--Metabolism--ME; Blastoderm--Ultrastructure--UL; Drosophila--Embryology--EM; DNA, Recombinant; Gene Expression Regulation; Histochemistry; Immunologic Tests

CAS Registry No.: 0 (DNA, Recombinant)

Enzyme No.: EC 3.2.1.23 (beta-Galactosidase)

15/5/34 (Item 1 from file: 55)
DIALOG(R)File 55:BIOSIS PREVIEWS(R)
(c) 1996 BIOSIS. All rts. reserv.

11510853 BIOSIS Number: 98110853

Regulation of DNA replication-related gene expression during Drosophila development

Yamaguchi M; Hirose F; Matsukage A

Lab. Cell Biol., Aichi Cancer Cent Res. Inst., Nagoya 464, Japan

Cell Structure and Function 19 (6). 1994. 463.

Full Journal Title: Forty-seventh Annual Meeting of the Japan Society for Cell Biology, Nagasaki, Japan, September 28-30, 1994. Cell Structure and Function

ISSN: 0386-7196

Language: ENGLISH

Document Type: CONFERENCE PAPER

Print Number: Biological Abstracts/RRM Vol. 047 Iss. 003 Ref. 045215

Descriptors/Keywords: MEETING ABSTRACT; DROSOPHILA; PROLIFERATING CELL NUCLEAR ANTIGEN; MESSENGER RNA; BETA-GALACTOSIDASE; PROMOTER; EMBRYOGENESIS; DEVELOPMENT

Concept Codes:

*03506 Genetics and Cytogenetics-Animal

*10300 Replication, Transcription, Translation

*10808 Enzymes-Physiological Studies

*25502 Developmental Biology-Embryology-General and Descriptive

*25508 Developmental Biology-Embryology-Morphogenesis, General

*34502 Immunology and Immunochemistry-General; Methods

*64076 Invertebrata, Comparative and Experimental Morphology, Physiology and Pathology-Insecta-Physiology

00520 General Biology-Symposia, Transactions and Proceedings of Conferences, Congresses, Review Annuals

10062 Biochemical Studies-Nucleic Acids, Purines and Pyrimidines

10064 Biochemical Studies-Proteins, Peptides and Amino Acids

Biosystematic Codes:

75314 Diptera

Super Taxa:

Animals; Invertebrates; Arthropods; Insects

15/5/35 (Item 2 from file: 55)
DIALOG(R)File 55:BIOSIS PREVIEWS(R)
(c) 1996 BIOSIS. All rts. reserv.

10405441 BIOSIS Number: 96005441

DEVELOPMENTAL REGULATORY ELEMENTS IN THE 5' FLANKING DNA OF THE
DROSOPHILA CHOLINE ACETYLTRANSFERASE GENE

KITAMOTO T; SALVATERRA P M

DIV. NEUROSCI., BECKMAN RESEARCH INST. CITY HOPE, 1450 EAST DUARTE RD.,
DUARTE, CA 91010, USA.

ROUX'S ARCH DEV BIOL 202 (3). 1993. 159-169. CODEN: RADBE

Full Journal Title: Roux'S Archives of Developmental Biology

Language: ENGLISH

Choline acetyltransferase (ChAT, EC 2.3.1.6) catalyzes the production of the neurotransmitter acetylcholine, and is an essential factor for neurons to be cholinergic. We have analyzed regulation of the Drosophila ChAT gene during development by examining the .beta.-galactosidase expression pattern in transformed lines carrying different lengths of 5' flanking DNA fused to a lacZ reporter gene. The largest fragment tested, 7.4 kb, resulted in the most extensive expression pattern in embryonic and larval nervous system and likely reflects all the cis-regulatory elements necessary for ChAT expression. We also found that 5' flanking DNA located between 3.3 kb and 1.2 kb is essential for the receptor gene expression in most of the segmentally arranged embryonic sensory neurons as well as other distinct cells in the CNS. The existence of negative regulatory elements was suggested by the observation that differentiating photoreceptor cells in eye imaginal discs showed the reporter gene expression in several 1.2 kb and 3.3 kb transformants but not in 7.4 kb transformants. Furthermore, we have fused the 5' flanking DNA fragments to a wild type ChAT cDNA and used these constructs to transform Drosophila with a Cha mutant background. Surprisingly, even though different amounts of 5' flanking DNA resulted in different spatial expression patterns, all of the positively expressing cDNA transformed lines were rescued from lethality. Our results suggest that developmental expression of the ChAT gene is regulated both positively and negatively by the combined action of several elements located in the 7.4 kb upstream region, and that the more distal 5' flanking DNA is not necessary for embryonic survival and development to adult flies.

Descriptors/Keywords: EMBRYO LARVAE ACETYLCHOLINE CHOLINE ACETYLTRANSFERASE

EC 2.3.1.6 NEURON CENTRAL NERVOUS SYSTEM

Concept Codes:

- *02506 Cytology and Cytochemistry-Animal
- *03506 Genetics and Cytogenetics-Animal
- *10062 Biochemical Studies-Nucleic Acids, Purines and Pyrimidines
- *10808 Enzymes-Physiological Studies
- *20504 Nervous System-Physiology and Biochemistry
- *25502 Developmental Biology-Embryology-General and Descriptive
- *25508 Developmental Biology-Embryology-Morphogenesis, General
- *64076 Invertebrata, Comparative and Experimental Morphology, Physiology and Pathology-Insecta-Physiology
- 10064 Biochemical Studies-Proteins, Peptides and Amino Acids

Biosystematic Codes:

75314 Diptera

Super Taxa:

Animals; Invertebrates; Arthropods; Insects

15/5/36 (Item 3 from file: 55)

DIALOG(R)File 55:BIOSIS PREVIEWS(R)

(c) 1996 BIOSIS. All rts. reserv.

10086391 BIOSIS Number: 95086391

A CIS-ELEMENT MEDIATING ULTRABITHORAX AUTOREGULATION IN THE CENTRAL

NERVOUS SYSTEM

CHRISTEN B; BIENZ M

MRC LABORATORY MOLECULAR BIOLOGY, HILLS ROAD, CAMBRIDGE CB2 2QH, UK.

MECH DEV 39 (1-2). 1992. 73-80. CODEN: MEDVE

Language: ENGLISH

We dissected an upstream control region (a BXD fragment) from the homeotic gene Ultrabithorax (Ubx) of Drosophila which confers a Ubx-like expression pattern in the embryonic ectoderm. We found several distinct enhancer elements spread through the whole BXD fragment each of which is active in transformed embryos, mediating a different pattern of .beta.-galactosidase expression in the ventral nerve cord. The strongest of these patterns mimics Ubx expression within the Ubx domain. This pattern is strictly dependent on Ubx function. Thus, the BXD control region contains a Ubx response element, suggesting that positive autoregulation of Ubx may occur in the central nervous system of the developing embryo.

Descriptors/Keywords: DROSOPHILA EMBRYO BETA GALACTOSIDASE VENTRAL NERVE CORD TRANSCRIPTIONAL ACTIVATION DEVELOPMENT

Concept Codes:

- *03506 Genetics and Cytogenetics-Animal
- *10300 Replication, Transcription, Translation
- *10808 Enzymes-Physiological Studies
- *20504 Nervous System-Physiology and Biochemistry
- *25502 Developmental Biology-Embryology-General and Descriptive
- *25508 Developmental Biology-Embryology-Morphogenesis, General
- *64076 Invertebrata, Comparative and Experimental Morphology, Physiology and Pathology-Insecta-Physiology
- 10064 Biochemical Studies-Proteins, Peptides and Amino Acids

Biosystematic Codes:

75314 Diptera

Super Taxa:

Animals; Invertebrates; Arthropods; Insects

?s patched and (gal or galactosidase)

388 PATCHED

7514 GAL

16555 GALACTOSIDASE

S16 4 PATCHED AND (GAL OR GALACTOSIDASE)

?t s16/6/1-4

16/6/1 (Item 1 from file: 154)

09241451 95171451

Protein kinase A and hedgehog signaling in Drosophila limb development.

16/6/2 (Item 2 from file: 154)

07664999 91183999

The Drosophila segment polarity gene patched is involved in a position-signalling mechanism in imaginal discs.

16/6/3 (Item 3 from file: 154)

05982167 86283167

The influence of serosal patch size on the growth of small intestinal neomucosa.

16/6/4 (Item 1 from file: 55)

7767662 BIOSIS Number: 90135662

THE DROSOPHILA SEGMENT POLARITY GENE PATCHED IS INVOLVED IN A
POSITION-SIGNALING MECHANISM IN IMAGINAL DISCS
?t s16/7/1-3

16/7/1 (Item 1 from file: 154)

DIALOG(R) File 154:MEDLINE(R)

(c) format only 1996 Knight-Ridder Info. All rts. reserv.

09241451 95171451

Protein kinase A and hedgehog signaling in Drosophila limb development.

Jiang J; Struhl G

Howard Hughes Medical Institute, Department of Genetics and Development
Columbia University College of Physicians and Surgeons, New York, New York
10032.

Cell (UNITED STATES) Feb 24 1995, 80 (4) p563-72, ISSN 0092-8674

Journal Code: CQ4

Languages: ENGLISH

Document type: JOURNAL ARTICLE

The Drosophila hedgehog (hh) gene encodes a secreted protein involved in organizing growth and patterning in many developmental processes. Hh appears to act by inducing the localized expression of at least two other signaling molecules, decapentaplegic (dpp) and wingless (wg), which then govern cell proliferation and patterning in surrounding tissue. Here, we demonstrate that cyclic AMP (cAMP)-dependent protein kinase A (PKA) is essential during limb development to prevent inappropriate dpp and wg expression. We also show that a constitutively active form of PKA can prevent inappropriate dpp and wg expression, but does not interfere with their normal induction by hh. We propose that the basal activity of PKA imposes a block on the transcription of dpp and wg and that hh exerts its organizing influence by alleviating this block.

16/7/2 (Item 2 from file: 154)

DIALOG(R) File 154:MEDLINE(R)

(c) format only 1996 Knight-Ridder Info. All rts. reserv.

07664999 91183999

The Drosophila segment polarity gene patched is involved in a position-signalling mechanism in imaginal discs.

Phillips RG; Roberts IJ; Ingham PW; Whittle JR

School of Biological Sciences, University of Sussex, Brighton, UK.

Development (ENGLAND) Sep 1990, 110 (1) p105-14, ISSN 0950-1991

Journal Code: ECW

Languages: ENGLISH

Document type: JOURNAL ARTICLE

We demonstrate the role of the segment polarity gene patched (ptc) in patterning in the cuticle of the adult fly. Genetic mosaics of a lethal allele of patched show that the contribution of patched varies in a position-specific manner, defining three regions in the wing where ptc clones, respectively, behave as wild-type cells, affect vein formation, or are rarely recovered. Analysis of twin clones demonstrates that the reduced clone frequency results from a proliferation failure or cell loss. In the region where clones upset venation, they autonomously fail to form veins and also non-autonomously induce ectopic veins in adjacent wild-type cells. In heteroallelic combinations with lethal alleles, two viable alleles produce distinct phenotypes: (1) loss of structures and mirror-image

duplications in the region where patched clones fail to proliferate; (2) vein abnormalities in the anterior compartment. We propose that these differences reflect independently mutable functions within the gene. We show the pattern of patched transcription in the developing imaginal wing disc in relation to the expression of certain other reporter genes using a novel double-labelling method combining non-radioactive detection of in situ hybridization with beta-galactosidase detection. The patched transcript is present throughout the anterior compartment, with a stripe of maximal intensity along the A/P compartment border extending into the posterior compartment. We propose that the patched product is a component of a cell-to-cell position-signalling mechanism, a proposal consistent with the predicted structure of the patched protein.

16/7/3 (Item 3 from file: 154)
 DIALOG(R) File 154:MEDLINE(R)
 (c) format only 1996 Knight-Ridder Info. All rts. reserv.

05982167 86283167

The influence of serosal patch size on the growth of small intestinal neomucosa.

Bragg LE; Thompson JS

J Surg Res (UNITED STATES) May 1986, 40 (5) p426-31, ISSN 0022-4804

Journal Code: K7B

Languages: ENGLISH

Document type: JOURNAL ARTICLE

Several factors might affect the growth of neomucosa after serosal patching of small intestinal defects. Often only short segments of small intestine can be patched because of limited serosal surface and anatomic factors. The purpose of this study was to determine the influence of patch size on neomucosal growth. Twenty male New Zealand white rabbits underwent patching with colon serosa of either a 2 X 15-cm distal ileal defect (n = 10) or three 2 X 5-cm ileal defects (n = 10). There was significantly greater coverage of the patched defect by neomucosa in the triple patch group (99.4% vs 93.1% P less than 0.005) and significantly more of the smaller defects were completely covered by neomucosa than the larger defects (12 of 15 vs 0 of 5, P less than 0.05) at 8 weeks. The final area of the defect was 27.5 and 32.8% of the initial patched area respectively for the single and triple patches. Microscopically there was no difference in villous height or crypt depth, but crypt density was significantly greater in the triple group (207 +/- 11 vs 186 +/- 17 crypts/mm, P less than 0.05). In vitro glucose uptake and disaccharidase activity were similar in both groups. Patching multiple small intestinal defects results in more rapid neomucosal growth than a single large defect of the same surface area. This might be due to a greater circumference exposed to surrounding normal mucosa with a resultant increase in crypt density. Since function and villous development of the neomucosa are similar, multiple patches should result in a greater increase in absorptive capacity.

?ds

Set	Items	Description
S1	497	AU="SCOTT M" OR AU="SCOTT M P" OR AU="SCOTT MP"
S2	6	S1 AND PATCHED
S3	4	RD (unique items)
S4	388	PATCHED
S5	157.	S4 AND (HUMAN OR MOUSE OR MOSQUITO OR BUTTERFLY OR BEETLE)
S6	14	S5 AND (GENE? OR CLONE? OR DNA?)

S7 11 RD (unique items)
 S8 4 DROSOPHILA AND (GAL OR GALACTOSIDASE)
 S9 502 DROSOPHILA AND (GAL OR GALACTOSIDASE)
 S10 210 S9 AND EMBRYO?
 S11 0 S10A ND DEVELOP?
 S12 125 S10 AND DEVELOP?
 S13 181 DROSOPHILA(10N) (GAL OR GALACTOSIDASE)
 S14 40 S13 AND DEVELOP? AND EMBRYO?
 S15 36 RD (unique items)
 S16 4 PATCHED AND (GAL OR GALACTOSIDASE)
 ?s s4 and review?

388 S4
 391802 REVIEW?
 S17 10 S4 AND REVIEW?
 ?rd

...completed examining records
 S18 8 RD (unique items)
 ?t s18/6/1-8

18/6/1 (Item 1 from file: 154)
 09145457 95075457
 Distinct pathways for autocrine and paracrine Wingless signalling in Drosophila embryos.

18/6/2 (Item 2 from file: 154)
 07988474 92126474
 Blowout of carotid venous patch angioplasty.

18/6/3 (Item 3 from file: 154)
 07961663 92099663
 Ventricular septal defect with tricuspid pouch with and without transposition. Anatomic and surgical considerations.

18/6/4 (Item 4 from file: 154)
 07944050 92082050
 A review of carotid endarterectomy at a large teaching hospital.

18/6/5 (Item 5 from file: 154)
 07660744 91179744
 Perilymph fistulas: the House Ear Clinic experience.

18/6/6 (Item 1 from file: 55)
 11821956 BIOSIS Number: 98421956
 A consideration of epistemology in systematic biology, with special reference to species
 Print Number: Biological Abstracts Vol. 100 Iss. 007 Ref. 099548

18/6/7 (Item 2 from file: 55)
 7236094 BIOSIS Number: 38016615

GENES THAT CONTROL PATTERN FORMATION DURING DEVELOPMENT

18/6/8 (Item 3 from file: 55)
5862168 BIOSIS Number: 83124475

A CASE OF CONTACT DERMATITIS DUE TO BETAMETHASONE VALERATE WITH A REVIEW
OF JAPANESE ARTICLES
?t s18/7/7

18/7/7 (Item 2 from file: 55)
DIALOG(R)File 55:BIOSIS PREVIEWS(R)
(c) 1996 BIOSIS. All rts. reserv.

7236094 BIOSIS Number: 38016615

GENES THAT CONTROL PATTERN FORMATION DURING DEVELOPMENT
SCOTT M P; HAYASHI S; WINSLOW G M; HOOPER J E; SONODA S
DEP. MOL. CELL. DEV. BIOL., UNIV. COLO., BOULDER, COLO. 80309, USA.
CAPECCHI, M. R. (ED.). CURRENT COMMUNICATIONS IN MOLECULAR BIOLOGY:
MOLECULAR GENETICS OF EARLY DROSOPHILA AND MOUSE DEVELOPMENT; MEETING, COLD
SPRING HARBOR, NEW YORK, USA, APRIL 20-23, 1989. XIII+141P. COLD SPRING
HARBOR LABORATORY PRESS: COLD SPRING HARBOR, NEW YORK, USA. ILLUS. PAPER.
ISBN 0-87969-339-8. 0 (0). 1989. 7-10. CODEN: 28365

Language: ENGLISH
?s patched and mammal?

388 PATCHED
137749 MAMMAL?
S19 13 PATCHED AND MAMMAL?
?rd

...completed examining records
S20 12 RD (unique items)
?t s20/6/1-12

20/6/1 (Item 1 from file: 154)
09506686 96028286
Morphogenetic signalling. Responses to hedgehog.

20/6/2 (Item 2 from file: 154)
09394383 95324383
Subcellular localization of the segment polarity protein patched suggests
an interaction with the wingless reception complex in Drosophila embryos.

20/6/3 (Item 3 from file: 154)
09081589 95011589
Localized expression of sloppy paired protein maintains the polarity of
Drosophila parasegments.

20/6/4 (Item 4 from file: 154)
08848979 94163979
A central role for epidermal segment border cells in the induction of
muscle patterning in the Drosophila embryo.

20/6/5 (Item 5 from file: 154)

08801069 94116069

Hedgehog is a signaling protein with a key role in patterning *Drosophila* imaginal discs.

20/6/6 (Item 6 from file: 154)

08668871 93378871

Contrasting distributions of patched and hedgehog proteins in the *Drosophila* embryo.

20/6/7 (Item 7 from file: 154)

08460175 93170175

The consequences of ubiquitous expression of the wingless gene in the *Drosophila* embryo.

20/6/8 (Item 8 from file: 154)

07151658 90058658

The *Drosophila* patched gene encodes a putative membrane protein required for segmental patterning.

20/6/9 (Item 9 from file: 154)

07108164 90015164

A protein with several possible membrane-spanning domains encoded by the *Drosophila* segment polarity gene patched.

20/6/10 (Item 10 from file: 154)

07004625 89306625

The role of segment polarity genes during *Drosophila* neurogenesis.

20/6/11 (Item 11 from file: 154)

06729922 89031922

Patch clamp analysis of Na channel gating in mammalian myocardium: reconstruction of double pulse inactivation and voltage dependence of Na currents.

20/6/12 (Item 12 from file: 154)

06712381 89014381

Patch clamp analysis of chemically activated and modulated ionic channels in isolated mammalian cardiomyocytes.

?

=> s patched

L1 1183 PATCHED

=> s patched(5a) (mouse or mammalian or butterfly or beetle)

1183 PATCHED

22421 MOUSE

14401 MAMMALIAN

7917 BUTTERFLY

2388 BEETLE

L2 1 PATCHED(5A) (MOUSE OR MAMMALIAN OR BUTTERFLY OR BEETLE)

=> d l2 cit ab

1. 4,556,560, Dec. 3, 1985, Methods for the treatment and prophylaxis of diaper rash and diaper dermatitis; Kent W. Buckingham, 424/641; 15/206; 514/494, 502, 865; 604/360 [IMAGE AVAILABLE]

US PAT NO: 4,556,560 [IMAGE AVAILABLE]

L2: 1 of 1

ABSTRACT:

Methods for the treatment and prevention of diaper rash and diaper dermatitis caused by the prolonged contact of human skin with body waste are disclosed. The methods of the present invention employ the topical application of a minimum inhibitory concentration of a pharmaceutically-acceptable lipase-inhibiting agent to the area in need of such treatment, or the area where prevention is desired. The lipase-inhibiting agent is preferably a water-soluble metallic salt, such as ZnCl₂, and is preferably applied in combination with a barrier-like vehicle. The effectiveness of these methods is surprising in light of the present confusion and controversy surrounding the actual causes of diaper rash, and the heretofore unrecognized role of lipase as a factor in the cause of diaper rash and diaper dermatitis.

=> s patched(5a) (gene# or clon? or DNA)

1183 PATCHED

9994 GENE#

11736 CLON?

13758 DNA

L3 1 PATCHED(5A) (GENE# OR CLON? OR DNA)

=> d l3 cit ab

1. 5,066,596, Nov. 19, 1991, Bacterial strains harboring cloned genes controlling Vibrio cholerae O-antigen biosynthesis; Paul A. Manning, et al., 435/252.33; 424/200.1, 235.1, 242.1, 257.1, 258.1, 261.1; 435/69.1, 69.3, 91.41, 172.1, 172.3, 320.1, 848; 536/24.1; 935/6, 9, 22, 26, 60, 73 [IMAGE AVAILABLE]

US PAT NO: 5,066,596 [IMAGE AVAILABLE]

L3: 1 of 1

ABSTRACT:

The invention relates to a fragment of DNA containing genes encoding the synthesis of the O-antigen of Vibrio cholerae serotypes Inaba or Ogawa and being at least 16 kb in length. The invention further related to a cosmid comprising a cloned DNA fragment containing genes encoding the synthesis of O-antigen of Vibrio cholerae serotypes Inaba or Ogawa and to a strain of E.coli that includes the fragment.

=> s patched(5a) drosophila

1183 PATCHED

974 DROSOPHILA

L4 0 PATCHED(5A) DROSOPHILA

=> s patched and drosophila

1183 PATCHED

974 DROSOPHILA

L5 5 PATCHED AND DROSOPHILA

=> d 15 1-5

1. 5,449,755, Sep. 12, 1995, Human cyclin E; James M. Roberts, et al., 530/350, 387.1; 536/23.5 [IMAGE AVAILABLE]

2. 5,350,671, Sep. 27, 1994, HCV immunoassays employing C domain antigens; Michael Houghton, et al., 435/5, 6, 975; 436/512, 518; 530/300, 326, 327, 328, 812, 826; 930/220, 223 [IMAGE AVAILABLE]

3. 5,324,659, Jun. 28, 1994, Yeast mutants useful for indentifying immunosupressants; Stephen A. Parent, et al., 435/255.2, 942 [IMAGE AVAILABLE]

4. 5,198,346, Mar. 30, 1993, Generation and selection of novel DNA-binding proteins and polypeptides; Robert C. Ladner, et al., 435/69.1, 172.3, 252.3, 320.1 [IMAGE AVAILABLE]

5. 5,096,815, Mar. 17, 1992, Generation and selection of novel DNA-binding proteins and polypeptides; Robert C. Ladner, et al., 435/69.1, 172.3, 252.3, 320.1 [IMAGE AVAILABLE]

=> s patched and (gene# or clon? or DNA)

1183 PATCHED

9994 GENE#

11736 CLON?

13758 DNA

L6 61 PATCHED AND (GENE# OR CLON? OR DNA)

=> s l6 and sequence#

271178 SEQUENCE#

L7 50 L6 AND SEQUENCE#

=> s l7 and drosophila

974 DROSOPHILA

L8 5 L7 AND DROSOPHILA

=> d 18 1-5

1. 5,449,755, Sep. 12, 1995, Human cyclin E; James M. Roberts, et al., 530/350, 387.1; 536/23.5 [IMAGE AVAILABLE]

2. 5,350,671, Sep. 27, 1994, HCV immunoassays employing C domain antigens; Michael Houghton, et al., 435/5, 6, 975; 436/512, 518; 530/300, 326, 327, 328, 812, 826; 930/220, 223 [IMAGE AVAILABLE]

3. 5,324,659, Jun. 28, 1994, Yeast mutants useful for indentifying immunosupressants; Stephen A. Parent, et al., 435/255.2, 942 [IMAGE AVAILABLE]

4. 5,198,346, Mar. 30, 1993, Generation and selection of novel **DNA**-binding proteins and polypeptides; Robert C. Ladner, et al., 435/69.1, 172.3, 252.3, 320.1 [IMAGE AVAILABLE]

5. 5,096,815, Mar. 17, 1992, Generation and selection of novel **DNA**-binding proteins and polypeptides; Robert C. Ladner, et al., 435/69.1, 172.3, 252.3, 320.1 [IMAGE AVAILABLE]

=> s l7 and (development or embryo?)

194785 DEVELOPMENT

4636 EMBRYO?

L9 31 L7 AND (DEVELOPMENT OR EMBRYO?)

=> d 19 1-10

1. 5,470,971, Nov. 28, 1995, Stress-induced proteins, ****genes**** coding therefor, transformed cells of organisms, methods and applications; Keiji Kondo, et al., 536/23.7; 435/69.1, 172.3, 252.3, 254.2, 254.21, 320.1; 536/24.1 [IMAGE AVAILABLE]

2. 5,468,485, Nov. 21, 1995, Avirulent microbes and uses therefor; Roy Curtiss, III, 424/184.1, 93.1, 93.2, 200.1; 435/69.1, 71.1, 172.1, 252.3, 252.33, 252.8 [IMAGE AVAILABLE]

3. 5,449,755, Sep. 12, 1995, Human cyclin E; James M. Roberts, et al., 530/350, 387.1; 536/23.5 [IMAGE AVAILABLE]

4. 5,427,785, Jun. 27, 1995, Rhizosheric bacteria; Clive W. Ronson, et al., 424/93.2; 47/57.6, 58; 71/7; 435/172.3, 252.2, 878; 935/64 [IMAGE AVAILABLE]

5. 5,408,037, Apr. 18, 1995, Methods for detecting glucagon antagonists; Robert A. Smith, et al., 530/308 [IMAGE AVAILABLE]

6. 5,369,766, Nov. 29, 1994, Object-oriented loader system with support for different load formats; Russell T. Nakano, et al., 395/700; 364/280, DIG.1 [IMAGE AVAILABLE]

7. 5,360,901, Nov. 1, 1994, ****Gene**** ****sequence**** encoding *Aspergillus niger* catalase-R; Randy M. Berka, et al., 536/23.2; 435/69.1, 71.1, 172.3, 192, 254.3, 320.1 [IMAGE AVAILABLE]

8. 5,360,732, Nov. 1, 1994, Production of *Aspergillus niger* catalase-R; Randy M. Berka, et al., 435/192, 69.1, 71.1, 172.3, 254.3, 320.1; 536/23.2; 935/14, 27, 36, 56, 68 [IMAGE AVAILABLE]

9. 5,350,671, Sep. 27, 1994, HCV immunoassays employing C domain antigens; Michael Houghton, et al., 435/5, 6, 975; 436/512, 518; 530/300, 326, 327, 328, 812, 826; 930/220, 223 [IMAGE AVAILABLE]

10. 5,349,127, Sep. 20, 1994, Expression of herbicide metabolizing cytochromes P450; Caroline Dean, et al., 800/205; 435/172.3, 320.1; 800/250, 255, DIG.71; 935/64, 67 [IMAGE AVAILABLE]

=> d 19 1 ab

US PAT NO: 5,470,971 [IMAGE AVAILABLE]

L9: 1 of 31

ABSTRACT:

****Genes**** (and portions thereof) which are stress-inducible, e.g., by cold-shock which encode useful proteins. The proteins contribute to confer thermo-tolerance and/or contribute to confer low temperature tolerance to organisms, like eucaryotes or procaryotes. Typical ****genes**** encoding such proteins and homologous ****genes**** encoding proteins with equivalent properties are discussed. Nucleotide ****sequences**** encoding such proteins, recombinant replicable expression vehicles which comprise ****DNA**** constructs which encode such proteins and competent transformed organisms like eucaryotes are discussed. The production of valuable fermentation products and of biologically active proteins under conditions outside the normal or optimum physiological growth conditions are described.

=> d 19 11-20

11. 5,324,659, Jun. 28, 1994, Yeast mutants useful for indentifying immunosupressants; Stephen A. Parent, et al., 435/255.2, 942 [IMAGE AVAILABLE]

12. 5,321,828, Jun. 14, 1994, High speed microcomputer in-circuit emulator; Michael D. Phillips, et al., 395/500; 364/232.3, 264.3, DIG.1 [IMAGE AVAILABLE]

13. 5,268,274, Dec. 7, 1993, Methods and nucleic acid ****sequences**** for the expression of the cellulose synthase operon; Arie Ben-Bassat, et al., 435/69.1, 101, 194, 252.3, 252.33, 320.1, 823; 536/23.2; 935/9, 14, 29, 40, 60, 72, 73 [IMAGE AVAILABLE]

14. 5,229,112, Jul. 20, 1993, Combatting plant insect pests with plant-colonizing microorganisms containing the toxin ****gene**** B. thuringiensis as a chromosomal insertion; Mark G. Obukowicz, et al., 424/93.2; 435/252.3, 252.34 [IMAGE AVAILABLE]

15. 5,212,296, May 18, 1993, Expression of herbicide metabolizing cytochromes; Caroline Dean, et al., 536/23.2; 424/93.2, 93.21; 435/320.1; 536/23.7 [IMAGE AVAILABLE]

16. 5,198,346, Mar. 30, 1993, Generation and selection of novel ****DNA****-binding proteins and polypeptides; Robert C. Ladner, et al., 435/69.1, 172.3, 252.3, 320.1 [IMAGE AVAILABLE]

17. 5,190,871, Mar. 2, 1993, Use of the site-specific integrating function of phage .phi.C31; Karen L. Cox, et al., 435/172.3, 252.35, 320.1 [IMAGE AVAILABLE]

18. 5,175,272, Dec. 29, 1992, ****DNA**** ****sequences**** with increased expression of HBcAg; Richard L. Mallonee, 536/26.3 [IMAGE AVAILABLE]

19. 5,175,094, Dec. 29, 1992, Increased expression of HBcAg; Richard L. Mallonee, 435/69.3 [IMAGE AVAILABLE]

20. 5,173,427, Dec. 22, 1992, Vectors and hosts with increased expression of HBCAG; Richard L. Mallonee, 435/252.33, 240.2, 252.3, 320.1 [IMAGE AVAILABLE]

=> s p4502c11

L10 0 P4502C11

=> s p4502c?

L11 0 P4502C?

=> s p450?

L12 114 P450?

=> s l12 and hypertension

7510 HYPERTENSION

L13 9 L12 AND HYPERTENSION

=> d l13 1-9

1. 5,294,725, Mar. 15, 1994, Scopularin; Donald R. Kirsch, et al., 549/417 [IMAGE AVAILABLE]

2. 5,288,721, Feb. 22, 1994, Substituted epoxyalkyl xanthines; J. Peter Klein, et al., 514/263; 544/267 [IMAGE AVAILABLE]

08/31/91 145

?b 154 55

18jan96 08:34:27 User208654 Session D59.1

\$0.12 0.004 Hrs File1

\$0.12 Estimated cost File1

\$0.12 Estimated cost this search

\$0.12 Estimated total session cost 0.004 Hrs.

SYSTEM:OS - DIALOG OneSearch

File 154:MEDLINE(R) 1985-1996/Jan W4

(c) format only 1996 Knight-Ridder Info

File 55:BIOSIS PREVIEWS(R) 1985-1995/Jan W1

(c) 1996 BIOSIS

Set Items Description

--- ---

?s pcr(5n)clon?

43619 PCR

253749 CLON?

S1 2678 PCR(5N)CLON?

?s pcr(3n)clon?

43619 PCR

253749 CLON?

S2 2010 PCR(3N)CLON?

?s s2 and review?

2010 S2

391802 REVIEW?

S3 8 S2 AND REVIEW?

?rd

...completed examining records

S4 6 RD (unique items)

?t s4/6/1-6

4/6/1 (Item 1 from file: 154)

09461990 95391990

Clonal relationship between lymphocytic predominance Hodgkin's disease and concurrent or subsequent large-cell lymphoma of B lineage.

4/6/2 (Item 2 from file: 154)

08946062 94261062

Isolated bone relapse during hematologic remission in childhood acute lymphoblastic leukemia: report of a metatarsal relapse and review of the literature.

4/6/3 (Item 3 from file: 154)

08336712 93046712

Protein engineering of antibodies.

4/6/4 (Item 4 from file: 154)

08135143 92273143

In vitro antibodies: strategies for production and application.

4/6/5 (Item 1 from file: 55)
11313321 BIOSIS Number: 97513321

The role of molecular genetics in field studies on malaria parasites
Print Number: Biological Abstracts Vol. 098 Iss. 011 Ref. 149060

4/6/6 (Item 2 from file: 55)
10804539 BIOSIS Number: 97004539

A family of serine protease genes expressed in adult buffalo fly
(*Haematobia irritans exigua*)

Print Number: Biological Abstracts Vol. 097 Iss. 001 Ref. 004116
?t s4/7/3-6

4/7/3 (Item 3 from file: 154)
DIALOG(R) File 154:MEDLINE(R)

(c) format only 1996 Knight-Ridder Info. All rts. reserv.

08336712 93046712

Protein engineering of antibodies.

Sandhu JS

Samuel Lunenfeld Research Institute, Mount Sinai Hospital, Toronto,
Ontario, Canada.

Crit Rev Biotechnol (UNITED STATES) 1992, 12 (5-6) p437-62, ISSN
0738-8551 Journal Code: CRB

Languages: ENGLISH

Document type: JOURNAL ARTICLE; REVIEW; REVIEW, TUTORIAL

This article reviews the technical advances in antibody engineering and the clinical applications of these molecules. Recombinant DNA technology facilitates the construction and expression of engineered antibodies. These novel molecules are designed to meet specific applications. Although genomic and cDNA cloning have been used widely in the past to isolate the relevant antibody V domains, at present, the PCR-based cloning is the preferred system. Bacterial and mammalian expression systems are used commonly for the production of antibodies, antibody fragments, and antibody fusion proteins. A range of chimeric antibodies with murine V domains joined to C regions from human and other species have been produced and found to exhibit the expected binding characteristics and effector functions. Humanized antibodies have been developed to minimize the HAMA response, and bifunctional immunoglobulins are being used in tumor therapy and diagnosis. Single chain antibodies and fusion proteins with antibody specificities jointed to nonimmunoglobulin sequences provide a source of antibody-like molecules with novel properties. The potential applications of minimal recognition units and antigenized antibodies are described. Combinatorial libraries produced in bacteriophage present an alternative to hybridomas for the production of antibodies with the desired antigen binding specificities. Future developments in this field are discussed also. (182 Refs.)

4/7/4 (Item 4 from file: 154)
DIALOG(R) File 154:MEDLINE(R)

(c) format only 1996 Knight-Ridder Info. All rts. reserv.

08135143 92273143

In vitro antibodies: strategies for production and application.

Morrison SL

Department of Microbiology and Molecular Genetics, University of California, Los Angeles 90024-1489.

Annu Rev Immunol (UNITED STATES) 1992, 10 p239-65, ISSN 0732-0582

Journal Code: ALO

Contract/Grant No.: CA 16858, CA, NCI; AI29470, AI, NIAID

Languages: ENGLISH

Document type: JOURNAL ARTICLE; REVIEW; REVIEW, ACADEMIC

The approaches to the production of antibodies (Ab) using the techniques of genetic engineering and expression are reviewed. Genetic engineering facilitates the production of proteins tailor-made for an intended use. Bacterial and mammalian expression systems are commonly used for the production of Ab and Ab-like molecules. While genomic or cDNA cloning can be used to obtain the relevant variable regions, PCR-based cloning approaches facilitate the acquisition of additional binding specificities. Large numbers of different chimeric Abs with murine variable regions joined to constant regions from human and other species have been expressed and found to exhibit the expected binding specificities and effector functions. These molecules have been used to study the structural basis of effector functions such as complement activation and Fc receptor binding, and potentially they may be used as therapeutic agents. Carbohydrate has been shown to influence both variable and constant region function. Single-chain Abs and fusion proteins with Ab binding specificities joined to nonimmunoglobulin sequences provide a source of Ab-like molecules with novel properties, and genetically engineered Ab-like molecules provide a source of useful antigens. Combinatorial libraries produced in bacteriophage present an alternative to hybridomas for the production of Abs with desired combining specificities. Issues of the immunogenicity of the recombinant molecules are addressed. (142 Refs.)

4/7/5 (Item 1 from file: 55)

DIALOG(R) File 55:BIOSIS PREVIEWS(R)

(c) 1996 BIOSIS. All rts. reserv.

11313321 BIOSIS Number: 97513321

The role of molecular genetics in field studies on malaria parasites

Walliker D

Div. Biol. Sci., Univ. Edinburgh, West Mains Rd., Edinburgh EH9 3JN, UK

International Journal for Parasitology 24 (6). 1994. 799-808.

Full Journal Title: International Journal for Parasitology

ISSN: 0020-7519

Language: ENGLISH

Print Number: Biological Abstracts Vol. 098 Iss. 011 Ref. 149060

Molecular genetics is having an important impact on the study of genes in natural populations of malaria parasites. The polymerase chain reaction (PCR) is proving particularly valuable for identifying genes in parasites taken directly from their hosts, without the need to establish them in culture. This is leading to novel methods of diagnosis, for example of drug-resistant parasites. Molecular techniques are also greatly assisting understanding of the genetic structure of parasite populations. This is relevant to the current debate on whether *Plasmodium falciparum* has a clonal or randomly interbreeding structure. Many patients are infected with mixtures of genetically distinct clones. PCR is being used to examine the genotypes of individual oocysts in the mosquito vector. In wild-caught mosquitoes in areas highly endemic for *P. falciparum*, a large proportion of oocysts are heterozygous, showing that cross-mating occurs frequently

between clones during mosquito feeds. In areas of lower endemicity, there is evidence of less frequent crossing.

4/7/6 (Item 2 from file: 55)
DIALOG(R)File 55:BIOSIS PREVIEWS(R)
(c) 1996 BIOSIS. All rts. reserv.

10804539 BIOSIS Number: 97004539

A family of serine protease genes expressed in adult buffalo fly
(*Haematobia irritans exigua*)

Elvin C M; Whan V; Riddles P W

CSIRO, Div. Tropical Animal Production, Lond Pocket Lab., Private Bag No
3 PO, Indooroopilly 4068, QLD, AUL

Molecular & General Genetics 240 (1). 1993. 132-139.

Full Journal Title: Molecular & General Genetics

ISSN: 0026-8925

Language: ENGLISH

Print Number: Biological Abstracts Vol. 097 Iss. 001 Ref. 004116

Gene fragments encoding serine proteases expressed in adult buffalo fly (*Haematobia irritans exigua*) were amplified from cDNA using generic oligonucleotide PCR primers, based on conserved residues surrounding the active-site His and Ser amino acids found in all serine proteases. The PCR product consisted of a broad band extending from about 450 bp to 520 bp, which suggested that the PCR product actually consisted of numerous DNA fragments of slightly variable sizes. Seventeen independent clones of these fragments, each with an insert of approximately 480 bp, were digested with HaeIII. Comparison of restriction fragment patterns indicated that 13 of these clones harboured different PCR products. This was confirmed by DNA sequence analysis of 9 clones. Each of the sequenced clones contained all open reading frame which included structurally conserved regions characteristic of the serine protease superfamily. This study reveals the expression of a large and highly variable repertoire of serine proteases in adult buffalo fly. Importantly, these data also demonstrate the utility of such an approach in obtaining DNA probes for use in further investigations of gene family organization and expression, as well as providing recombinant antigens in the form of fusion proteins which may be used as candidates for vaccine production.

?ds

Set	Items	Description
S1	2678	PCR (5N) CLON?
S2	2010	PCR (3N) CLON?
S3	8	S2 AND REVIEW?
S4	6	RD (unique items)

?s s1 and review?

	2678	S1
	391802	REVIEW?
S5	12	S1 AND REVIEW?

?s s5 not s3

	12	S5
	8	S3
S6	4	S5 NOT S3

?rd

...completed examining records
S7 3 RD (unique items)
?t s7/6/1-3

7/6/1 (Item 1 from file: 154)
09095781 95025781

[Low-grade MALT-type non-Hodgkin lymphoma of the stomach with local recurrence 14 years following resection. Demonstration of clonal identity using polymerase-chain-reaction (PCR)]

Niedrig malignes Non-Hodgkin-Lymphom vom MALT-Typ im Magen mit Lokalrezidiv 14 Jahre nach Resektion. Demonstration der klonalen Identität mittels Polymerase-Kettenreaktion (PCR).

7/6/2 (Item 2 from file: 154)
09081894 95011894

Utility of microsporidian rRNA in diagnosis and phylogeny: a review.

7/6/3 (Item 3 from file: 154)
08677862 93387862

Gene transfer and cardiovascular disorders.

?t s7/7/3

7/7/3 (Item 3 from file: 154)
DIALOG(R) File 154:MEDLINE(R)

(c) format only 1996 Knight-Ridder Info. All rts. reserv.

08677862 93387862

Gene transfer and cardiovascular disorders.

French BA

Department of Medicine, Baylor College of Medicine, Houston, Texas.

Herz (GERMANY) Aug 1993, 18 (4) p222-9, ISSN 0340-9937

Journal Code: F88

Contract/Grant No.: P50-HL42267-01, HL, NHLBI

Languages: ENGLISH

Document type: JOURNAL ARTICLE; REVIEW; REVIEW, TUTORIAL

Within the past four years, basic recombinant techniques (such as molecular cloning, sequencing, site-directed mutagenesis, PCR, and transfection) have been combined to yield a "second generation" of recombinant DNA technology with experimental potential which could barely have been envisioned only a decade ago. This review will focus upon the genesis and cardiovascular application of two recent developments in gene transfer technology: gene targeting by homologous recombination and direct in vivo gene transfer. Gene targeting evolved from transgenic mouse technology but is distinguished by its ability to precisely disrupt or "knock-out" specific genes in the murine genome. This not only provides decisive answers to functional questions, but also produces accurate models of human genetic disorders. In vivo gene transfer provides for the direct introduction of genetic information into living tissues. In vivo gene transfer not only facilitates basic research by providing a simple and direct way to analyze gene structure and function in intact animals, but may also find direct clinical application in the treatment of genetic and acquired disorders such as familial hypercholesterolemia and restenosis. (34 Refs.)

?s pcr-based cloning

S8 0 PCR-BASED CLONING
?s pcr(w)based(w)cloning

43619 PCR
378332 BASED
98200 CLONING

S9 57 PCR(W)BASED(W)CLONING
?rd

...examined 50 records (50)
...completed examining records
S10 33 RD (unique items)
?s s10 and review?

33 S10
391802 REVIEW?

S11 2 S10 AND REVIEW?
?t s11/6/1-2

11/6/1 (Item 1 from file: 154)
08336712 93046712
Protein engineering of antibodies.

11/6/2 (Item 2 from file: 154)
08135143 92273143
In vitro antibodies: strategies for production and application.
?t s10/6/1-33

10/6/1 (Item 1 from file: 154)
09440942 95370942
Cloning of rat interleukin-3 receptor beta-subunit from cultured
microglia and its mRNA expression in vivo.

10/6/2 (Item 2 from file: 154)
09437008 95367008
Molecular characterization of the murine syk protein tyrosine kinase
cDNA, transcripts and protein.

10/6/3 (Item 3 from file: 154)
09343326 95273326
Induced oleoresin biosynthesis in grand fir as a defense against bark
beetles.

10/6/4 (Item 4 from file: 154)
09291530 95221530
Cloning and subcellular localization of novel rab proteins reveals
polarized and cell type-specific expression.

10/6/5 (Item 5 from file: 154)
09274052 95204052

Receptor tyrosine kinases expressed in metastatic colon cancer.

10/6/6 (Item 6 from file: 154)
09245019 95175019

Expression of cyclic nucleotide-gated cation channels in non-sensory tissues and cells.

10/6/7 (Item 7 from file: 154)
09214459 95144459

PCR-based cloning, sequencing, and exon mapping of lymphocyte-derived neuroendocrine peptides.

10/6/8 (Item 8 from file: 154)
09210431 95140431

Enhanced expression of multiple protein tyrosine phosphatases in the regenerating mouse liver: isolation of PTP-RL10, a novel cytoplasmic-type phosphatase with sequence homology to cytoskeletal protein 4.1.

10/6/9 (Item 9 from file: 154)
09207784 95137784

HLA class I allele (HLA-A2) expression defect associated with a mutation in its enhancer B inverted CAT box in two families.

10/6/10 (Item 10 from file: 154)
09115227 95045227

The two nonallelic insulin-like growth factor-I genes in *Xenopus laevis* are differentially regulated during development.

10/6/11 (Item 11 from file: 154)
09064984 94379984

PCR-based cloning of the full-length *Neurospora* eukaryotic initiation factor 5A cDNA: polyhistidine-tagging and overexpression for protein affinity binding.

10/6/12 (Item 12 from file: 154)
08953853 94268853

PISSLRE, a human novel CDC2-related protein kinase.

10/6/13 (Item 13 from file: 154)
08905079 94220079

Isolation of three novel POU-domain containing cDNA clones from lactating mouse mammary gland.

10/6/14 (Item 14 from file: 154)
08888677 94203677

Expression of two novel eph-related receptor protein tyrosine kinases in mammary gland development and carcinogenesis [published erratum appears in *Oncogene* 1994 Aug;9(8):2431]

10/6/15 (Item 15 from file: 154)

08866267 94181267

Molecular cloning of a novel non-receptor tyrosine kinase, HYL (hematopoietic consensus tyrosine-lacking kinase).

10/6/16 (Item 16 from file: 154)

08841206 94156206

Characterization of a novel murine testis-specific serine/threonine kinase.

10/6/17 (Item 17 from file: 154)

08782361 94097361

A single amino acid substitution in the H-2Kb molecule generates a defined allogeneic epitope.

10/6/18 (Item 18 from file: 154)

08754961 94069961

The maize stripe virus major noncapsid protein messenger RNA transcripts contain heterogeneous leader sequences at their 5' termini.

10/6/19 (Item 19 from file: 154)

08534794 93244794

The organization of the intron-containing human S6 ribosomal protein (rpS6) gene and determination of its location at chromosome 9p21.

10/6/20 (Item 20 from file: 154)

08452680 93162680

The structure of the human intron-containing S8 ribosomal protein gene and determination of its chromosomal location at 1p32-p34.1.

10/6/21 (Item 21 from file: 154)

08373979 93083979

A small plasmid for recombination-based screening.

10/6/22 (Item 22 from file: 154)

08336712 93046712

Protein engineering of antibodies.

10/6/23 (Item 23 from file: 154)

08241086 92379086

Human inter-alpha-trypsin inhibitor: full-length cDNA sequence of the heavy chain H1.

10/6/24 (Item 24 from file: 154)

08155162 92293162

PCR based cloning and sequencing of isogenes encoding the tree pollen major allergen Car b I from *Carpinus betulus*, hornbeam.

10/6/25 (Item 25 from file: 154)

08135143 92273143

In vitro antibodies: strategies for production and application.

10/6/26 (Item 26 from file: 154)

08072010 92210010

The complexity of the Rab and Rho GTP-binding protein subfamilies revealed by a PCR cloning approach.

10/6/27 (Item 27 from file: 154)

07821705 91340705

Isolation of a cDNA encoding a mammalian multiubiquitinating enzyme (E225K) and overexpression of the functional enzyme in Escherichia coli.

10/6/28 (Item 1 from file: 55)

11965316 BIOSIS Number: 98565316

Protein tyrosine kinase expression during the estrous cycle and carcinogenesis of the mammary gland

Print Number: Biological Abstracts Vol. 101 Iss. 001 Ref. 008121

10/6/29 (Item 2 from file: 55)

11930714 BIOSIS Number: 98530714

Three new putative voltage gated sodium channels (VGSC) from human neuroblastoma (SH-SY5Y) cells: A PCR based cloning strategy

Print Number: Biological Abstracts/RRM Vol. 047 Iss. 012 Ref. 200589

10/6/30 (Item 3 from file: 55)

11470290 BIOSIS Number: 98070290

Identification of new genes expressed in hemopoietic cell lines: Yield of three PCR-based cloning approaches

Print Number: Biological Abstracts/RRM Vol. 047 Iss. 002 Ref. 031877

10/6/31 (Item 4 from file: 55)

11236030 BIOSIS Number: 97436030

PCR-based cloning and sequence analysis of fixL homologue from Frankia species strain CeS15

Print Number: Biological Abstracts/RRM Vol. 046 Iss. 010 Ref. 160794

10/6/32 (Item 5 from file: 55)

10038487 BIOSIS Number: 95038487

MOLECULAR CLONING AND CHARACTERIZATION OF RKLK10 A CDNA ENCODING T KININOGENASE FROM RAT SUBMANDIBULAR GLAND AND KIDNEY

10/6/33 (Item 6 from file: 55)

7883667 BIOSIS Number: 40084667

PCR-BASED CLONING STRATEGY FOR G PROTEIN-COUPLED RECEPTORS

?t s10/7/4 7-9 11 15 17 24 26 27 30-33

10/7/4 (Item 4 from file: 154)

DIALOG(R)File 154:MEDLINE(R)

(c) format only 1996 Knight-Ridder Info. All rts. reserv.

09291530 95221530

Cloning and subcellular localization of novel rab proteins reveals polarized and cell type-specific expression.

Lutcke A; Parton RG; Murphy C; Olkkonen VM; Dupree P; Valencia A; Simons K; Zerial M

European Molecular Biology Laboratory, Heidelberg, FRG.

J Cell Sci (ENGLAND) Dec 1994, 107 (Pt 12) p3437-48, ISSN 0021-9533
Journal Code: HNK

Languages: ENGLISH

Document type: JOURNAL ARTICLE

Small GTPases of the rab subfamily are specific regulators of vesicular transport. The intracellular localization of these proteins has been mostly investigated in cultured cells where they have been found associated with distinct compartments of the exocytic and endocytic pathways. Using a PCR-based cloning approach we have recently identified several novel rab proteins, extending the total number of this family to more than 30 members. Here, we have investigated the mRNA expression in different tissues and the intracellular localization in organ cryosections of two rab proteins, rab18 and rab20. Both northern blot analysis and confocal immunofluorescence microscopy demonstrated that these proteins are expressed in a tissue- and cell type-dependent manner. Despite their presence in non-polarized cells and polarized cells, both proteins are highly expressed on the apical side of kidney tubule epithelial cells. Electron microscopic studies revealed that rab18 and rab20 are located in apical dense tubules, endocytic structures underlying the apical plasma membrane, suggesting that they play a role in apical endocytosis/recycling. In intestinal epithelial cells as well, both proteins were localized apically, but, in addition, rab18 was found associated with the basolateral domain, suggesting that this protein is not restricted to the apical transport machinery of polarized epithelial cells. The results demonstrate that, depending on the epithelial cell type, rab proteins that are also expressed in non-polarized cells may be enriched in one or both surface domains. Together with the observed tissue- and cell type-dependent variation in the expression of the rab proteins, this suggests that the large number of mammalian rab proteins might reflect the specific requirements in the organization of membrane traffic encountered by different cell types.

10/7/7 (Item 7 from file: 154)

DIALOG(R)File 154:MEDLINE(R)

(c) format only 1996 Knight-Ridder Info. All rts. reserv.

09214459 95144459

PCR-based cloning, sequencing, and exon mapping of lymphocyte-derived neuroendocrine peptides.

Maier CC; Blalock JE

Department of Physiology and Biophysics, University of Alabama at Birmingham 35294.

Immunomethods (UNITED STATES) Aug 1994, 5 (1) p3-7, ISSN 1058-6687
Journal Code: B3R

Contract/Grant No.: P01 NS29719, NS, NINDS; DK38024, DK, NIDDK; T3207335

Languages: ENGLISH

Document type: JOURNAL ARTICLE

In this report a procedure for the analysis of mRNA expression in cells of limited availability by the reverse transcriptase-polymerase chain reaction (RT-PCR) method is described. The cells are lysed with Nonidet P-40, and the mRNA in the lysate is used directly as template for the cDNA synthesis reaction. Target cDNA is then amplified by PCR, and the products can be analyzed that same day by agarose gel electrophoresis. The oligonucleotide primers used for amplification are designed to include restriction sites to facilitate cloning for subsequent sequencing. We have demonstrated that luteinizing hormone-releasing hormone mRNA can be amplified from the hypothalamus and thymus of a 7-day rat pup, in which the starting cell number was limited. Furthermore, exon usage by target cDNA in different cell types can be easily determined by amplifying with exon-specific primers. Proopiomelanocortin (POMC) mRNA expressed in the pituitary utilizes all three exons, while a majority of POMC mRNA expressed in lymphocytes lacks exons 1 and 2. Thus, this provides an extremely rapid and sensitive means not only for analyzing mRNA expression but also for differential exon usage.

10/7/8 (Item 8 from file: 154)

DIALOG(R)File 154:MEDLINE(R)

(c) format only 1996 Knight-Ridder Info. All rts. reserv.

09210431 95140431

Enhanced expression of multiple protein tyrosine phosphatases in the regenerating mouse liver: isolation of PTP-RL10, a novel cytoplasmic-type phosphatase with sequence homology to cytoskeletal protein 4.1.

Higashitsuji H; Arii S; Furutani M; Imamura M; Kaneko Y; Takenawa J; Nakayama H; Fujita J

First Department of Surgery, Faculty of Medicine, Kyoto University, Japan.

Oncogene (ENGLAND) Jan 19 1995, 10 (2) p407-14, ISSN 0950-9232

Journal Code: ONC

Languages: ENGLISH

Document type: JOURNAL ARTICLE

To elucidate the role that protein tyrosine phosphatase (PTPs) may play in liver regeneration, PTPs expressed in the mouse liver after partial hepatectomy (PH) were investigated by a PCR-based cloning method. Sequencing of 115 cDNA clones identified 10 different sequences including MPTP (T cell PTP), PTP-1B, PTP-P19, mR-PTP mu, R-PTP alpha, PTP NE-3 (PTP-P1), R-PTP-kappa and the murine homologue of human LAR. The remaining two sequences, PTP-RL9 and PTP-RL10, encoded novel PTPs. PTP-RL10 cDNA contained an open reading frame of 1176 amino acids with no apparent membrane-spanning region. The amino-terminal region had sequence homology to those of human erythrocyte protein 4.1 and ezrin, cytoskeletal proteins. In the regenerating liver, the levels of five PTP gene mRNAs (MPTP, PTP-P19, R-PTP alpha, LAR homologue, and PTP-RL9) increased within 6 h, decreased to the normal level by 24 h, and increased again at 48 to 72 h after PH. The levels of PTP-1B and R-PTP-kappa mRNAs peaked within 6 h, decreased gradually, and returned to the normal level by 168 h after PH. In contrast, the levels of two PTP mRNAs (mR-PTP mu and PTP-RL10) peaked at 48 to 72 h, and returned to the normal level by 168 h after PH. No expression of PTP NE-3 was detected in the liver by Northern blotting. The differential expression of multiple PTPs during the pre-replicative and post-replicative stages of liver regeneration suggests that PTPs are involved in the regulation of growth and differentiation of liver cells.

10/7/9 (Item 9 from file: 154)
DIALOG(R) File 154:MEDLINE(R)
(c) format only 1996 Knight-Ridder Info. All rts. reserv.

09207784 95137784

HLA class I allele (HLA-A2) expression defect associated with a mutation in its enhancer B inverted CAT box in two families.

Balas A; Garcia-Sanchez F; Gomez-Reino F; Vicario JL

Laboratory of Histocompatibility, Regional Transfusion Center, Madrid, Spain.

Hum Immunol (UNITED STATES) Sep 1994, 41 (1) p69-73, ISSN 0198-8859
Journal Code: G9W

Languages: ENGLISH

Document type: JOURNAL ARTICLE

The present study shows a very highly diminished HLA-A2 cell surface expression with mendelian segregation in two nonrelated Spanish families. The associated haplotype included Cblank-B38-DRB1*1301-DQ6 in both families. cDNA sequence analysis in two members, one of each pedigree, revealed the presence of the commonest HLA-A2 allele (A*0201), without repetitive mutations that could indicate inappropriate or inefficient translation. Further, the coamplified 3'-untranslated region sequence was also the same described for HLA-A2. HLA-A transcription frequency by means of cDNA PCR-based cloning experiments and by Northern blotting pointed out a relatively low number of HLA-A2 mRNA molecules compared with the complementary HLA-A allele. 5'-Regulatory region sequences from two low-expressing HLA-A2 nonrelated individuals showed a unique and identical single point mutation at position -101 (T to C), when compared with all MHC class I alleles sequenced so far. Position -101 is located in the inverted CAT box associated with the MHC class I enhancer B. The fact that this is an extremely well-conserved position leads us to postulate that this change may be the only responsible for the defective expression of HLA-A2.

10/7/11 (Item 11 from file: 154)
DIALOG(R) File 154:MEDLINE(R)
(c) format only 1996 Knight-Ridder Info. All rts. reserv.

09064984 94379984

PCR-based cloning of the full-length Neurospora eukaryotic initiation factor 5A cDNA: polyhistidine-tagging and overexpression for protein affinity binding.

Tao Y; Chen KY

Department of Chemistry, Rutgers University, Piscataway, NJ 08855-0939.

Biochem J (ENGLAND) Sep 1 1994, 302 (Pt 2) p517-25, ISSN 0264-6021

Journal Code: 9YO

Contract/Grant No.: R01 CA49695, CA, NCI

Languages: ENGLISH

Document type: JOURNAL ARTICLE

Eukaryotic initiation factor 5A (eIF-5A) is the only cellular protein known to contain a hypusine residue that is formed by transferring the aminobutyl moiety from spermidine to a specific lysine residue, followed by hydroxylation at the aminobutyl group. A simple PCR-based strategy was developed to obtain a full-length cDNA of Neurospora crassa eIF-5A. The strategy consists of (i) the design of a pair of key primers (21-mer) based on the highly conserved eIF-5A cDNA domains known in other species, (ii) PCR amplification of Neurospora cDNA using the two key primers to obtain the core sequence for the design of core primers, and (iii) combined use of the key primers, core primers and the universal primers, T3 and T7, to

amplify the target sequence in a *Neurospora* cDNA library. The longest cDNA obtained was cloned into pBlueScript phagemid, and sequence analysis indicated that it encodes a polypeptide of 163 amino acid residues with a codon usage preference characteristic of abundant *Neurospora* genes. The *Neurospora* polypeptide showed 59% and 67% identity with human and yeast eIF-5A precursor protein respectively. We subcloned the *Neurospora* eIF-5A cDNA into pQE-30, which introduces six adjacent histidine residues to the N-terminus of the recombinant protein. The resulting plasmid, pQTy21, was overexpressed in *Escherichia coli*, and the soluble polyhistidine-tagged protein was purified by metal chelation chromatography. We obtained about 60 mg of purified eIF-5A precursor from 1 litre of culture in a single step using a Ni(II)-nitrilotriacetic acid (NTA)-agarose column. The histidine-tagged eIF-5A precursor protein could be recognized by anti-*Neurospora crassa* 21 kDa protein serum raised against wild-type eIF-5A precursor and could serve as the substrate protein for deoxyhypusine synthase. Using the histidine-tagged recombinant protein and the Ni(II)-NTA-agarose column, we constructed a protein affinity column and demonstrated an affinity binding between eIF-5A precursor and deoxyhypusine synthase in the presence of NAD⁺. One-step eIF-5A precursor affinity-column chromatography could lead to a 30-fold purification of deoxyhypusine synthase.

10/7/15 (Item 15 from file: 154)

DIALOG(R) File 154:MEDLINE(R)

(c) format only 1996 Knight-Ridder Info. All rts. reserv.

08866267 94181267

Molecular cloning of a novel non-receptor tyrosine kinase, HYL (hematopoietic consensus tyrosine-lacking kinase).

Sakano S; Iwama A; Inazawa J; Ariyama T; Ohno M; Suda T

Department of Cell Differentiation, Kumamoto University School of Medicine, Japan.

Oncogene (ENGLAND) Apr 1994, 9 (4) p1155-61, ISSN 0950-9232

Journal Code: ONC

Languages: ENGLISH

Document type: JOURNAL ARTICLE

We identified a novel non-receptor tyrosine kinase from a human megakaryoblastic cell line, UT-7, by means of a PCR-based cloning method. The HYL gene contained a SH2 and SH3 domain and a tyrosine kinase catalytic domain. The deduced amino acid sequence of the protein encoded by this gene was most homologous to CSK (c-src kinase). This gene and CSK shared some unique structural properties such as the absence of a myristylation signal and phosphorylation sites of tyrosine residues corresponding to tyrosines 416 and 527 of chicken p60c-src. Unlike CSK, the SH3 domain of HYL was unique since the ALYDY motif was absent. Northern blot analysis revealed a 2.2 kb transcript in various myeloid cell lines but not in adult tissues except for the brain and the lung, whereas CSK mRNA was ubiquitously expressed. The expression of HYL was upregulated when these myeloid cells were differentiated by induction with phorbol myristate acetate. We named this gene, hematopoietic consensus tyrosine-lacking kinase, HYL. The HYL gene was assigned to chromosome 19 at band p13. It is suggested that HYL plays a significant role in the signal transduction of hematopoietic cells.

10/7/17 (Item 17 from file: 154)

DIALOG(R) File 154:MEDLINE(R)

(c) format only 1996 Knight-Ridder Info. All rts. reserv.

08782361 94097361

A single amino acid substitution in the H-2Kb molecule generates a defined allogeneic epitope.

Kesari KV; Van Bleek G; Nathenson SG; Geliebter J

Howard Hughes Medical Institute, Rockefeller University, New York, NY 10021.

Mol Immunol (ENGLAND) Dec 1993, 30 (18) p1671-7, ISSN 0161-5890

Journal Code: NG1

Contract/Grant No.: AI07289, AI, NIAID; AI10702, AI, NIAID

Languages: ENGLISH

Document type: JOURNAL ARTICLE

Using Mitomycin C mutagenesis and negative and positive selection with monoclonal antibodies specific for H-2Kb and H-2Kbm10, respectively, a mutant cell line clone, Mitc-182, was isolated. Direct sequencing of uncloned cDNA as well as PCR based cloning and sequencing of the H-2Kb182 transcript from this mutant revealed a single G-->T transversion resulting in the substitution of Trp167 by cysteine. Serologically, the mutant Kb182 and Kbm10 are almost identical as each has lost at least five Kb specific mAb epitopes and gained several new epitopes. Interestingly, the mutant cell line, Mitc-182, is efficiently recognized by alloreactive CTLs raised in reciprocal combinations, e.g. CB6 anti Cbm10 and Cbm10 anti CB6, indicating that Kb182 contains both Kb and Kbm10 specific epitopes. The mutation has not affected the ability of Kb182 to present Kb restricted antigenic peptides of Sendai and vesicular stomatitis viruses. In addition to underscoring the importance of amino acid residue 167 in alloreactivity, these results indicate a positive correlation between the gain of both an mAb epitope and a defined alloreactive CTL epitope.

10/7/24 (Item 24 from file: 154)

DIALOG(R) File 154:MEDLINE(R)

(c) format only 1996 Knight-Ridder Info. All rts. reserv.

08155162 92293162

PCR based cloning and sequencing of isogenes encoding the tree pollen major allergen Car b I from Carpinus betulus, hornbeam.

Larsen JN; Stroman P; Ipsen H

ALK Research, Horsholm, Denmark.

Mol Immunol (ENGLAND) Jun 1992, 29 (6) p703-11, ISSN 0161-5890

Journal Code: NG1

Languages: ENGLISH

Document type: JOURNAL ARTICLE

Cloning of the gene encoding the major allergen, Car b I, from Carpinus betulus (hornbeam) pollen was performed using the Polymerase Chain Reaction (PCR) to specifically amplify the gene of interest using single stranded cDNA as template. Specific primers, deduced from the aminoterminal sequence of the purified protein, were tailored to facilitate direct expression of plasmic clones, and the large fraction of positive clones obtained, revealed the presence of isogenic variation. Three clones were characterized in detail by antibody based assays and nucleotide sequencing. The recombinant allergens were shown by crossed immunoelectrophoresis (CIE) to precipitate with monospecific polyclonal rabbit antibodies raised against purified Bet v I, by crossed radioimmuno-electrophoresis (CRIE) to bind tree pollen allergic patient serum IgE, and by immunoblotting to bind murine monoclonal antibodies, raised against purified Car b I from pollen. Car b I is encoded by a 159-triplets open reading frame. The molecular masses (M(r) = 17272, 17355 and 17217 Da, respectively), the amino acid

composition, and the aminoterminal sequence of the predicted polypeptides agree well with data obtained by analysis of the protein purified from pollen. The deduced amino acid sequences show pronounced homology (73, 75 and 74% identities respectively) to Bet v I, the major allergen from *Betula verrucosa* (white birch) pollen. Soluble recombinant Car b I, without a fusion partner, was produced in *Escherichia coli* with an immunochemical reactivity closely resembling that of the native pollen allergen. The tree pollen major allergens therefore constitute an ideal system for the study of allergenic epitopes.

10/7/26 (Item 26 from file: 154)

DIALOG(R) File 154:MEDLINE(R)

(c) format only 1996 Knight-Ridder Info. All rts. reserv.

08072010 92210010

The complexity of the Rab and Rho GTP-binding protein subfamilies revealed by a PCR cloning approach.

Chayrier P; Simons K; Zerial M

Cell Biology Programme, European Molecular Biology Laboratory, Heidelberg, F.R.G.

Gene (NETHERLANDS) Mar 15 1992, 112 (2) p261-4, ISSN 0378-1119

Journal Code: FOP

Languages: ENGLISH

Document type: JOURNAL ARTICLE

Partial sequences corresponding to eleven novel Rab proteins and one new Rho protein have been isolated using a PCR-based cloning approach. These results confirm that the overall diversity of the Rab and Rho protein subfamilies account for more than thirty different members in mammalian cells.

10/7/27 (Item 27 from file: 154)

DIALOG(R) File 154:MEDLINE(R)

(c) format only 1996 Knight-Ridder Info. All rts. reserv.

07821705 91340705

Isolation of a cDNA encoding a mammalian multiubiquitinating enzyme (E225K) and overexpression of the functional enzyme in *Escherichia coli*.

Chen ZJ; Niles EG; Pickart CM

Department of Biochemistry, State University of New York, Buffalo 14214.

J Biol Chem (UNITED STATES) Aug 25 1991, 266 (24) p15698-704, ISSN

0021-9258 Journal Code: HIV

Contract/Grant No.: AI28824, AI, NIAID

Languages: ENGLISH

Document type: JOURNAL ARTICLE

The ubiquitin (Ub)-conjugating enzyme E2(25K) catalyzes the synthesis of multi-Ub chains in which successive Ub units are linked by an isopeptide bond involving the epsilon-amino group of Lys-48 of Ubn, and the COOH-terminal Gly residue of Ubn+1 (Chen, Z., and Pickart, C. M. (1990) J. Biol. Chem., 265, 21835-21842). We now describe the polymerase chain reaction (PCR)-based cloning of an E2(25K)-encoding cDNA from a bovine thymus library, using degenerate oligonucleotide primers based on the sequences of two E2(25K) peptides. The cDNA encodes a 200-residue protein whose sequence bears similarities of 66 and 59%, respectively, to the sequences of the Ub-conjugating enzymes encoded by the UBC1 and UBC4/UBC5 genes of the yeast *Saccharomyces cerevisiae*. These three yeast E2s play key roles in Ub-dependent proteolysis (Seufert, W., McGrath, J. P., and

Jentsch, S. (1990) EMBO J. 9, 4535-4541). Comparison of the amino acid sequence of E2(25K) with other known E2 sequences strongly suggests that Cys-92, one of two E2(25K) Cys residues, forms the Ub thiol ester adduct that is an intermediate in E2-catalyzed multiubiquitination. The E2(25K)-encoding cDNA was overexpressed in Escherichia coli, and the recombinant E2(25K) protein was purified to electrophoretic homogeneity; enzymatic assays showed that its multiubiquitinating activity was quantitatively identical with that of the native protein. The availability of a cloned cDNA will allow us to assess the physiological role of E2(25K).

10/7/30 (Item 3 from file: 55)
DIALOG(R)File 55:BIOSIS PREVIEWS(R)
(c) 1996 BIOSIS. All rts. reserv.

11470290 BIOSIS Number: 98070290
Identification of new genes expressed in hemopoietic cell lines: Yield of three PCR-based cloning approaches
Furukawa T; Nakamoto B; Papayannopoulou T; Stamatoyannopoulos G
Div. Hematol., Univ. Wash., Seattle, WA, USA
Blood 84 (10 SUPPL. 1). 1994. 414A.
Full Journal Title: Abstracts Submitted to the 36th Annual Meeting of the American Society of Hematology, Nashville, Tennessee, USA, December 2-6, 1994. Blood
ISSN: 0006-4971
Language: ENGLISH
Print Number: Biological Abstracts/RRM Vol. 047 Iss. 002 Ref. 031877

10/7/31 (Item 4 from file: 55)
DIALOG(R)File 55:BIOSIS PREVIEWS(R)
(c) 1996 BIOSIS. All rts. reserv.

11236030 BIOSIS Number: 97436030
PCR-based cloning and sequence analysis of fixL homologue from Frankia species strain CeS15
Murry M A; Stigter J; De Bruijn F J
Dep. Bot. and Plant Pathol., Mich. State Univ., East Lansing, MI 48824, USA
0 (0). 1993. 485.
Full Journal Title: Palacios, R., J. Mora and W. E. Newton (Ed.). Current Plant Science and Biotechnology in Agriculture, Vol. 17. New horizons in nitrogen fixation; 9th International Congress on Nitrogen Fixation, Cancun, Mexico, December 6-12, 1992. xvi+788p. Kluwer Academic Publishers: Dordrecht, Netherlands; Norwell, Massachusetts, USA. ISBN 0-7923-2207-X.
ISSN: *****
Language: ENGLISH
Print Number: Biological Abstracts/RRM Vol. 046 Iss. 010 Ref. 160794

10/7/32 (Item 5 from file: 55)
DIALOG(R)File 55:BIOSIS PREVIEWS(R)
(c) 1996 BIOSIS. All rts. reserv.

10038487 BIOSIS Number: 95038487
MOLECULAR CLONING AND CHARACTERIZATION OF RKLK10 A CDNA ENCODING T KININOGENASE FROM RAT SUBMANDIBULAR GLAND AND KIDNEY